

A man with short dark hair and a goatee, wearing a black shirt, is sitting in a black office chair. He is looking towards the camera with a slight smile. His hands are clasped together on a desk in front of him. The background is slightly blurred, showing what appears to be an office setting with a desk and some papers. Overlaid on the image is Arabic text in a white, stylized font. At the bottom left, there are two small blue heart icons.

اللهم أنر قبر من حنّ له القلب
اللهم هب له سعة في قبره لا يراها نهاية
هب لمضجعه طيباً و لظلمته نوراً
ولذنوبه غفراناً و برّد قبره
وزده احساناً فوق احسانه
واجعل الجنة مسكنه

Back to Filters

Question 1 of 203

Which of the following renal pathologies is most likely to recur in a renal transplant patient?

- | | |
|---|--|
| A | Diabetic renal disease |
| B | Focal segmental glomerulosclerosis |
| C | Minimal change glomerulonephritis |
| D | Membranous glomerulonephritis |
| E | Membranoproliferative glomerulonephritis |

2211

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Question 1 of 203

A	Diabetic renal disease
B	Focal segmental glomerulosclerosis
C	Minimal change glomerulonephritis
D	Membranous glomerulonephritis
E	Membranoproliferative glomerulonephritis

Membranoproliferative glomerulonephritis is associated with a 30-90% recurrence rate post renal transplant, (Type 2 much greater than Type 1). Whilst the recurrence rate is also high with FSGS, (40%), it does not quite approach these levels. For membranous glomerulonephritis, recurrence rates from case series approach 30%, minimal change disease is not usually a cause of renal transplant, and diabetic renal disease takes years to appear in a transplanted kidney.

Rate this question:      

Next Question

Previous Question	Tag Question
Feedback	End Session

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	1
Responses Total:	1
Responses - % Correct:	0%

Question 2 of 203



Creatinine	250 μ mol/l
Hb	14 g/dl
Phosphate	2.0 mmol/l
Calcium	2.1 mmol/l

A	Adult polycystic kidney disease
B	Contrast nephropathy
C	Diabetes
D	Diffuse proliferative glomerulonephritis
E	Hypertensive nephropathy

Submit

[Previous Question](#)
[Skip Question](#)

Question 2 of 203

Creatinine	250 μmol/l
Hb	14 g/dl
Phosphate	2.0 mmol/l
Calcium	2.1 mmol/l

A	Adult polycystic kidney disease
B	Contrast nephropathy
C	Diabetes
D	Diffuse proliferative glomerulonephritis
E	Hypertensive nephropathy

© 2005 Blackwell Publishing Ltd

Rate this question: 

End Session

Peer Responses %

Session Progress

Responses incorrect. 2

Responses - % Correct: 0%

[Blog](#) [About Pastest](#) [Contact Us](#) [Help](#)

© Pastest 2017

Back to Filters

Question 3 of 203

A patient presents with a history of poor urine output over the past 2 days. Other than diarrhoea about a week ago he has no other significant history.

Which of the following tests would be useful for establishing the cause of his acute renal failure?

- A Potassium measurement
- B Stool culture
- C Clotting screen
- D Blood film
- E AXR

2215

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Question 3 of 203

Which of the following tests would be useful for establishing the cause of his acute renal failure?

A	Potassium measurement
B	Stool culture
C	Clotting screen
D	Blood film
E	AXR

You should suspect a diagnosis of haemolytic-uraemic syndrome associated with diarrhoea caused by *Escherichia coli*. The potassium concentration may well be high in cases of both acute renal failure (ARF) and haemolysis, but it will not tell you the diagnosis. The clotting screen will be normal unless the patient becomes so unwell they show a disseminated intravascular coagulation (DIC) picture.

Rate this question:

End Session

Peer Responses %

Responses Correct:	0
Responses Incorrect:	3
Responses Total:	3
Responses - % Correct:	0%

Back to Filters

Question 4 of 203

A 40-year-old man is referred by his GP to A&E with severe acute left-sided flank pain. This is the first time he has had such severe sudden pain. He has no past medical history of cardiovascular or renal disease. On examination his BMI is 30 kg/m² and pulse 78 bpm and regular. His jugular venous pressure is not raised, heart sounds are normal and his chest is clear. His liver is not palpable but he has a mass in the left flank. The mass is bimanually palpable and you can get above it.

Investigations are as follows:

Na ⁺	140 mmol/l
K ⁺	4.2 mmol/l
Urea	10.2 mmol/l
Creatinine	203 mmol/l
Hb	15.0 g/dl
WCC	5.1 × 10 ⁹ /l
MCV	81 fl
PLT	243 × 10 ⁹ /l
ESR	8 mm/h
Urine dipstick	blood +++, protein +

What other disorder is this patient likely to suffer from?

- A

Aortic valve prolapse
- B

Berry aneurysms in 50–75%
- C

Hypertension
- D

Gallstones
- E

Diabetes mellitus

2523

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 4 of 203

Investigations are as follows:

Na ⁺	140 mmol/l
K ⁺	4.2 mmol/l
Urea	10.2 mmol/l
Creatinine	203 mmol/l
Hb	15.0 g/dl
WCC	5.1×10^9 /l
MCV	81 fl
PLT	243×10^9 /l
ESR	8 mm/h
Urine dipstick	blood +++, protein +

- | | |
|---|---------------------------|
| A | Aortic valve prolapse |
| B | Berry aneurysms in 50-75% |
| C | Hypertension |
| D | Gallstones |
| E | Diabetes mellitus |

- | | |
|---|--------------|
| C | Hypertension |
|---|--------------|

A	Aortic valve prolapse
---	-----------------------

B	Berry aneurysms in 50-75%
---	---------------------------

D	Gallstones
---	------------

E	Diabetes mellitus
---	-------------------

Rate this question:

[Next Question](#)

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	4
Responses Total:	4
Responses - % Correct:	0%

Back to Filters

Question 5 of 203

A 31-year-old housewife attends the diabetic clinic for her annual review. She has been a Type 1 diabetic for 8 years and her blood sugar level is usually between 5 and 9 mmol/l. She leads a sedentary lifestyle, is obese and smokes about 10 cigarettes/day. On examination her blood pressure is 140/95 mmHg and her pulse is 68 bpm and regular.

Blood results:

Hb	13.4 g/dl
WCC	5.2 x10 ⁹ /l
MCV	78 fl
plts	214 x10 ⁹ /l
HbA _{1c}	69.40 mmol/mol (8.5%)
Na	135 mmol/l
K	4.0 mmol/l
urea	7.3 mmol/l
creatinine	131 mol/l
24 hour protein	20 mg

There is evidence of background diabetic retinopathy.

What is the likely outcome of intensively treating her hyperglycaemia?

- A

There would be no significant effect on the occurrence of neuropathy
- B

Reduce the occurrence of microalbuminuria by at least 30%
- C

Significantly reduce the occurrence of retinopathy by at least 70%
- D

Reduce the risk of myocardial infarction by 75%
- E

She is likely to need renal replacement therapy within the next 10 years

2524

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 5 of 203

A 31-year-old housewife attends the diabetic clinic for her annual review. She has been a Type 1 diabetic for 8 years and her blood sugar level is usually between 5 and 9 mmol/l. She leads a sedentary lifestyle, is obese and smokes about 10 cigarettes/day. On examination her blood pressure is 140/95 mmHg and her pulse is 68 bpm and regular.

Blood results:

Hb	13.4 g/dl
WCC	5.2 x10 ⁹ /l
MCV	78 fl
plt's	214 x10 ⁹ /l
HbA _{1c}	69.40 mmol/mol (8.5%)
Na	135 mmol/l
K	4.0 mmol/l
urea	7.3 mmol/l
creatinine	131 mol/l
24 hour protein	20 mg

There is evidence of background diabetic retinopathy.

What is the likely outcome of intensively treating her hyperglycaemia?

- A

There would be no significant effect on the occurrence of neuropathy
- B

Reduce the occurrence of microalbuminuria by at least 30%
- C

Significantly reduce the occurrence of retinopathy by at least 70%
- D

Reduce the risk of myocardial infarction by 75%
- E

She is likely to need renal replacement therapy within the next 10 years

Explanation

The Diabetes Control and Complications Trial was a long-term trial that examined whether intensive treatment, with the goal of maintaining blood glucose concentrations close to the normal range, could decrease the microvascular complications of insulin-dependent diabetes. Intensive therapy slowed the progression of retinopathy by 54% compared with conventional therapy for the secondary-intervention cohort. Intensive therapy reduced the occurrence of microalbuminuria (urinary albumin excretion of = >30 mg) by 39% and that of clinical neuropathy by 60%. The main side effect of intensive therapy was an increase in severe hypoglycaemia. The patient in the above scenario did not have microalbuminuria. The link EDIC follow up study showed these benefits persisted even after the end of the trial.

2524

Rate this question: ⓪☆☆☆☆☆

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	5
Responses Total:	5
Responses - % Correct:	0%

Back to Filters

Question 6 of 203

A 23-year-old woman presents with a 3-week history of increasing swelling of her ankles. There are no preceding symptoms and there is no past medical or drug history of note. She then develops hand and facial swelling and finds her trousers increasingly tight around the waist. She has no rash, joint pains or other symptoms. On examination, her blood pressure is 160/70 mmHg, she has gross pitting oedema of both legs, ascites and facial, hand and sacral oedema.

Investigations show:

Na ⁺	140 mmol/l
K ⁺	4.2 mmol/l
Urea	10.2 mmol/l
Creatinine	67 μmol/l
Hb	8.5 g/dl
WCC	5.1 × 10 ⁹ /l
PLT	243 × 10 ⁹ /l
Albumin	16 g/l
Glucose	4 mmol/l
Cholesterol	10 mmol/l
24-hour urine collection	4.2 g/24-hours proteinuria
Chest X-ray	Small bilateral pleural effusions
Urine dipstick	4+ proteinuria and no blood

What is the most likely diagnosis?

- A

Post-infectious glomerulonephritis
- B

Goodpasture’s syndrome
- C

Minimal change disease
- D

IgA nephropathy
- E

Diabetic nephropathy

6358

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 6 of 203

Investigations show.

Na ⁺	140 mmol/l
K ⁺	4.2 mmol/l
Urea	10.2 mmol/l
Creatinine	67 μmol/l
Hb	8.5 g/dl
WCC	5.1×10^9 /l
PLT	243×10^9 /l
Albumin	16 g/l
Glucose	4 mmol/l
Cholesterol	10 mmol/l
24-hour urine collection	4.2 g/24-hours proteinuria
Chest X-ray	Small bilateral pleural effusions
Urine dipstick	4+ proteinuria and no blood

What is the most likely diagnosis?

- | | |
|---|------------------------------------|
| A | Post-infectious glomerulonephritis |
| B | Goodpasture's syndrome |
| C | Minimal change disease |
| D | IgA nephropathy |
| E | Diabetic nephropathy |

Explanation



- C Minimal change disease

A Post-infectious glomerulonephritis

B Goodpasture's syndrome

D IgA nephropathy

E	Diabetic nephropathy
---	----------------------

Rate this question:

[Next Question](#)

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Easy

Peer Responses %

Responses Correct:	0
Responses Incorrect:	6
Responses Total:	6
Responses - % Correct:	0%

Question 7 of 203

Investigations show:

Na ⁺	140 mmol/l
K ⁺	4.2 mmol/l
Urea	45 mmol/l
Creatinine	800 μmol/l
Hb	6 g/dl
WCC	$5.1 \times 10^9/l$
PLT	$243 \times 10^9/l$
Chest X-ray	Bilateral air space shadowing throughout both lung fields
Urine microscopy	Red blood cells and red blood cell casts

3

- | | |
|---|--|
| A | Goodpasture's syndrome |
| B | Cryoglobulinaemia |
| C | Antineutrophil cytoplasmic antibody (ANCA)-positive vasculitis |
| D | IgA nephropathy |
| E | Uric acid nephropathy |

Skip Question

Question 7 of 203

Investigations show that

Na ⁺	140 mmol/l
K ⁺	4.2 mmol/l
Urea	45 mmol/l
Creatinine	800 μmol/l
Hb	6 g/dl
WCC	$5.1 \times 10^9/l$
PLT	$243 \times 10^9/l$
Chest X-ray	Bilateral air space shadowing throughout both lung fields
Urine microscopy	Red blood cells and red blood cell casts

3

- | | |
|---|--|
| A | Goodpasture's syndrome |
| B | Cryoglobulinaemia |
| C | Antineutrophil cytoplasmic antibody (ANCA)-positive vasculitis |
| D | IgA nephropathy |
| E | Uric acid nephropathy |

- | | |
|---|--|
| C | Antineutrophil cytoplasmic antibody (ANCA)-positive vasculitis |
|---|--|

This is a classic history for Wegener's granulomatosis. She would need treatment with dialysis and immunosuppression and most centres would advocate plasma exchange with a creatinine above 400 $\mu\text{mol/l}$. None of the other diagnoses would classically cause sinus or joint involvement, and pulmonary haemorrhage is typically seen only in Goodpasture's syndrome and ANCA-positive vasculitis. Note left ventricular failure can lead to pink sputum and may be confused with pulmonary haemorrhage. To confirm the diagnosis of pulmonary haemorrhage, a raised KCO would be seen on lung function tests. The low haemoglobin in the setting of acute renal failure and the disproportionate degree of abnormality seen on chest X-ray compared with clinical findings suggest pulmonary haemorrhage rather than infection or fluid to explain the chest X-ray findings.

- A Goodpasture's syndrome

Goodpasture's syndrome would present with rapidly progressive glomerulonephritis and pulmonary haemorrhage. Severe pulmonary haemorrhage may require intubation. Sinus congestion and upper ear, nose and throat symptoms suggest Wegener's granulomatosis rather than Goodpasture's syndrome.

- | | |
|---|-------------------|
| B | Cryoglobulinaemia |
|---|-------------------|

Cryoglobulins deposited in medium and small vessels in the glomeruli lead to complement fixation and inflammation injury which leads to glomerulonephritis. This may be related to lymphoproliferative diseases such as myeloma and Waldenström's macroglobulinaemia or infection such as hepatitis C.

- D IgA nephropathy

There are no ear, nose and throat symptoms nor pulmonary haemorrhage with IgA nephropathy. Furthermore ANCA would be negative.

- | | |
|---|-----------------------|
| E | Uric acid nephropathy |
|---|-----------------------|

Uric acid nephropathy may present with renal colic due to uric acid stones. There are no ear, nose and throat symptoms as in this case.

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	7
Responses Total:	7
Responses - % Correct:	0%

Back to Filters

Question 8 of 203

A 25-year-old mechanic presents to his general practitioner (GP) with a 3-week history of worsening leg swelling. The rest of the history is unremarkable. He is on no medication. Examination confirms pitting oedema to the thighs. Blood pressure is 125/63 mmHg with no postural drop.

Blood results are shown belowm (the sample was noted to be lipaemic).

Na ⁺	121 mmol/l
K ⁺	4.6 mmol/l
Urea	4.4 mmol/l
Creatinine	90 mol/l
Albumin	17 g/l
Glucose	4.6 mmol/l
Plasma osmolality	280 mOsmol/kg
24 hour urinary protein	6.2 g

What investigation would help explain this patient’s hyponatraemia?

- A

Plasma antidiuretic hormone
- B

Plasma lipids
- C

Spot urinary sodium
- D

Short synacthen test
- E

Urine osmolality

6369

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 8 of 203

Blood results are shown belowm (the sample was noted to be lipaemic).

Na ⁺	121 mmol/l
K ⁺	4.6 mmol/l
Urea	4.4 mmol/l
Creatinine	90 μmol/l
Albumin	17 g/l
Glucose	4.6 mmol/l
Plasma osmolality	280 mOsmol/kg
24 hour urinary protein	6.2 g

三

- | | |
|---|-----------------------------|
| A | Plasma antidiuretic hormone |
| B | Plasma lipids |
| C | Spot urinary sodium |
| D | Short synacthen test |
| E | Urine osmolality |

- | | |
|---|---------------|
| B | Plasma lipids |
|---|---------------|

A Plasma antidiuretic hormone

C	Spot urinary sodium
---	---------------------

- Spot urinary sodium may be useful to differentiate the cause of acute renal failure. In cases of acute renal failure with a high urinary sodium would suggest a pre-renal cause.

- | | |
|---|----------------------|
| D | Short synacthen test |
|---|----------------------|

E	Urine osmolality
---	------------------

- Urine osmolality would be high in cases of pre-renal renal failure as the urine is more concentrated. It would not help in this case.

Rate this question:

Next Question

Tag Question

End Session

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	8
Responses Total:	8
Responses - % Correct:	0%

Back to Filters

Question 9 of 203

A 44-year-old female received a cadaveric renal transplant 8 weeks previously. At a routine transplant clinic appointment she is well and on the following medication: ciclosporin, azathioprine, prednisolone, co-trimoxazole and lisinopril.

Results of investigations are shown below.

Sodium	138 mmol/l
Potassium	6.9 mmol/l
Urea	11.2 mmol/l
Creatinine	139 mol/l
Bicarbonate	24 mmol/l

Electrocardiogram (ECG) Peaked T waves and prolongation of PR interval

What is the most appropriate initial therapy?

- A

Inhaled salbutamol
- B

intravenous (IV) insulin with glucose
- C

PO polystyrene sulphonate resin
- D

Immediate withdrawal of lisinopril
- E

IV calcium gluconate

6370

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 9 of 203

A 44-year-old female received a cadaveric renal transplant 8 weeks previously. At a routine transplant clinic appointment she is well and on the following medication: ciclosporin, azathioprine, prednisolone, co-trimoxazole and lisinopril.

Results of investigations are shown below.

Sodium	138 mmol/l
Potassium	6.9 mmol/l
Urea	11.2 mmol/l
Creatinine	139 mol/l
Bicarbonate	24 mmol/l

Electrocardiogram (ECG) Peaked T waves and prolongation of PR interval

What is the most appropriate initial therapy?

- A

Inhaled salbutamol
- B

intravenous (IV) insulin with glucose
- C

PO polystyrene sulphonate resin
- D

Immediate withdrawal of lisinopril
- E

IV calcium gluconate

Explanation

The answer is - IV calcium gluconate

This patient has severe hyperkalaemia with typical changes on the electrocardiogram (ECG) indicating that the hyperkalaemia is having an adverse effect on the cardiac conduction system. Although some patients show a gradual progression of ECG findings, many progress rapidly without warning. As such, patients with potassium levels greater than 6.5 mmol/l and ECG changes should be treated as a medical emergency. Protection of the heart is paramount. Calcium gluconate acts within minutes and works by raising the depolarisation threshold for myocytes. The effect of bolus calcium gluconate is transient, and repeated boluses may be necessary until the plasma potassium has been lowered. The fastest way to decrease the extracellular potassium is to shift the potassium to the intracellular compartment. This can be accomplished by insulin (along with glucose to prevent hypoglycaemia).

6370

Rate this question:

⊖

★

★

★

★

★

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	9
Responses Total:	9
Responses - % Correct:	0%

Question 10 of 203

Water deprivation phase:

DDAVP Phase:

What diagnosis would best fit the clinical picture and investigation result?

- 6446

[Previous Question](#)

Skip Question

Back to Filters

Question 10 of 203

A 21-year-old Nigerian woman was referred by her GP with a progressive history of polydipsia and polyuria of 6 months duration. She has a history of sickle cell disease and had been admitted on two previous occasions to hospital with chest pain precipitated by crisis. Fasting blood glucose is 4.5 mmol/l. She is not on any medication. On examination, her blood pressure is 120/60 mmHg and heart rate is 90 bpm. A water deprivation test was performed.

Water deprivation phase:

Plasma osmolality	298 mosm/kg (278–305 mosm/kg)
Urine osmolality	300 mosm/kg (350-1000 mosm/kg)

DDAVP Phase:

Plasma osmolality	295 mosm/kg (278–305 mosm/kg)
Urine osmolality	325 mosm/kg (350-1000 mosm/kg)

What diagnosis would best fit the clinical picture and investigation result?

- A

Analgesic nephropathy
- B

Cranial diabetes insipidus
- C

Lithium-induced nephrotoxicity
- D

Nephrogenic diabetes insipidus
- E

Psychogenic polydipsia

Explanation

- D

Nephrogenic diabetes insipidus

Nephrogenic diabetes insipidus (NDI) is characterised by the inability to concentrate urine due to the lack of response of renal tubules to antidiuretic hormone (ADH). Normally the kidneys concentrate urine by increasing water resorption by the collecting duct in the presence of ADH; this mechanism helps to maintain plasma osmolality and extracellular volume. Without ADH, a large amount of dilute urine is excreted. The water deprivation test after an overnight fast assesses the kidneys’ urine-concentrating ability and response to ADH. The result shows abnormally low (<350 mosmol/kg) urine osmolality, which increases only slightly after exogenous ADH (1-deamino-8-d-arginine-vasopressin (DDAVP), vasopressin). NDI is X-linked recessive.

Homozygous affected people (all males) are completely unresponsive to ADH while heterozygous females show normal or slight impairment. Acquired NDI occurs in disorders that disrupt the medulla or distal nephron and impair concentrating ability. Causes are: sickle cell nephropathy, polycystic kidney disease, pyelonephritis, amyloidosis, and certain nephrotoxins such as lithium and demeclocycline.

- A

Analgesic nephropathy

The injuries that may be caused by analgesia include renal papillary necrosis and chronic interstitial nephritis. There is also an increased risk of cancers of the urinary system. Water deprivation test is normal.

- B

Cranial diabetes insipidus

Although there is polyuria and polydipsia associated with cranial diabetes insipidus, the water deprivation test is normal and serum ADH levels low.

- C

Lithium-induced nephrotoxicity

When lithium induces acute renal failure, there is associated severe dehydration, natriuresis and water diuresis. Other symptoms include altered mental status and poor oral intake. Water deprivation test is normal.

- E

Psychogenic polydipsia

There is excessive fluid intake despite physiological stimuli to drink, therefore plasma sodium is low and urinary osmolality is low. However, water deprivation test is not affected.

6446

Rate this question: ⚙️☆☆☆☆☆

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	10
Responses Total:	10
Responses - % Correct:	0%

Back to Filters

Question 11 of 203

A 72-year-old woman was reviewed in outpatients having recently been treated with furosemide and amiloride combination tablets for heart failure. You understand that the GP inappropriately increased the number of combination tablets 6 weeks ago, rather than just increasing the furosemide component. She also has a history of angina for which she takes a slow-release nitrate preparation. On examination her blood pressure is 130/80 mmHg, chest examination was unremarkable and heart sounds were normal.

The results of her blood tests were:

Na ⁺	134 mmol/l
K ⁺	5.9 mmol/l
Urea	14.2 mmol/l
Creatinine	258 mol/l

An electrocardiogram (ECG) showed deep Q waves in the anterior leads but was otherwise unremarkable. Her amiloride has been stopped that day.

What further urgent action needs to be taken with respect to the potassium?

- A

Intravenous calcium gluconate 10%
- B

Intravenous insulin 10 U and dextrose 5 %
- C

None
- D

Nebulised salbutamol 5 mg
- E

Oral calcium resonium

6509

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Back to Filters

Question 11 of 203

A 72-year-old woman was reviewed in outpatients having recently been treated with furosemide and amiloride combination tablets for heart failure. You understand that the GP inappropriately increased the number of combination tablets 6 weeks ago, rather than just increasing the furosemide component. She also has a history of angina for which she takes a slow-release nitrate preparation. On examination her blood pressure is 130/80 mmHg, chest examination was unremarkable and heart sounds were normal.

The results of her blood tests were:

Na ⁺	134 mmol/l
K ⁺	5.9 mmol/l
Urea	14.2 mmol/l
Creatinine	258 mol/l

An electrocardiogram (ECG) showed deep Q waves in the anterior leads but was otherwise unremarkable. Her amiloride has been stopped that day.

What further urgent action needs to be taken with respect to the potassium?

- A

Intravenous calcium gluconate 10%
- B

Intravenous insulin 10 U and dextrose 5 %
- C

None
- D

Nebulised salbutamol 5 mg
- E

Oral calcium resonium

Explanation ⚙

- C

None

This woman has renal impairment and slightly elevated potassium. In the absence of ECG changes suggestive of hyperkalaemia (peaked T waves, widened QRS), no further action is required immediately. It is likely that the deep Q waves are indicative of previous myocardial infarction. Where peaked T waves and widened QRS exist in conjunction with hyperkalaemia, calcium gluconate and intravenous insulin and dextrose would be initial therapies of choice.

- A

Intravenous calcium gluconate 10%

Calcium prevents cardiac arrhythmias that occur as a result of hyperkalaemia. Caution must be exercised when administering intravenously because extravasation may lead to localised tissue necrosis.

- B

Intravenous insulin 10 U and dextrose 5 %

Insulin moves the potassium from extracellular to intracellular compartment thereby reducing serum potassium levels.

- D

Nebulised salbutamol 5 mg

Salbutamol, similar to insulin, moves potassium from extracellular to intracellular compartment. A side effect of nebulised salbutamol is tachycardia therefore care must be taken in patients who are already tachycardic.

- E

Oral calcium resonium

Calcium resonium acts by preventing absorption of potassium by the gut. As a consequence, it is slower acting. A common side effect is constipation.

6509

Rate this question: ⊖ ★ ★ ★ ★ ★

Next Question

- Previous Question

Feedback
- Tag Question

End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	11
Responses Total:	11
Responses - % Correct:	0%

Back to Filters

Question 12 of 203

A 56-year-old man with chronic renal failure is admitted for an arthroscopy of the right knee. His admission U&E reveals potassium of 5.9 mmol/l, creatinine of 450 μ mol/l and urea of 28 mmol/l.

Which of the following stems represents the best plan with respect to his surgery?

A	He is safe to continue with surgery	<div></div>
B	Surgery should be delayed until his potassium is below 5.5 mmol/l	
C	He should be commenced on calcium resonium and surgery can proceed	
D	He should be given insulin and dextrose and surgery may proceed	
E	He should be given nebulised salbutamol and surgery be postponed by 6 hours	

6622

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Question 12 of 203

Which of the following stems represents the best plan with respect to his surgery?

- | | |
|---|---|
| A | He is safe to continue with surgery |
| B | Surgery should be delayed until his potassium is below 5.5 mmol/l |
| C | He should be commenced on calcium resonium and surgery can proceed |
| D | He should be given insulin and dextrose and surgery may proceed |
| E | He should be given nebulised salbutamol and surgery be postponed by 6 hours |

Guidelines for surgery in patients with chronic renal failure suggest that surgery should be postponed until serum potassium is below 5.5 mmol/l. In addition, in patients undergoing surgery potassium should be monitored immediately after and then again 4–6 hours later. Where surgery is likely to involve significant circulatory compromise and acute on chronic renal failure may be precipitated, such as in coronary artery bypass surgery, patients should ideally be stabilised on dialysis first.

Rate this question: 

End Session

Peer Responses %

Responses Correct:	0
Responses Incorrect:	12
Responses Total:	12
Responses - % Correct:	0%

Question 13 of 203

3

3

3

3

- 3

3

3

3

3

3

3

Question 13 of 203

Investigations show:

Na ⁺	134 mmol/l
K ⁺	5.9 mmol/l
Urea	48 mmol/l
Creatinine	570 μmol/l
Hb	8.9 g/dl
WCC	$5.1 \times 10^9/l$
PLT	$243 \times 10^9 /l$

Which of the following represents the best treatment option in this patient?

- | | |
|---|------------------------------|
| A | Blood transfusion |
| B | Erythropoietin |
| C | Urgent haemodialysis |
| D | Intravenous furosemide |
| E | Intravenous iron replacement |

Explanation



- | | |
|---|----------------------|
| C | Urgent haemodialysis |
|---|----------------------|

This man has uraemic pericarditis, an indication for urgent haemodialysis. Haemorrhagic pericardial effusion and atrial fibrillation are often associated. Anticoagulation is not advised because of the risk of precipitating cardiac tamponade. Uraemic pericarditis usually resolves after a period of intensive dialysis. Uraemic neuropathy is also an indication for urgent dialysis, as is evidence of fluid overload coupled with renal failure which is non-responsive to diuretics.

- | | |
|---|-------------------|
| A | Blood transfusion |
|---|-------------------|

Giving blood transfusion in this scenario would exacerbate the problem as the patient is already fluid overloaded.

- | | |
|---|----------------|
| B | Erythropoietin |
|---|----------------|

Erythropoietin will increase the haemoglobin, but will need time to act. It may be given subcutaneously or intravenously. However, it would not be a treatment of uraemic pericarditis and has no effect on uraemia.

- D Intravenous furosemide

Furosemide may be useful to reduce fluid overload if the patient is still passing urine. However, it would potentially exacerbate uraemia and would not be the definitive treatment for uraemic pericarditis.

- | | |
|---|------------------------------|
| E | Intravenous iron replacement |
|---|------------------------------|

Since giving intravenous iron adds to fluid load, it would especially not be recommended in this case.

6623

10. $\frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$

[Previous Question](#)

Test Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Responses Correct:	0
Responses Incorrect:	1
Responses Total:	1
Responses % Correct:	0%

Back to Filters

Question 14 of 203

A 40-year-old woman with end-stage renal failure and undergoing regular haemodialysis is due to undergo renal transplantation from her sister the following day. The primary cause of her renal failure is hypertension and she has no other past medical history. On examination, her blood pressure is 145/70 mmHg, chest and abdominal examination is unremarkable.

Investigations show:

Na ⁺	134 mmol/l
K ⁺	5.8 mmol/l
Urea	15 mmol/l
Creatinine	370 mol/l
Hb	8.9 g/dl
WCC	5.1 × 10 ⁹ /l
PLT	243 × 10 ⁹ /l

Which of the following represents the best treatment plan?

- A

Give calcium resonium
- B

Consider dextrose and insulin infusion
- C

Do nothing
- D

Arrange an extra haemodialysis session
- E

Give regular salbutamol nebulisers

6624

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 14 of 203

Investigations show:

Na ⁺	134 mmol/l
K ⁺	5.8 mmol/l
Urea	15 mmol/l
Creatinine	370 μmol/l
Hb	8.9 g/dl
WCC	$5.1 \times 10^9/l$
PLT	$243 \times 10^9 /l$

Which of the following represents the best treatment plan?

- | | |
|---|--|
| A | Give calcium resonium |
| B | Consider dextrose and insulin infusion |
| C | Do nothing |
| D | Arrange an extra haemodialysis session |
| E | Give regular salbutamol nebulisers |

Explanation



- | | |
|---|--|
| D | Arrange an extra haemodialysis session |
|---|--|

A Give calcium resonium

B Consider dextrose and insulin infusion

C	Do nothing
---	------------

E Give regular salbutamol nebulisers

Rate this question:

Rate this question: [Next Question](#)[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	2
Responses Total:	2
Responses - % Correct:	0%

Back to Filters

Question 15 of 203

A 48-year-old man presents with painless microscopic haematuria discovered by his GP at a new patient medical. He also has vague right iliac fossa pain. There is a past history of tuberculosis and chest X-ray is suggestive of right apical fibrosis. His father died of a brain haemorrhage. On examination, his blood pressure is 160/100 mmHg and there is appears to be a palpable mass on balloting the right kidney.

Investigations show:

Na ⁺	134 mmol/l
K ⁺	5.5 mmol/l
Urea	15 mmol/l
Creatinine	370 mol/l
Hb	10.4 g/dl
WCC	5.1 × 10 ⁹ /l
PLT	243 × 10 ⁹ /l
Plasma viscosity	Normal

Which of the following stems fits best with this diagnosis?

- A

Medullary sponge kidney
- B

Autosomal dominant polycystic kidney disease (APKD)
- C

Wilms' tumour
- D

Renal cell carcinoma
- E

Transitional cell carcinoma

6625

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 15 of 203

A 48-year-old man presents with painless microscopic haematuria discovered by his GP at a new patient medical. He also has vague right iliac fossa pain. There is a past history of tuberculosis and chest X-ray is suggestive of right apical fibrosis. His father died of a brain haemorrhage. On examination, his blood pressure is 160/100 mmHg and there is appears to be a palpable mass on balloting the right kidney.

Investigations show:

Na ⁺	134 mmol/l
K ⁺	5.5 mmol/l
Urea	15 mmol/l
Creatinine	370 mol/l
Hb	10.4 g/dl
WCC	5.1 × 10 ⁹ /l
PLT	243 × 10 ⁹ /l
Plasma viscosity	Normal

Which of the following stems fits best with this diagnosis?



- A

Medullary sponge kidney
- B

Autosomal dominant polycystic kidney disease (APKD)
- C

Wilms' tumour
- D

Renal cell carcinoma
- E

Transitional cell carcinoma

Explanation



- B

Autosomal dominant polycystic kidney disease (APKD)

APKD may present at any age from the second decade with acute loin pain or haematuria due to cyst haemorrhage or infection, loin pain due to increased renal size, subarachnoid haemorrhage due to berry aneurysm rupture, complications of hypertension, complications of liver cyst formation or symptoms of chronic renal failure. The picture seen here with mild anaemia, hypertension and a palpable right kidney points towards the diagnosis; the normal viscosity counts against renal cell carcinoma.

- A

Medullary sponge kidney

The diagnosis of medullary sponge kidney is based on the finding of dilated collecting ducts (precalyceal ducts) that may or may not contain calculi and/or nephrocalcinosis. The classic radiological appearance is described as 'papillary blush' or 'paint brush' on early and delayed films during intravenous urography.

- C

Wilms' tumour

Wilms' tumour (nephroblastoma) is a rare kidney cancer and it affects mainly children. In fact, it is the most common cancer in children.

- D

Renal cell carcinoma

Renal cell carcinoma is the commonest type of kidney cancer in adults, accounting for between 90% and 95% of cases. The classical triad of symptoms includes haematuria, flank pain and abdominal mass, although this triad of symptoms occurs in only 10–15% of patients and is usually associated with advanced disease. Mostly, nowadays, renal cell carcinoma is asymptomatic and detected incidentally when investigating for other conditions.

- E

Transitional cell carcinoma

This is also known as urothelial carcinoma; it is the most common type of bladder cancer and cancer of the ureter, urethra and uterus. This condition may asymptomatic or present with haematuria, back pain, weight loss, or painful or frequent urination. The normal viscosity, anaemia and hypertension are not due to transitional cell carcinoma.

6625

Rate this question:

Next Question

- Previous Question

Feedback
- Tag Question

End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	3
Responses Total:	3
Responses - % Correct:	0%

Back to Filters

Question 16 of 203

A 24-year-old man presents with macroscopic haematuria. He is seen a few days following an upper respiratory tract infection. Urine is positive for blood and protein, but no growth is seen after urine culture. Blood pressure is normal. Renal biopsy reveals focal proliferative glomerulonephritis with mesangial IgA deposition.

Which of the following diagnoses fits best with the clinical picture?

- A Post-streptococcal glomerulonephritis
- B IgA nephropathy
- C Goodpasture’s syndrome
- D Wegener’s granulomatosis
- E Minimal change disease

6626

Submit

Previous Question Skip Question

Calculator 

Normal Values 

Question 16 of 203



A	Post-streptococcal glomerulonephritis
B	IgA nephropathy
C	Goodpasture's syndrome
D	Wegener's granulomatosis
E	Minimal change disease

Explanation

This is the most common form of glomerulonephritis seen world-wide and consists of focal proliferative glomerulonephritis with mesangial deposits of IgA. It is thought to occur due to an exaggerated immune response to viral or other antigens and may follow respiratory or gastrointestinal infection. They may present with asymptomatic microscopic haematuria or macroscopic haematuria, proteinuria occurs and 5% may be nephrotic. Good prognosis is associated with normal blood pressure, renal function and absence of proteinuria at presentation. The main distinguishing feature here from post-streptococcal glomerulonephritis is the presence of mesangial IgA. Although small amounts of IgA may be present in post-strep GN, significant quantities suggest a different diagnosis.

Rate this question:      

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	4
Responses Total:	4
Responses - % Correct:	0%

Back to Filters

Question 17 of 203

A 68-year-old man with a long history of hypertension and chronic renal failure (creatinine 190 μ mol/l on admission) is admitted for angiography. He is kept in over the weekend and you are asked to review him 3 days post procedure. Nursing staff have noticed a purpuric rash over his feet and think the blood supply to his lower limbs is impaired. Blood tests reveal erythrocyte sedimentation rate (ESR) of 86, eosinophilia and a creatinine level of 450 μ mol/l.

Which of the following stems represents the most likely diagnosis?

- | | |
|---|-------------------------|
| A | Acute vasculitis |
| B | Contrast nephropathy |
| C | Renal artery thrombosis |
| D | Cholesterol embolism |
| E | Renal vein thrombosis |

6627

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Question 17 of 203



osis?

osis?

osis?

osis?

osis?

osis?



osis?

osis?

osis?

osis?

osis?

osis?

Question 18 of 203

11

Sodium	140 mmol/l
Potassium	5.1 mmol/l
Chloride	99 mmol/l
Bicarbonate	21 mmol/l
Urea	19 mmol/l
Creatinine	221 mmol/l
Eosinophils	Mildly elevated

Specific gravity	1.020
Protein	2+
Blood	4+
Red cell casts	Many
24 hour protein	1.9 g

A	Antiglomerular basement membrane antibody
B	Antistreptolysin titre
C	cANCA (antineutrophil cytoplasmic antibodies)
D	pANCA
E	Serum immunoglobulin A (IgA) levels

Submit

Skip Question

Back to Filters

Question 18 of 203

A 35-year-old man with a history of mild asthma comes to your clinic with 10 days' history of malaise, fever, and tightness in the chest. On further inquiry he also mentions shortness of breath, weight loss and swelling in the legs and scrotum. There is no significant past medical history. On examination, his temperature is 38°C, blood pressure is 175/90 mmHg, pulse is 77 beats/min and respiratory rate 14 breaths/min. Physical examination confirms bilateral pedal and scrotal oedema. There are also scattered bilateral wheezes. Chest X-ray is unremarkable.

Some laboratory results are:

Sodium	140 mmol/l
Potassium	5.1 mmol/l
Chloride	99 mmol/l
Bicarbonate	21 mmol/l
Urea	19 mmol/l
Creatinine	221 mmol/l
Eosinophils	Mildly elevated

Urinalysis:

Specific gravity	1.020
Protein	2+
Blood	4+
Red cell casts	Many
24 hour protein	1.9 g

Which laboratory finding would support the most likely diagnosis?

A	Antiglomerular basement membrane antibody
B	Antistreptolysin titre
C	cANCA (antineutrophil cytoplasmic antibodies)
D	pANCA
E	Serum immunoglobulin A (IgA) levels

Explanation

This patient has Churg-Strauss syndrome: a nephritic syndrome associated with eosinophilia and asthma. It is associated with positive pANCA. The patient has nephritic syndrome as evidenced by hypertension, oedema, and haematuria. Asthma is suggested by his chest tightness and shortness of breath.

The American College of Rheumatology has defined criteria for a diagnosis of Churg-Strauss syndrome:

- Asthma
- High numbers of eosinophils
- Sensory neuropathy
- Lung abnormalities
- Sinus problems such as abnormal growths (polyps)
- A tissue sample showing a blood vessel with eosinophils around it.

A person with Churg-Strauss syndrome may have some or all of these symptoms.

Besides the criteria for diagnosis listed above, the person with Churg-Strauss syndrome may have:

- A nodular or purpuric skin rash
- Renal, intestinal or cardiac involvement
- Malaise
- Loss of appetite (anorexia) and weight loss
- Other symptoms specific to the part of the body affected in the person

Frequency

Internationally: incidence is approximately 2.5 cases per 100 000 adults per year.

Mortality/morbidity

- The principal causes of morbidity and mortality are myocarditis and myocardial infarction secondary to coronary arteritis.
- With treatment, the 1-year survival rate is 90% and the 5-year survival rate is 62%.
- Overall, without treatment, the 5-year survival rate is about 25%.

Treatment

Glucocorticoids alone usually are adequate for treatment of Churg-Strauss syndrome. Cytotoxic drugs are necessary in fewer than 20% of patients. Major life-threatening organ involvement may require treatment with pulse doses of intravenous (iv) corticosteroids as well as other cytotoxic agents. Other treatments include intravenous immune globulin, interferon-alpha, and plasma exchange. Plasma exchange has not improved the course of the disease.

Complications

Complications of Churg-Strauss syndrome depend on the specific organ system involvement.

6720

Rate this question: ⚙️ ⭐️ ⭐️ ⭐️ ⭐️ ⭐️

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Difficult

Peer Responses %

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

Question 19 of 203

What is the most appropriate next step?

- 6721

© Pastest 2017

Back to Filters

Question 19 of 203

You saw a 38-year-old gentleman in your clinic 2 days ago with sore throat, cough and fever. He says that he has been feeling increasingly tired and lethargic over the past 3 weeks. You suspected upper respiratory tract infection and prescribed cephalexin, took some routine blood tests and asked the man to return in 2 days. Today in your clinic he mentions that his urinary volume has decreased dramatically within last few days, and he is only passing few millilitres of urine. His blood results show urea of 24 mmol/l and creatinine 186 μ mol/l. You immediately admit him in the hospital and repeat bloods today show urea of 38 mmol/l and creatinine of 372 μ mol/l. His urine output in hospital is only 5–10 ml in past 2 h. The temperature is 38.3°C. He has an erythematous oropharynx with tonsillar exudates. Urinalysis shows red cell casts and dysmorphic red blood cells.

What is the most appropriate next step?

- A

Administer penicillin intravenously
- B

Prescribe a high dose of ciclosporin
- C

Give a low dose of methylprednisolone intravenously (IV)
- D

Give a high dose of methylprednisolone IV and cyclophosphamide
- E

Administer broad-spectrum antibiotic

Explanation

This is rapidly progressive glomerulonephritis. In adults where the outcome is much worse, early initiation of a high dose of steroids has been associated with improved mortality and more rapid return of kidney function.

Rapidly progressive glomerulonephritis (RPGN) is a disease of the kidney that results in a rapid decrease in the glomerular filtration rate of at least 50% over a short period (a few days to 3 months). The main pathological finding is fibrinoid necrosis (>90% of biopsy specimens); extensive crescent formation is present in at least 50% of glomeruli.

This type of glomerulonephritis is also known as crescentic glomerulonephritis after the ‘crescents’ seen on light microscopy. These crescents are collections of epithelial cells and macrophages within Bowman’s space.

The presence of crescents indicates that glomerular damage is rapid and progressive (from onset to end-stage renal failure within weeks to months).

Immunofluorescence detects deposits of immunoglobulin G (IgG) or complement factor C3 in the glomerular basement membrane. There are three distributions of immunofluorescence in crescentic glomerulonephritis:

- absent immunofluorescence
- granular immunofluorescence
- linear immunofluorescence.

Frequency

Internationally: in the United Kingdom, the frequency is estimated at 2 cases per 100 000 persons. In Sweden, the frequency is estimated at 1 case per 100 000 persons.

Mortality/morbidity

Massive pulmonary hemorrhage is the most common cause of death in patients presenting with ANCA-associated disease. However, once immunosuppressive therapy has begun, infection is more common.

Two major patterns of ANCA are

- a- proteinase 3 – PR3 ANCA (formerly cytoplasmic or cANCA): Positive in Wegener’s granulomatosis in 90% of cases
- b- Myeloperoxidase – MPO ANCA (formerly p ANCA): Positive in Chrug Strauss syndrome, and rarely reported as positive in PAN.

Clinical features

Symptoms and signs of renal failure. There may be loin pain, haematuria and systemic symptoms (fever, malaise, myalgia, weight loss).

Histological findings

Renal biopsy specimens show a diffuse, proliferative, necrotising glomerulonephritis with crescent formation.

Treatment

High-dose corticosteroids; cyclophosphamide +/- plasma exchange/renal transplantation.

Prognosis

Poor if initial creatinine >600 μ mol/l

6721

Rate this question:      

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Difficult

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	7
Responses Total:	7
Responses - % Correct:	0%

Back to Filters

Question 20 of 203

A 39-year-old man comes to the office because of having an achy low back pain for the past few weeks. He describes pain as a dull ache that came on gradually. There are no other specific symptoms. He has been taking hydrochlorothiazide and enalapril for hypertension for the last five years. His father and younger sister also have hypertension. On examination, his pulse rate was 65 beats/min, blood pressure 135/90 mmHg, and temperature 36.6°C. He has got mild tenderness on the left flank and a mid systolic click on cardiac examination. Urine dipstick showed microscopic haematuria.

What is the most appropriate next step?

- A

Arrange a kidney biopsy
- B

Ultrasound scan of the kidneys
- C

Computed tomography (CT) scan of the kidneys
- D

Intravenous pyelogram
- E

Genetic linkage analysis

6722

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 20 of 203

A 39-year-old man comes to the office because of having an achy low back pain for the past few weeks. He describes pain as a dull ache that came on gradually. There are no other specific symptoms. He has been taking hydrochlorothiazide and enalapril for hypertension for the last five years. His father and younger sister also have hypertension. On examination, his pulse rate was 65 beats/min, blood pressure 135/90 mmHg, and temperature 36.6°C. He has got mild tenderness on the left flank and a mid systolic click on cardiac examination. Urine dipstick showed microscopic haematuria.

What is the most appropriate next step?

A	Arrange a kidney biopsy
B	Ultrasound scan of the kidneys
C	Computed tomography (CT) scan of the kidneys
D	Intravenous pyelogram
E	Genetic linkage analysis

Explanation

The answer is - Ultrasound scan of the kidneys

This patient is most likely to have adult polycystic kidney disease. Ultrasound scan will be the most appropriate test as it will show multiple renal cysts. It presents with hypertension, flank pain, micro- or macroscopic haematuria and chronic renal failure. It is associated with intracranial aneurysms, colonic diverticula, mitral valve prolapse, and cysts in other organs - the liver, pancreas and spleen. Renal biopsy is not indicated in the diagnosis of adult polycystic kidney disease. Computed tomography (CT) is a sensitive test, but it is done after the ultrasound scan. Genetic linkage analysis will give the diagnosis in suspected cases when the radiographic imaging is negative. Intravenous pyelogram is the least sensitive of all the tests.

6722

Rate this question:      

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	8
Responses Total:	8
Responses - % Correct:	0%

Back to Filters

Question 21 of 203

A 58-year-old man previously in good health suddenly develops severe abdominal pain radiating from the left loin to the groin and associated with nausea, perspiration and frequent urination. There is a history of diverticular disease but nil else of note. He is restless, tossing and turning in bed. His BP is elevated at 180/90 mmHg, his pulse is 95/min and he is in obvious pain.

What is the most likely diagnosis?



- | | |
|---|--|
| A | Herpes zoster involving the left lower rib dermatome |
| B | Sigmoid diverticulitis |
| C | Torsion of the left testicle |
| D | Left ureteric calculus |
| E | Retroperitoneal haemorrhage |

6724

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 21 of 203

A 58-year-old man previously in good health suddenly develops severe abdominal pain radiating from the left loin to the groin and associated with nausea, perspiration and frequent urination. There is a history of diverticular disease but nil else of note. He is restless, tossing and turning in bed. His BP is elevated at 180/90 mmHg, his pulse is 95/min and he is in obvious pain.

What is the most likely diagnosis?

- A

Herpes zoster involving the left lower rib dermatome
- B

Sigmoid diverticulitis
- C

Torsion of the left testicle
- D

Left ureteric calculus
- E

Retroperitoneal haemorrhage

Explanation

This is a typical case of ureteric colic. It is sudden in onset radiates from loin to groin, and is associated with urge to urinate frequently. It is the pain radiating to the groin that points more towards ureteric colic versus diverticulitis, the major differential here. The usual causes are blood clots and calculi. Urine examination reveals macroscopic or microscopic haematuria.

6724

Rate this question: ⓪★ ★ ★ ★ ★

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	9
Responses Total:	9
Responses - % Correct:	0%

Back to Filters

Question 22 of 203

A 72-year-old woman comes to the emergency department with severe flank pain. On examination she is mentally obtunded with fever of 39°C, blood pressure of 80/40 mmHg and pulse rate of 110 beats/min. On further evaluation, her blood results show a white cell count of 22×10^9 with neutrophilia. Urgent ultrasound scan shows left hydronephrosis and hydroureter. Computed tomography (CT) scan confirms the presence of an obstructing stone. Intravenous vasopressors are initiated.

What will be the most appropriate next step in management?



- | | |
|---|--|
| A | Perform extracorporeal shock wave lithotripsy |
| B | Schedule a nephrectomy |
| C | Commence antibiotics and review response before inserting a nephrostomy tube |
| D | Administer antibiotics intravenously and insert a nephrostomy immediately |
| E | Commence antibiotics and schedule cystoscopy and ureteral stent placement |

6726

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Question 22 of 203

What will be the most appropriate next step in management?

- 11

This patient has obstructing pyelonephritis with sepsis. The patient needs urgent IV antibiotics with assessment of the response over the next 24 hours. If they are still spiking fevers, or complaining of significant colic, insertion of a nephrostomy tube is the next step. Lithotripsy will be an elective procedure to break large stones so that they can be passed naturally. This is not suitable for an urgent next step in treatment.

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Session

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	10
Responses Total:	10
Responses - % Correct:	0%

Back to Filters

Question 23 of 203

A 25-year-old woman comes to the office with a 3-day history of lower abdominal pain that is now beginning to radiate to her right flank. She denies nausea, vomiting or diarrhoea but states she feels feverish and is worried that she may have the flu. She has noticed an increase in her urinary frequency.

Vital signs are temperature 39.2°C, blood pressure 120/70 mmHg, pulse 95/min, and respiration 16/min. Physical examination shows tenderness in the suprapubic region, hyperactive bowel sounds and no signs of peritoneal irritation or masses. Her right flank is tender to light percussion. The remainder of the physical examination is within normal limits.

Some of the laboratory results are:

Urinalysis

Leucocytes	+++
Proteinuria	+
Gram stain	Gram-negative bacilli

What is the most likely diagnosis?

- A

Acute appendicitis
- B

Acute pyelonephritis
- C

Ovarian torsion
- D

Perinephric abscess
- E

Pelvic inflammatory disease

6727

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 23 of 203

Vital signs are temperature 39.2°C, blood pressure 120/70 mmHg, pulse 95/min, and respiration 16/min. Physical examination shows tenderness in the suprapubic region, hyperactive bowel sounds and no signs of peritoneal irritation or masses. Her right flank is tender to light percussion. The remainder of the physical examination is within normal limits.

Some of the laboratory results are:

Urinalysis

Leucocytes	+++
Proteinuria	+
Gram stain	Gram-negative bacilli

What is the most likely diagnosis?

- | | |
|---|-----------------------------|
| A | Acute appendicitis |
| B | Acute pyelonephritis |
| C | Ovarian torsion |
| D | Perinephric abscess |
| E | Pelvic inflammatory disease |

Explanation

This is a typical picture of pyelonephritis. Additional testing such as renal ultrasonography or intravenous pyelography is indicated only if there is suspicion of a complication such as a stone stricture or tumour causing the pyelonephritis. These tests are not necessarily part of routine management of a simple uncomplicated pyelonephritis. If symptoms persist beyond 3 to 5 days of effective treatment an ultrasound or computed tomography (CT) scan of the kidney should be obtained to exclude a perinephric abscess or other drainable collection of fluid.

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	11
Responses Total:	11
Responses - % Correct:	0%

Back to Filters

Question 24 of 203

A 69-year-old man with hypertension comes to the emergency department complaining of dysuria, flank pain, fever and chills for 2 days. He takes metoprolol and hydrochlorothiazide for his hypertension. Vital signs are temperature 39.3°C, blood pressure 100/60 mmHg, pulse 120/min, respiration 28/min, oxygen saturation 98% on room air. He has mild tenderness on the left costovertebral angle. The rest of the physical examination is normal. The chest X Ray is normal and urine dipstick positive for blood, protein and nitrites. Blood and urine are sent for analysis and cultures.

What is the most appropriate next step in management?

A

Call for a urological consultation

B

Insert a foley catheter

C

Obtain an ultrasound of the kidneys

D

Start a dopamine infusion

E

Start intravenous antibiotic therapy

6730

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 24 of 203

What is the most appropriate next step in management?

- ### Explanation

Rate this question:

End Session

Peer Responses %

Session Progress

[Blog](#) [About Pastest](#) [Contact Us](#) [Help](#)

Back to Filters

Question 25 of 203

A 64-year-old man comes to the office complaining of left flank pain and gross haematuria. The pain is non-radiating, dull in nature and not associated with any fever or chills. The patient has not had any medical care in 20 years. His vital signs and physical examination are unremarkable. A complete blood count and biochemical profile are normal. A computed tomography (CT) scan of the abdomen and pelvis reveals a contrast enhancing 5 cm mass in the lower pole of the left kidney. A chest X-ray and ultrasound of the abdomen are unremarkable, as is a CT thorax.

What is the most appropriate way to manage this patient?



- | | |
|---|--|
| A | Order a MIBI scan |
| B | order a magnetic resonance image (MRI) with gadolinium |
| C | Schedule chemotherapy |
| D | Schedule radiation therapy |
| E | Send him to a urologist for radical nephrectomy |

6731

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Back to Filters

Question 25 of 203

A 64-year-old man comes to the office complaining of left flank pain and gross haematuria. The pain is non-radiating, dull in nature and not associated with any fever or chills. The patient has not had any medical care in 20 years. His vital signs and physical examination are unremarkable. A complete blood count and biochemical profile are normal. A computed tomography (CT) scan of the abdomen and pelvis reveals a contrast enhancing 5 cm mass in the lower pole of the left kidney. A chest X-ray and ultrasound of the abdomen are unremarkable, as is a CT thorax.

What is the most appropriate way to manage this patient?

- | | |
|---|--|
| A | Order a MIBI scan |
| B | order a magnetic resonance image (MRI) with gadolinium |
| C | Schedule chemotherapy |
| D | Schedule radiation therapy |
| E | Send him to a urologist for radical nephrectomy |

Explanation

This patient has renal cell carcinoma of the left kidney. Computed tomography (CT) scan with and without contrast that reveals a contrast-enhancing mass is the primary radiological modality used to diagnose this disease. This patient presents with two of the three symptoms of the classic triad of renal cell carcinoma: costovertebral pain, hematuria and palpable mass. Unfortunately this triad is only seen in 10% of the patients diagnosed with renal cell carcinoma. Surgical removal remains the only potential curative therapy for renal cell carcinoma. While a variety of therapeutic modalities are available (i.e. partial nephrectomy), radical nephrectomy provides this patient with the best possible chance of cure. Prognosis for patients with T1 renal cell carcinoma (less than 7 cm in size) is >85% with appropriate treatment.

6731

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	13
Responses Total:	13
Responses - % Correct:	0%

Back to Filters

Question 26 of 203

A 70-year-old-man was admitted with anuria and lower back pain of 24 h duration. His only medication is bendroflumethiazide 2.5 mg for hypertension. He is awaiting an appointment for urinary frequency, urgency and dribbling, otherwise he is fit and well.

Investigations reveal:

Haemoglobin (Hb)	12.5 g/dl
White blood count (WBC)	5.6 × 10 ⁹ /l
Platelets	160 × 10 ⁹ /l
Sodium	134 mmol/l
Potassium	6.5 mmol/l
Urea	20.6 mmol/l
Creatinine	400 μmol/l
Corrected calcium	2.70 mmol/l
Albumin	35 g/l
Total protein	80 g/l



You have given him IV insulin and dextrose to reduce his potassium, a follow up is measured at 5.4 mmol/l and you catheterize him, which of the following is the next most appropriate management?

- A

Urgent haemodialysis
- B

Central venous line for monitoring intravascular volume
- C

Urgent intravenous urography (IVU)
- D

Restrict oral fluid
- E

Urgent renal tract ultrasound

6971

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 26 of 203

A 70-year-old-man was admitted with anuria and lower back pain of 24 h duration. His only medication is bendroflumethiazide 2.5 mg for hypertension. He is awaiting an appointment for urinary frequency, urgency and dribbling, otherwise he is fit and well.

Investigations reveal:

Haemoglobin (Hb)	12.5 g/dl
White blood count (WBC)	5.6 × 10 ⁹ /l
Platelets	160 × 10 ⁹ /l
Sodium	134 mmol/l
Potassium	6.5 mmol/l
Urea	20.6 mmol/l
Creatinine	400 μmol/l
Corrected calcium	2.70 mmol/l
Albumin	35 g/l
Total protein	80 g/l



You have given him IV insulin and dextrose to reduce his potassium, a follow up is measured at 5.4 mmol/l and you catheterize him, which of the following is the next most appropriate management?

- A

Urgent haemodialysis
- B

Central venous line for monitoring intravascular volume
- C

Urgent intravenous urography (IVU)
- D

Restrict oral fluid
- E

Urgent renal tract ultrasound

Explanation

The symptom of urinary frequency and the age of the patient suggest probable prostatic hypertrophy. Prostatic obstruction is the commonest cause of obstructive uropathy in men. In community studies, urinary tract obstruction is responsible for 25% of cases of acute renal failure. It is easily treatable if diagnosis is made accurately and early. The cause of the acute presentation in this case is most likely due to obstructive uropathy. The first investigation should be renal and pelvic ultrasound to exclude hydronephrosis which, if present, warrant urgent nephrostomy for ureteric obstruction, or transurethral/supra-pubic catheter as appropriate for bladder obstruction. Renal function often normalises with relief of the obstruction.

6971

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	14
Responses Total:	14
Responses - % Correct:	0%

Back to Filters

Question 27 of 203

An 18-year-old girl presented with a 3-day history of fever, abdominal pain, diarrhoea and vomiting. She had returned from holiday in Morocco 1 week ago. She has no significant past medical history and she was well while on holiday.

On examination, she is unwell and slightly jaundiced. There is no lymphadenopathy and abdominal examination was normal. Temperature was 37.8°C, blood pressure was 98/56 mmHg, pulse of 110 beats/min and Glasgow Coma score of 15/15.

Investigations are shown below:

Haemoglobin (Hb)	6.9 g/l
White cell count (WCC)	15 × 10 ⁹ g/l
Platelets	22 × 10 ⁹ /l
Mean corpuscular volume (MCV)	86 fl
Reticulocyte count	160 × 10 ⁹ /l (25-100 ×10 ⁹ /l)
Blood film	Multiple fragmented red cells
Sodium	134 mmol/l
Potassium	4.5 mmol/l
Urea	21.5 mmol/l
Creatinine	350 μmol/l
Bilirubin	58 μmol/l
Lactate dehydrogenase	300 IU/l (70-250 IU/l)
Aspartate transaminase	38 IU/l
Alanine aminotransferase	36 IU/l

What is the most likely underlying diagnosis?

- A

Thrombotic thrombocytopaenic purpura (TTP)
- B

Leptospirosis
- C

Familial Mediterranean fever
- D

Haemolytic-uraemic syndrome (HUS)
- E

Acute hepatitis B infection

6973

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 27 of 203

Haemoglobin (Hb)

Thrombotic thrombocytopenic purpura (TTP)

B	Leptospirosis
---	---------------

C Familial Mediterranean Fever

D	Haemolytic-uraemic syndrome (HUS)
---	-----------------------------------

E	Acute hepatitis B infection
---	-----------------------------

10

- Children account for 90% of the cases of HUS, while adults are more frequently affected with TTP.
- Renal failure and fever predominates in HUS, while in TTP neurological symptoms are more common.
- The prognosis is worse in TTP compared with HUS.
- Treatment is supportive with plasma exchange until the activity of the disease has subsided. Steroids are also beneficial.

B	Leptospirosis
---	---------------

C Familial Mediterranean fever

E Acute hepatitis B infection

Rate this question:

100

Tag Question

End Session

Difficulty: Average

Responses incorrect:	15
Responses Total:	15
Responses - % Correct:	0%

Blog About D

Question 28 of 203

On examination he is dyspnoeic and slightly jaundiced, with widespread spider naevi. He also has multiple purpuric rashes on his legs with a reticulated pattern to the skin. The leg ulcer is below the right knee with large ulceration and necrosis.

PH	7.45
p _a (O ₂)	7.2 kPa
p _a (CO ₂)	3.5 kPa
Chest X-ray	normal
Sodium	135 mmol/l
Potassium	4.8 mmol/l
Urea	18 mmol/l
Creatinine	280 μmol/l
Corrected calcium	2.5 mmol/l
Albumin	27 g/l
Bilirubin	60 μmol/l
Aspartate transaminase (AST)	200 IU/l
Alanine aminotransferase (ALT)	289 IU/l
Haemoglobin (Hb)	11.5 g/dl
White cell count (WCC)	10.5 × 10 ⁹ /l
Platelets	200 × 10 ⁹ /l
Mean corpuscular volume (MCV)	88 fl
Urinalysis	Protein +++, blood +++

3

- | | |
|---|---------------------------------|
| A | Multiple myeloma |
| B | Rheumatoid arthritis |
| C | Waldestrom's macroglobulinaemia |
| D | Cryoglobulinaemia |
| E | Hepatocellular carcinoma |

Submit

Skip Question

Question 28 of 203

A 50-year -old Turkish male with a 3-year history of liver cirrhosis secondary to hepatitis C was admitted with a right gangreneous leg ulcer of 2 months’ duration with pain in his joints. He has developed pleuritic chest pain with dyspnoea over the last few days. He is on interferon for his liver disease. He is a non-smoker and does not drink.

On examination he is dyspnoeic and slightly jaundiced, with widespread spider naevi. He also has multiple purpuric rashes on his legs with a reticulated pattern to the skin. The leg ulcer is below the right knee with large ulceration and necrosis.

Investigations reveal:

PH	7.45
p _a (O ₂)	7.2 kPa
p _a (CO ₂)	3.5 kPa
Chest X-ray	normal
Sodium	135 mmo/l
Potassium	4.8 mmol/l
Urea	18 mmol/l
Creatinine	280 mol/l
Corrected calcium	2.5 mmol/l
Albumin	27 g/l
Bilirubin	60 mol/l
Aspartate transaminase (AST)	200 IU/l
Alanine aminotransferase (ALT)	289 IU/l
Haemoglobin (Hb)	11.5 g/dl
White cell count (WCC)	10.5 × 10 ⁹ /l
Platelets	200 × 10 ⁹ /l
Mean corpuscular volume (MCV)	88 fl
Urinalysis	Protein +++, blood +++

What is the underlying diagnosis?

- A

Multiple myeloma
- B

Rheumatoid arthritis
- C

Waldestrom’s macroglobulinaemia
- D

Cryoglobulinaemia
- E

Hepatocellular carcinoma

Explanation



- D

Cryoglobulinaemia

Cryoglobulins are immunoglobulins that precipitate when blood is cooled while flowing through the skin and subcutaneous tissues. This leads to small vessel damage in the limbs and deposition on the wall of small vessels resulting in generalised vasculitis, which presents with a reticulated skin pattern of microthrombosis and areas of gangrene. Pulmonary embolism, and arterial and venous thrombosis are common. Vasculitis results in glomerulonephritis, polyneuropathy, Raynaud’s phenomenon and leg ulcers. Recognised associations are chronic infections such as hepatitis C, systemic lupus erythematosus, rheumatoid arthritis and Sjögren’s syndrome. Management is with plasma exchange, chemotherapy and treatment of the underlying condition.

- A

Multiple myeloma

Multiple myeloma is a haematological malignancy. It is characterised by an aberrant clone of plasma cells producing immunoglobulin (IgG or IgA) or immunoglobulin light chain. It generally presents in the over 60s with a male preponderance. Median survival is 3 years. The usual presentation is unexplained renal impairment with normal-sized kidneys and a bland urine deposit. Bone pain, weakness, fatigue, easy bruising, pallor and hepatomegaly (20%) are other features.

- B

Rheumatoid arthritis

This history includes large ulcerations which are not a usual feature of rheumatoid arthritis per se, although patients with rheumatoid arthritis are at increased risk of ulcers.

- C

Waldestrom’s macroglobulinaemia

Waldestrom’s macroglobulinaemia is a type of non-Hodgkin lymphoma. There is a large amount of macroglobulin produced. Features include weakness, fatigue, weight loss, and chronic oozing of blood from the nose and gums. Peripheral neuropathy occurs in 10% of patients. Lymphadenopathy and hepatosplenomegaly are present in 30-40% of cases. Other possible signs and symptoms include blurring or loss of vision, headache, and (rarely) stroke or coma.

- E

Hepatocellular carcinoma

Although hepatitis C with chronic liver disease is associated with hepatocellular carcinoma; ulceration and skin necrosis is not a feature of hepatocellular carcinoma.

6974

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	16
Responses Total:	16
Responses - % Correct:	0%

Question 29 of 203

A water deprivation test was performed:

Plasma osmolality	330 mOsmol/kg (285-295 mOsmol/kg)
Urine osmolality	220 mOsmol/kg

Plasma osmolality	300 mOsmol/kg
Urine osmolality	250 mOsmol/kg

- | | |
|---|------------------------------------|
| A | Psychogenic polydipsia |
| B | Cranial diabetes insipidus |
| C | Partial cranial diabetes insipidus |
| D | Nephrogenic diabetes insipidus |
| E | Diabetes mellitus |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 29 of 203

A 28-year-old female sustained a minor head injury, was seen in the emergency department with headache and was discharged following a normal computed tomography of the head. She presented again 2 months later with polyuria and polydipsia. She has a 2-year history of recurrent pyelonephritis and is under investigation by the urologists.

A water deprivation test was performed:

Water deprivation phase

Plasma osmolality	330 mOsmol/kg (285–295 mOsmol/kg)
Urine osmolality	220 mOsmol/kg

Desmopressin (DDAVP) phase (after administration of 2 μg desmopressin)

Plasma osmolality	300 mOsmol/kg
Urine osmolality	250 mOsmol/kg

What is the diagnosis?



- A

Psychogenic polydipsia
- B

Cranial diabetes insipidus
- C

Partial cranial diabetes insipidus
- D

Nephrogenic diabetes insipidus
- E

Diabetes mellitus

Explanation



- D

Nephrogenic diabetes insipidus

In nephrogenic diabetes insipidus, the water deprivation test after overnight fast shows the maximal osmolality of urine is abnormally low <250 mosmol/kg. After administration of exogenous vasopressin (DDAVP), there is only slight increase in urine osmolality <50%.

- A

Psychogenic polydipsia

In psychogenic polydipsia, a low initial plasma and urine osmolality is usual at the start of the test. This increases after water deprivation since vasopressin can be stimulated normally.

- B

Cranial diabetes insipidus

In cranial diabetes insipidus, DDAVP produces a rise in urine osmolality of about 50%.

- C

Partial cranial diabetes insipidus

In partial cranial diabetes insipidus, there is urinary osmolality of 300–750 mosmol/kg after water deprivation and this remains below 750 mosmol/kg after administration of vasopressin.

- E

Diabetes mellitus

There is polyuria and polydipsia in uncontrolled diabetes with high blood sugar. But the water deprivation test would yield normal results.

6976

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	17
Responses Total:	17
Responses - % Correct:	0%

Back to Filters

Question 30 of 203

A 65-year-old male was admitted with pneumonia and hyponatraemia with sodium of 116 mmol/l. He was treated with antibiotics and rapid infusion of 3% hypertonic sodium chloride, but despite initial improvement deteriorated neurologically with fits and subsequent coma.

What is the possible cause of his deterioration?



- | | |
|---|--|
| A | Syndrome of inappropriate antidiuretic hormone secretion |
| B | Acute cerebrovascular event |
| C | Central pontine myelinolysis |
| D | Normal pressure hydrocephalus |
| E | Encephalomyelitis |

6977

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Question 30 of 203

- | | |
|---|--|
| A | Syndrome of inappropriate antidiuretic hormone secretion |
| B | Acute cerebrovascular event |
| C | Central pontine myelinolysis |
| D | Normal pressure hydrocephalus |
| E | Encephalomyelitis |

- | | |
|---|------------------------------|
| C | Central pontine myelinolysis |
|---|------------------------------|

A Syndrome of inappropriate antidiuretic hormone secretion

B	Acute cerebrovascular event
---	-----------------------------

D Normal pressure hydrocephalus

E	Encephalomyelitis
---	-------------------

Rate this question:

[Next Question](#)

Previous Question

Tag Question

Feedback

End Session

End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	18
Responses Total:	18
Responses - % Correct:	0%

Back to Filters

Question 31 of 203

A 63-year-old man was admitted to hospital for an elective coronary angiogram. The procedure went ahead without any obvious complications. A day after the procedure he complained of generalised aching abdominal pain which improved following administration of lactulose, although it did not resolve entirely. A day later he became acutely unwell. On examination he appeared drowsy and confused. He had a pulse rate of 100/minute with a blood pressure of 135/80 mmHg. He had peripheral oedema and coarse crackles in both lung bases. His abdominal examination showed diffuse tenderness but no rebound or guarding. There was a bluish discoloration of his toes, and signs of peripheral emboli. His urgent blood test results are as follows:

Haemoglobin (Hb)	15.3 g/dl
White cell count (WCC)	10.0 × 10 ⁹ /l
Platelets	349 × 10 ⁹ /l
Neutrophils	6.0 × 10 ⁹ /l (1.5-7.0)
Lymphocytes	2.0 × 10 ⁹ /l (1.5-4.0)
Eosinophils	2.1 × 10 ⁹ /l (0.04-0.40)
Basophils	0.1 × 10 ⁹ /l (<0.1)
Monocytes	0.2 × 10 ⁹ /l (<0.8)
Na ⁺	143 mmol/l
K ⁺	6.2 mmol/l
Urea	18.5 mmol/l
Creatinine	490 μmol/l

The most likely diagnosis is:

- A

Acute renal failure secondary to contrast induced nephropathy
- B

Acute renal failure secondary to generalised sepsis
- C

Acute renal failure secondary to multiple cholesterol emboli
- D

Acute renal failure secondary to catastrophic intra-abdominal event
- E

Acute renal failure secondary to exacerbation of congestive cardiac failure

7014

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 31 of 203

His urgent blood test results are as follows:

Haemoglobin (Hb)	15.3 g/dl
White cell count (WCC)	$10.0 \times 10^9/\text{l}$
Platelets	$349 \times 10^9/\text{l}$
Neutrophils	$6.0 \times 10^9/\text{l}$ (1.5-7.0)
Lymphocytes	$2.0 \times 10^9/\text{l}$ (1.5-4.0)
Eosinophils	$2.1 \times 10^9/\text{l}$ (0.04-0.40)
Basophils	$0.1 \times 10^9/\text{l}$ (<0.1)
Monocytes	$0.2 \times 10^9/\text{l}$ (<0.8)
Na ⁺	143 mmol/l
K ⁺	6.2 mmol/l
Urea	18.5 mmol/l
Creatinine	490 $\mu\text{mol/l}$

A	Acute renal failure secondary to contrast induced nephropathy
B	Acute renal failure secondary to generalised sepsis
C	Acute renal failure secondary to multiple cholesterol emboli
D	Acute renal failure secondary to catastrophic intra-abdominal event
E	Acute renal failure secondary to exacerbation of congestive cardiac failure

This man has developed acute renal failure and is severely unwell. We are not told of any risk factors that would increase the chance of contrast-induced nephrotoxicity (such as age over 70 years; co-existing nephrotoxicity, especially diabetic; concomitant administration of nephrotoxic drugs; dehydration prior to procedure; or pre-existing congestive cardiac failure). His presentation with abdominal pain and eosinophilia all support a diagnosis of multiple cholesterol emboli. There are no convincing signs of severe sepsis. The abdominal pain is probably caused by bowel ischaemia, but the lack of signs of peritonism in this case make this a less likely primary cause for his deterioration, and the degree of cardiac insufficiency is unlikely to have caused such a significant deterioration. Data about cholesterol embolisation are largely based on post mortem examinations which are, by definition, a selected group of poor outcomes. However, some studies suggest that as many as 25–30% of patients undergoing angiography will have atherosclerotic emboli.

Rate this question:

Neurobiology

□ □ □

17-00000

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	19
Responses Total:	19
Responses - % Correct:	0%

Back to Filters

Question 32 of 203

A 70-year-old lady is admitted via the emergency department with complaints of weakness, nausea and malaise. Unfortunately she has not brought all of her regular medications with her. She tells you that she is a tablet-controlled diabetic with high blood pressure and kidney problems. She has with her a packet of lisinopril which she takes 10 mg/day. In addition she has a packet of gliclazide which she takes 80 mg/day. She takes several other tablets which she left on the sideboard at home when the ambulance collected her. She attended an outpatient appointment 2 weeks ago where she was prescribed a new medication. She does not know the name of this medication and does not have it with her, she was told that it was a medication also used to treat people with liver problems and had been taking one tablet a day.

On examination she is overweight and appears unwell. She has evidence of mild biventricular cardiac insufficiency with elevated jugular venous pressure and pitting oedema of her ankles. The remainder of the examination is unremarkable.

An electrocardiogram (ECG) shows increased PR interval with widening of the QRS complex.

Urgent biochemical analysis shows:

Na ⁺	140 mmol/l
K ⁺	7.1 mmol/l
Urea	11.0 mmol/l
Creatinine	146 mol/l

Appropriate treatment is instigated.

The most likely cause for this lady’s presentation on this occasion is:

- ≡
- A

Recent commencement of digoxin 62.5 g od
- B

Recent commencement of spironolactone 50 mg
- C

A recent 2-day course of ibuprofen 400 mg tds/prn for backache
- D

Recent commencement of bisoprolol fumarate 2.5 mg od
- E

Recent commencement of amiloride hydrochloride 5 mg bd

7015

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Question 32 of 203

Urgent biochemical analysis shows:

Na ⁺	140 mmol/l
K ⁺	7.1 mmol/l
Urea	11.0 mmol/l
Creatinine	146 μmol/l

The most likely cause for this lady's presentation on this occasion is:

- | | |
|---|--|
| A | Recent commencement of digoxin 62.5 g od |
| B | Recent commencement of spironolactone 50 mg |
| C | A recent 2-day course of ibuprofen 400 mg tds/prn for backache |
| D | Recent commencement of bisoprolol fumarate 2.5 mg od |
| E | Recent commencement of amiloride hydrochloride 5 mg bd |

This lady has life-threatening hyperkalaemia with evidence of cardiotoxicity on electrocardiogram (ECG), requiring urgent intervention to prevent potentially fatal cardiac arrhythmia. All of the drugs listed can be associated with hyperkalaemia. However, we know that this lady has renal disease. In addition we know that she is already taking an angiotensin-converting enzyme (ACE) inhibitor. Following the Randomised Aldactone Evaluation Study (RALES) clinical trial when it was shown that aldosterone antagonists can reduce mortality in severe chronic congestive cardiac failure when given in conjunction with an ACE inhibitor, the prescription rates of spironolactone increased along with hospital admission rates and deaths from hyperkalaemia. Although this medication can be given safely with ACE inhibitors and indeed be beneficial, there is a risk of causing life-threatening hyperkalaemia particularly in those with renal impairment. The combination of ACE inhibitor therapy with spironolactone therefore needs to be carefully considered and monitored to avoid significant side-effects. Standard dose used for heart failure is at lower level than seen here (around 25-50mg daily). Digoxin toxicity can be complicated by hypokalaemia although it's hyperkalaemia which is seen here. Non-steroidal anti-inflammatory agents can be associated with hyperkalaemia in the context of renal impairment, although we might expect this with a longer duration of use if this were the cause in this case. Non-selective beta-blockade can lead to hyperkalaemia by decreasing uptake of potassium from extracellular to intracellular spaces. This would not be expected with the highly selective beta-1 blocker, bisoprolol. Amiloride can cause hyperkalaemia in similar situations, however it is not so commonly used in current practice.

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	20
Responses Total:	20
Responses - % Correct:	0%

Back to Filters

Question 33 of 203

A 16-year-old young man presents with long-standing polyuria, polydipsia, occasional abdominal and calf cramping pains along with the following biochemical abnormalities. His parents are first cousins. He has a normal stature and has been noted to have a low-normal blood pressure at around 95/50 mmHg.

Investigations reveal:

Na ⁺	136 mmol/l
K ⁺	2.3 mmol/l
Urea	4.5 mmol/l
Creatinine	80 μmol/l
Bicarbonate	34 mmol/l
Mg ²⁺	0.59 mmol/l
Urinary calcium excretion Low	(urinary calcium/creatinine ratio 0.1)

The likely diagnosis is:

- A

Alport’s syndrome
- B

Liddle’s syndrome
- C

Bartter’s syndrome
- D

Severe Cushing’s syndrome
- E

Gitelman’s syndrome

7016

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 33 of 203

Investigations reveal:

Na ⁺	136 mmol/l
K ⁺	2.3 mmol/l
Urea	4.5 mmol/l
Creatinine	80 μmol/l
Bicarbonate	34 mmol/l
Mg ²⁺	0.59 mmol/l
Urinary calcium excretion Low	(urinary calcium/creatinine ratio 0.1)

- | | |
|---|---------------------------|
| A | Alport's syndrome |
| B | Liddle's syndrome |
| C | Bartter's syndrome |
| D | Severe Cushing's syndrome |
| E | Gitelman's syndrome |

The patient has features of Gitelman's syndrome with hypokalaemic metabolic alkalosis, hypomagnesaemia, and hypocalciuria. His symptoms are relatively mild and his age at presentation and low-normal blood pressure all support the diagnosis of Gitelman's syndrome. This can occur either as a sporadic or autosomal recessive condition. The consanguinity in this case suggests that this may be an autosomal recessive case. Bartter's syndrome shares some features (including hypokalaemia and hypotension) with Gitelman's but is more severe and patients therefore present earlier with profound hypokalaemia and hypotension in early childhood. In Gitelman's urinary calcium excretion is low, and this can also be used as a differentiator. The prognosis of Gitelman's is very good, and treatment is with potassium and magnesium supplements. Alport's syndrome is an inherited condition in which an abnormal form of glomerular basement membrane is produced. It is associated with bilateral sensorineural deafness and renal failure. Liddle's syndrome is associated with hypokalaemic metabolic alkalosis but usually in the context of hypertension. Cushing's syndrome would be expected to produce other characteristic features as well as hypertension without the magnesium and calcium abnormalities seen above.

Rate this question:

Next Question

End Session

Peer Responses %

Session Progress	
Responses Correct:	0
Responses Incorrect:	21
Responses Total:	21
Responses - % Correct:	0%

Back to Filters

Question 34 of 203

A 70-year-old man is admitted with shortness of breath and peripheral oedema. He is oliguric. He has a history of diabetes associated with renal impairment and has been followed up in the renal clinic where he has been relatively stable until now. His pulse is regular, rate 96/min and his blood pressure 190/100 mmHg. His respiratory rate is 30/min, oxygen saturation 92% on high-flow oxygen. He has a raised jugular venous pulse and gross pitting oedema of both his ankles. Respiratory examination reveals coarse crackles to the mid zones of both lungs. He has a third heart sound on auscultation. His electrocardiogram (ECG) shows prolongation of the PR interval and widening of the QRS complex.

Urgent blood results are as follows:

Na ⁺	136 mmol/l
K ⁺	6.7 mmol/l
Urea	36 mmol/l
Creatinine	780 mol/l
Bicarbonate	11 mmol/l

The best series of treatments while awaiting transfer to the renal unit for dialysis is:

- A

15 units actrapid insulin in 50 ml 50% dextrose intravenous (iv) plus calcium gluconate
- B

15 units actrapid insulin in 50 ml 50% dextrose iv with 500 ml 1.26% sodium bicarbonate plus calcium gluconate
- C

Salbutamol 5 mg nebulised, calcium gluconate, 15 units actrapid insulin in 50 ml 50% dextrose iv, furosemide bolus iv with glyceryl trinitrate infusion plus calcium resonium 15 g orally
- D

Salbutamol 5 mg nebulised, calcium resonium 15 g orally, 15 units actrapid insulin in 50 ml 50% dextrose iv plus calcium gluconate
- E

Salbutamol 5 mg nebulised, calcium resonium 15 g orally, 15 units actrapid insulin in 50 ml 50% dextrose iv with 500 ml 1.26 % sodium bicarbonate, furosemide bolus iv with glyceryl trinitrate infusion plus calcium gluconate

7017

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 34 of 203

A 70-year-old man is admitted with shortness of breath and peripheral oedema. He is oliguric. He has a history of diabetes associated with renal impairment and has been followed up in the renal clinic where he has been relatively stable until now. His pulse is regular, rate 96/min and his blood pressure 190/100 mmHg. His respiratory rate is 30/min, oxygen saturation 92% on high-flow oxygen. He has a raised jugular venous pulse and gross pitting oedema of both his ankles. Respiratory examination reveals coarse crackles to the mid zones of both lungs. He has a third heart sound on auscultation. His electrocardiogram (ECG) shows prolongation of the PR interval and widening of the QRS complex.

Urgent blood results are as follows:

Na ⁺	136 mmol/l
K ⁺	6.7 mmol/l
Urea	36 mmol/l
Creatinine	780 mol/l
Bicarbonate	11 mmol/l

The best series of treatments while awaiting transfer to the renal unit for dialysis is:

- A

15 units actrapid insulin in 50 ml 50% dextrose intravenous (iv) plus calcium gluconate
- B

15 units actrapid insulin in 50 ml 50% dextrose iv with 500 ml 1.26% sodium bicarbonate plus calcium gluconate
- C

Salbutamol 5 mg nebulised, calcium gluconate, 15 units actrapid insulin in 50 ml 50% dextrose iv, furosemide bolus iv with glyceryl trinitrate infusion plus calcium resonium 15 g orally
- D

Salbutamol 5 mg nebulised, calcium resonium 15 g orally, 15 units actrapid insulin in 50 ml 50% dextrose iv plus calcium gluconate
- E

Salbutamol 5 mg nebulised, calcium resonium 15 g orally, 15 units actrapid insulin in 50 ml 50% dextrose iv with 500 ml 1.26 % sodium bicarbonate, furosemide bolus iv with glyceryl trinitrate infusion plus calcium gluconate

Explanation

This man has acute renal failure with evidence of fluid overload and pulmonary oedema. He also has life-threatening hyperkalaemia with electrocardiogram (ECG) changes. These two features constitute medical emergencies and need to be treated accordingly. Calcium gluconate needs to be given in view of its cardioprotective and stabilising effect, followed by attempts to lower the potassium with salbutamol, and carefully monitored dextrose/insulin infusion. Calcium resonium with take some time to work but may improve matters over the next few days. In view of his respiratory distress, nitrates and a bolus dose of furosemide will offload the heart and improve his pulmonary oedema. Sodium bicarbonate is of questionable value in this case, and the large fluid load would worsen the cardiopulmonary situation, particularly the pulmonary oedema.

7017

Rate this question:      

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	22
Responses Total:	22
Responses - % Correct:	0%

Question 35 of 203

Investigations from his last visit are as follows:

A 24-h urine collection was carried out which demonstrated 180 mg of albumin in the sample.

A	Commencing an angiotensin-converting enzyme (ACE) inhibitor in the first instance
B	Optimising glycaemic control
C	Optimising blood pressure control with other agents besides ACE inhibitors in the first instance
D	Commencing betablockade
E	Commencing angiotensin receptor blockers (ARB) in the first instance

[Previous Question](#)

Skip Question

Back to Filters

Question 35 of 203

A 65-year-old man with type 2 diabetes mellitus is seen in the diabetic clinic for follow-up. He has been diabetic for nearly 10 years now, diet controlled for 4 years and since that time on gliclazide and metformin. Unfortunately he has missed many of his follow-up appointments. His blood pressure is 124/74 mmHg. He has no evidence of peripheral neuropathy.

Investigations from his last visit are as follows:

Haemoglobin A _{1c} (HbA _{1c})	45.36 mmol/mol (6.3 %)
Protein	Negative
Glucose	Negative
Leucocytes	Negative
Nitrites	Negative
Na ⁺	137 mmol/l
K ⁺	3.8 mmol/l
Urea	6.8 mmol/l
Creatinine	98 μmol/l

A 24-h urine collection was carried out which demonstrated 180 mg of albumin in the sample.

The management step most likely to improve his overall prognosis with respect to renal and other complications is:

- A

Commencing an angiotensin-converting enzyme (ACE) inhibitor in the first instance

≡
- B

Optimising glycaemic control
- C

Optimising blood pressure control with other agents besides ACE inhibitors in the first instance
- D

Commencing betablockade
- E

Commencing angiotensin receptor blockers (ARB) in the first instance

Explanation ⚙

- A

Commencing an angiotensin-converting enzyme (ACE) inhibitor in the first instance

ACE inhibitors have been shown to improve outcome in microalbuminuria in diabetic renal disease and reduce progression to frank proteinuria and renal failure. Additionally, they show cardiovascular (CV) outcome benefits in large randomised controlled trials (RCTs). Their use in this context has therefore become widespread.

- B

Optimising glycaemic control

Optimising glycaemic control is likely to reduce microvascular complications as evidenced by the UK Prospective Diabetes Study (UKPDS). In this case however, there is little further optimisation which could be made based on the HbA_{1c} result given, and further attempts to improve the situation might well lead to unacceptable side-effects such as hypoglycaemia.

- C

Optimising blood pressure control with other agents besides ACE inhibitors in the first instance

Although blood pressure control is important in delaying renal progression, ACE inhibition, in particular, has been proven to improve outcome in microalbuminuria in diabetic renal disease and reduce progression to frank proteinuria and renal failure. Therefore, ACE inhibition should be used in the first instance.

- D

Commencing betablockade

There is no evidence that β blockade will improve renal prognosis aside from blood pressure control.

- E

Commencing angiotensin receptor blockers (ARB) in the first instance

There is growing evidence that ARBs have similar benefits to ACE inhibitors in heart failure and other indications, and they may have a lower incidence of side-effects. Whilst evidence of renoprotection from clinical trials is strong however, evidence of ischaemic CV outcome benefit is less clear. Just taking valsartan as one example: whilst a reduction in hospitalisations was seen in the main heart failure study, there was no reduction in mortality seen. For this reason, at the present time ACE inhibitors remain first-line in most cases.

7018

Rate this question: ☹ ★ ★ ★ ★ ★

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	23
Responses Total:	23
Responses - % Correct:	0%

Question 36 of 203

The results are as follows:

Na ⁺	140 mmol/l
K ⁺	6.9 mmol/l
Urea	4.0 mmol/l
Creatinine	82 μmol/l
C-reactive protein	139 mg/l
Glucose	5.8 mmol/l

The best course of action is:

- | | |
|---|---|
| A | Urgent calcium gluconate 10 ml 10% intravenous (iv) followed by insulin/dextrose infusion |
| B | Urgent electrocardiogram (ECG), followed by calcium gluconate 10 ml 10% iv followed by insulin/dextrose infusion |
| C | Repeat laboratory urea and electrolyte sample, and hold off further treatment until this result is returned |
| D | Repeat laboratory urea and electrolyte sample and check potassium on an arterial blood gas machine if available |
| E | Urgent ECG. Repeat laboratory urea and electrolyte sample and check potassium on an arterial blood gas machine if available in the meantime |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Question 36 of 203

The results are as follows:

You visit him on the ward. He is having a nebuliser and reading the newspaper.

A	Urgent calcium gluconate 10 ml 10% intravenous (iv) followed by insulin/dextrose infusion
B	Urgent electrocardiogram (ECG), followed by calcium gluconate 10 ml 10% iv followed by insulin/dextrose infusion
C	Repeat laboratory urea and electrolyte sample, and hold off further treatment until this result is returned
D	Repeat laboratory urea and electrolyte sample and check potassium on an arterial blood gas machine if available
E	Urgent ECG. Repeat laboratory urea and electrolyte sample and check potassium on an arterial blood gas machine if available in the meantime

not on any medications which would predispose him to high potassium; indeed, his regular salbutamol treatment is likely to give him low serum potassium by driving potassium into the cells. His urea and electrolytes are otherwise normal. His glucose is normal. The most likely cause of the high potassium in this case is either mild haemolysis not picked up in the laboratory, or the long use of a tourniquet during blood taking, leading to relative tissue acidosis when the sample was taken and therefore a high potassium. It is likely to be artefactual. However, it would be prudent to check for evidence of cardiac toxicity which is easy to do with an ECG. It would also be worth rechecking the potassium, and if a blood gas machine is available it may be worth rechecking the potassium locally pending the formal laboratory result.

As explained above, it is important to determine whether there is any cardiac toxicity and also to confirm hyperkalaemia.

Calcium gluconate is given if there are any cardiac electrical changes. The high potassium may be an artefact therefore it is important to check validity.

If hyperkalaemia is true, then cardiac toxicity could be fatal; therefore although a repeat sample is needed, an urgent ECG should be performed

As above, urgent ECG is needed to check for any cardiac toxicity.

Rate this question:

End Session

Peer Responses %

Back to Filters

Question 37 of 203

A 65-year-old man, a retired factory worker presents with some troublesome urinary symptoms over the last 2 months. He reports feeling the need to empty his bladder frequently and often feels he still needs to void immediately after having been to the toilet. He admits to some terminal dribbling. He denies any systemic symptoms. He was seen by his general practitioner (GP) who advised that he stop drinking tea and coffee, although this has had little effect on his symptoms. His GP referred him because he found blood on urinalysis. He has a history of hypertension and diabetes mellitus for which he takes bendrofluazide and metformin. He is a smoker who has smoked around 10 cigarettes per day since he was a teenager. On examination he appears well, blood pressure is 160/95 mmHg, with a normal abdominal examination. Rectal examination reveals a smooth, moderately enlarged prostate with preservation of the median sulcus. A urine sample appears normal to visual inspection.

Blood results are as follows:

Na ⁺	146 mmol/l
K ⁺	3.2 mmol/l
Urea	8.5 mmol/l
Creatinine	129 mol/l
Alanine aminotransferase	33 IU/l
Alkaline phosphatase	98 IU/l
Bilirubin	24 mol/l
Prostate-specific antigen	4 ng/l

Urinalysis reveals:

Blood	2+
Leucocytes	1+
Protein	1+
Nitrites	Negative

Urine microscopy demonstrates erythrocytes at more than 5 per high power field (HPF) (microscopic haematuria defined as >3 per HPF). No casts or crystals are seen. No microorganisms are found on Gram stain.

The diagnosis is likely to be:



- A

Benign
- B

Squamous cell carcinoma of the bladder
- C

Prostatic carcinoma
- D

Transitional cell carcinoma of the bladder
- E

Non-infectious haemorrhagic cystitis

7020

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Question 37 of 203

Blood results are as follows:

Na ⁺	146 mmol/l
K ⁺	3.2 mmol/l
Urea	8.5 mmol/l
Creatinine	129 μmol/l
Alanine aminotransferase	33 IU/l
Alkaline phosphatase	98 IU/l
Bilirubin	24 μmol/l
Prostate-specific antigen	4 ng/l

Urinalysis reveals:

Blood	2+
Leucocytes	1+
Protein	1+
Nitrites	Negative

Urine microscopy demonstrates erythrocytes at more than 5 per high power field (HPF) (microscopic haematuria defined as >3 per HPF). No casts or crystals are seen. No microorganisms are found on Gram stain.

The diagnosis is likely to be:

A	Benign
B	Squamous cell carcinoma of the bladder
C	Prostatic carcinoma
D	Transitional cell carcinoma of the bladder
E	Non-infectious haemorrhagic cystitis

Explanation

The likely diagnosis is transitional cell carcinoma of the bladder. This man is a lifelong smoker and his work in a factory may have exposed him to recognised carcinogens associated with urothelial malignancy. The combination of symptoms suggesting bladder irritation and persistent albeit microscopic haematuria makes this a concern. Persistent microscopic haematuria needs to be taken seriously and investigated appropriately. Transitional cell carcinoma is much more common than squamous cell carcinoma. His symptoms are not suggestive of prostatic hypertrophy, although mild prostatic enlargement is common and might be found coincidentally. The normal rectal examination and low prostate-specific antigen make a diagnosis of prostatic carcinoma less likely, although it would still be on the differential and would need to be excluded. Non-infectious haemorrhagic cystitis is commonly associated with a specific insult to the bladder (such as metabolites of cyclophosphamide or radiotherapy of the pelvis and lower abdomen) none of which are reported in this case.

7020

[Next Question](#)

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	25
Responses Total:	25
Responses - % Correct:	0%

Back to Filters

Question 38 of 203

A 40-year-old lady is admitted to the medical assessment unit with a history of fatigue, headache and leg oedema. She also complains of pruritus and heartburn.

On examination she has slight oedema of the fingers and some perioral telangiectasia. Her mouth is dry and her jugular venous pressure (JVP) is slightly elevated. Auscultation of the lungs reveals fine bibasal end inspiratory crepitations. Her pulse is 105/min, respiratory rate 20/min, blood pressure 195/95 mmHg. Urine dipstick shows proteinuria and blood film shows helmet cells.

Given the likely diagnosis, what treatment is likely to have the most favourable outcome?

- A

Beta-blocker
- B

Nitroprusside
- C

Glucocorticoid
- D

Angiotensin-converting enzyme inhibitor
- E

Cyclophosphamide

7108

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 38 of 203

A 40-year-old lady is admitted to the medical assessment unit with a history of fatigue, headache and leg oedema. She also complains of pruritus and heartburn. On examination she has slight oedema of the fingers and some perioral telangiectasia. Her mouth is dry and her jugular venous pressure (JVP) is slightly elevated. Auscultation of the lungs reveals fine bibasal end inspiratory crepitations. Her pulse is 105/min, respiratory rate 20/min, blood pressure 195/95 mmHg. Urine dipstick shows proteinuria and blood film shows helmet cells.

Given the likely diagnosis, what treatment is likely to have the most favourable outcome?

- A

Beta-blocker
- B

Nitroprusside
- C

Glucocorticoid
- D

Angiotensin-converting enzyme inhibitor
- E

Cyclophosphamide

Explanation ⚙

- D

Angiotensin-converting enzyme inhibitor

The history and examination give features of scleroderma renal crisis. Histologically interlobular arteries are affected with intimal thickening. Features include malignant hypertension, oliguria, fluid retention, and microangiopathic haemolytic anaemia. ACE inhibitors will significantly reduce progression to renal failure and increase the chance of recovery if renal failure has already developed.

- A

Beta-blocker

A β -blocker may help control blood pressure, but it will not help control renal scleroderma renal crisis. It may make any Raynaud's phenomenon worse.

- B

Nitroprusside

An excessive reduction in pressure and hypovolaemia should be avoided in renal crisis because both can further diminish renal perfusion and superimpose acute tubular necrosis upon the lesions of scleroderma. Thus, parenteral antihypertensive agents (such as intravenous nitroprusside or labetalol) should be avoided, if possible, as should other nephrotoxins such as nonsteroidal anti-inflammatory drugs and radiocontrast agents.

- C

Glucocorticoid

Similarly, glucocorticoids are effective immunosuppressants; however, they are not used for the treatment of scleroderma. A side-effect of steroids is dyspepsia and may make heartburn symptoms worse.

- E

Cyclophosphamide

Cyclophosphamide is an immunosuppressant. It has not been shown to be an effective treatment for scleroderma renal crisis.

7108

Rate this question: ⊖ ★ ★ ★ ★ ★

Next Question

- Previous Question

Feedback
- Tag Question

End Session

Difficulty: Difficult

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	26
Responses Total:	26
Responses - % Correct:	0%

Back to Filters

Question 39 of 203

A 43-year-old gentleman is brought into hospital during the night. He was found on the street smelling strongly of alcohol. Staff recall that he also has a history of intravenous drug use. He is disorientated, localises pain and will open his eyes in response to a painful stimulus. Reflexes are symmetrically brisk. Pupils are 2 mm in diameter. His pulse is 110/min, blood pressure 95/60 mmHg, respiratory rate 12/min, oxygen saturation 97% on air

Initial Investigations reveal:

Electrocardiogram	Sinus tachycardia, inverted p wave aVr
Chest X-ray	Hyperinflated lung fields, old healed rib fractures
Urine dipstick	Protein +, blood +++
Haemoglobin	14g/dl
White cell count	11.5 × 10 ⁹ /l
Platelets	420 × 10 ⁹ /l
Sodium	135 mmol/l
Potassium	5.7 mmol/l
Urea	10.8 mmol/l
Creatinine	278 mol/l
Calcium	2.03 mmol/l
Phosphate	1.69 mmol/l (0.8-1.4)

What appropriate step should be taken next?

- A

Intravenous calcium gluconate
- B

Vitamin D replacement therapy
- C

Catheterise and intravenous saline
- D

Intravenous antibiotics
- E

Intravenous naloxone

7109

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 39 of 203

-

- The diagnosis is rhabdomyolysis. The patient has high risk factors with alcohol and intravenous drug use. He has most likely been immobile on the street, or could have suffered injury while intoxicated. Inverted P waves in aVR are usually normal. Myoglobinuria gives a 'false-positive' on urine dipstick for blood. Tests reveal typically hyperkalaemia, hypocalcaemia, raised creatinine:urea ratio, hyperphosphataemia. Further confirmatory investigations include urine myoglobin, serum creatine kinase and serum myoglobin. Fluid resuscitation is needed with urine output measurement. He does not have symptoms of opioid overdose which are sufficiently severe enough to require naloxone, and intravenous calcium can increase the destruction of muscle. The hyperkalaemia needs monitoring but no active treatment is needed currently as fluid hydration will help.

- Calcium gluconate helps stabilise cardiac membrane in hyperkalaemia. In this case rehydration will help hyperkalaemia. The latter stages of rhabdomyolysis may result in hypercalcaemia as calcium is released from the muscles. This is more likely to occur if supplementary calcium has been given.

- During later stages of rhabdomyolysis, the accumulated calcium is released from the storage sites. This is often associated with hyperparathyroidism and hypervitaminosis D and overt hypercalcaemia. However, the hyperparathyroidism and hypervitaminosis D are not seen in all cases. Therefore, vitamin D is not a treatment.

- There is no evidence of any infection. The white cell count is

- During later stages of rhabdomyolysis, the accumulated calcium is released from the storage sites. This is often associated with hyperparathyroidism and hypervitaminosis D and overt hypercalcaemia. However, the hyperparathyroidism and hypervitaminosis D are not seen in all cases. Therefore, vitamin D is not a treatment.

Rate this question:

Next Question

Tag Question

End Session

Peer Responses %

Session Progress

Back to Filters

Question 40 of 203

A 30-year-old woman was admitted yesterday with sudden onset breathlessness while eating dinner and associated cough with frothy pink sputum. She was treated in the emergency department and has been transferred to the ward. A similar episode prompted admission to hospital 2 months ago. She has a background medical history of hay fever. Her chest on auscultation is clear and she feels a lot better. Her blood pressure is 160/90 mmHg. The remainder of the examination is unremarkable.

Investigations on admission:

Electrocardiogram	Sinus rhythm at 120/min
Chest X-ray	Upper lobe diversion, Kerley B lines, cardiothoracic ratio of 0.4
Sodium	140 mmol/l
Potassium	3.8 mmol/l
Urea	7.5 mmol/l
Creatinine	110 μmol/l

What is the most likely underlying diagnosis?

- A

Wegener’s granulomatosis
- B

Conn’s syndrome
- C

Fibromuscular dysplasia
- D

Goodpasture’s syndrome
- E

Aortic stenosis

7110

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 40 of 203

Investigations on admission:

Electrocardiogram	Sinus rhythm at 120/min
Chest X-ray	Upper lobe diversion, Kerley B lines, cardiothoracic ratio of 0.4
Sodium	140 mmol/l
Potassium	3.8 mmol/l
Urea	7.5 mmol/l
Creatinine	110 μ mol/l

- | | |
|---|--------------------------|
| A | Wegener's granulomatosis |
| B | Conn's syndrome |
| C | Fibromuscular dysplasia |
| D | Goodpasture's syndrome |
| E | Aortic stenosis |

- | | |
|---|-------------------------|
| C | Fibromuscular dysplasia |
|---|-------------------------|

A Wegener's granulomatosis

B Conn's syndrome

D	Goodpasture's syndrome
---	------------------------

E	Aortic stenosis
---	-----------------

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	28
Responses Total:	28
Responses % Correct:	0%

Question 41 of 203

Tests show:

Blood pressure	145/80mmHg
Urine dipstick	++protein, ++blood
Fundoscopy	Bilateral yellow dots around the macula

- | | |
|---|--|
| A | Rapidly progressive glomerulonephritis |
| B | Mesangiocapillary glomerulonephritis |
| C | Primary focal segmental glomerulosclerosis |
| D | Membranous glomerulonephritis |
| E | Alport's syndrome |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 41 of 203

A general practitioner has referred a 20-year-old patient to the renal clinic with haematuria. The patient recalls having coryzal symptoms 2 weeks ago and then developing frank haematuria for a few days. He says the symptoms have resolved. The patient recalls his maternal grandfather had ‘kidney’ problems at a young age. He has a past history of hearing difficulties and wears hearing aids, and is under the ophthalmologists for treatment of corneal ulcers.

Tests show:

Blood pressure	145/80mmHg
Urine dipstick	++protein, ++blood
Fundoscopy	Bilateral yellow dots around the macula

Evidence of what underlying diagnosis is likely to be seen on the renal biopsy?

- A

Rapidly progressive glomerulonephritis
- B

Mesangiocapillary glomerulonephritis
- C

Primary focal segmental glomerulosclerosis
- D

Membranous glomerulonephritis
- E

Alport’s syndrome

Explanation ⚙

- E

Alport’s syndrome

Alport’s syndrome consists of: (1) nephritis causing microscopic haematuria and progressive renal failure with episodic frank haematuria; (2) ocular pathology – corneal ulcerations, bilateral anterior lenticonus, bilateral dot-fleck marks around the fovea; (3) progressive high-frequency sensorineuronal deafness. A light microscope examination can be unremarkable and electron microscope examination is needed. Occasionally focal and segmental glomerulosclerosis, interstitial fibrosis, tubular atrophy, and infiltration of lymphocytes and plasma cells may be seen on light microscopy. The syndrome is inherited mainly as X-linked.

- A

Rapidly progressive glomerulonephritis

This is a syndrome consisting of a rapid loss in renal function (usually a 50% decline in glomerular filtration rate within 3 months) with glomerular crescent formation in at least 50–75% of glomeruli on renal biopsy. If left untreated it rapidly progresses into acute renal failure.

- B

Mesangiocapillary glomerulonephritis

MPGN is an uncommon cause of chronic nephritis in children and young adults. There is a thickening of the peripheral capillary walls by subendothelial immune deposits and or intramembranous dense deposits.

- C

Primary focal segmental glomerulosclerosis

FGS is a common cause of nephrotic range proteinuria in adults. There would not be haematuria on dipstick.

- D

Membranous glomerulonephritis

This is the commonest cause of nephrotic syndrome (proteinuria, oedema and hypoalbuminaemia) in adults. There would not be blood in membranous on dipstick.

7111

Rate this question: ⊖ ★ ★ ★ ★ ★

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Difficult

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	29
Responses Total:	29
Responses - % Correct:	0%

Question 42 of 203

Initial investigations show:

Blood pressure	115/65mmHg
Haemoglobin	13.5 g/dl
White cell count	$8 \times 10^9/l$
Platelets	$320 \times 10^9/l$
Immunoglobulin G (IgG)	7.3 g/l (6.0-13.0)
Ig A	2.8 g/l (0.8-3.0)
Ig M	0.6 g/l (0.4-2.5)
Sodium	143 mmol/l
Potassium	3.9 mmol/l
Urea	8.5 mmol/l
Creatinine	115 μ mol/l
Bilirubin	30 μ mol/l
Aspartate aminotransferase	10 U/l
Alkaline phosphatase	60 U/l
Albumin	43 g/l
24-h urinary protein collection	1.8g/24h (<0.2g)
Urine microscopy	Red cell casts present
Renal biopsy light microscopy	Mesangial proliferation

3

- | | |
|---|------------------------------------|
| A | Dapsone side effect |
| B | Immunoglobulin A (IgA) nephropathy |
| C | Cryoglobulinaemia |
| D | Amyloidosis |
| E | Renal cell carcinoma |

Submit

Skip Question

Back to Filters

Question 42 of 203

A 26-year-old male is referred to you by his general practitioner with persistent microscopic haematuria. He has recently been diagnosed with coeliac disease and commenced on a gluten-free diet. The dermatologists have also started oral dapsone. The patient otherwise has remained asymptomatic.

Initial investigations show:

Blood pressure	115/65mmHg
Haemoglobin	13.5 g/dl
White cell count	8 × 10 ⁹ /l
Platelets	320 × 10 ⁹ /l
Immunoglobulin G (IgG)	7.3 g/l (6.0-13.0)
Ig A	2.8 g/l (0.8-3.0)
Ig M	0.6 g/l (0.4-2.5)
Sodium	143 mmol/l
Potassium	3.9 mmol/l
Urea	8.5 mmol/l
Creatinine	115 μmol/l
Bilirubin	30 μmol/l
Aspartate aminotransferase	10 U/l
Alkaline phosphatase	60 U/l
Albumin	43 g/l
24-h urinary protein collection	1.8g/24h (<0.2g)
Urine microscopy	Red cell casts present
Renal biopsy light microscopy	Mesangial proliferation

What is the most likely diagnosis?

- A

Dapsone side effect
- B

Immunoglobulin A (IgA) nephropathy
- C

Cryoglobulinaemia
- D

Amyloidosis
- E

Renal cell carcinoma

Explanation

Immunoglobulin A (IgA) nephropathy (Berger’s disease) is a mesangioproliferative glomerulonephritis. It usually presents with persistent microscopic haematuria or recurrent macroscopic haematuria usually associated with an upper respiratory tract infection. IgA levels are raised in 50% of cases. There is an association with cirrhosis and coeliac disease. Dapsone causes haemolytic anaemia. Cryoglobulinaemia usually presents with more cutaneous and articular manifestations. It is associated with haematological malignancies and connective tissue diseases. Amyloid gives rise to proteinuria and the nephrotic syndrome. Renal cell carcinoma arises from tubular epithelium.

7114

Rate this question: ⚙️ ⭐️ ⭐️ ⭐️ ⭐️ ⭐️

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	30
Responses Total:	30
Responses - % Correct:	0%

Question 43 of 203

Blood results show:

Today		Yesterday	
Sodium	138 mmol/l	Sodium	139 mmol/l
Potassium	4.3 mmol/l	Potassium	4.4 mmol/l
Urea	19.4 mmol/l	Urea	12.1 mmol/l
Creatinine	210 μmol/l	Creatinine	129 μmol/l
Bicarbonate	16 mmol/l	Bicarbonate	26 mmol/l

What is the most appropriate next management step?

- | | |
|---|--|
| A | Intravenous fluids with furosemide to encourage diuresis |
| B | Intravenous fluids and omit metformin, lisinopril |
| C | Sodium bicarbonate |
| D | Referral for dialysis |
| E | Intravenous antibiotics |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 43 of 203

A 49-year-old diabetic patient is on the ward for investigation of abdominal pain. You are asked to see her blood results on the computer during the night shift. Review of her medical notes on the ward reveals she has a background of nephropathy, neuropathy and ischaemic heart disease. She is currently taking gliclazide, metformin, atorvastatin, lisinopril, aspirin, and bisoprolol. While on the ward she developed a urinary tract infection and was commenced on trimethoprim. She is being investigated for abdominal pain and recently had an abdominal ultrasound scan and contrast-enhanced computerised tomography performed, which has revealed a solitary left adrenal nodule.

Blood results show:

Today		Yesterday	
Sodium	138 mmol/l	Sodium	139 mmol/l
Potassium	4.3 mmol/l	Potassium	4.4 mmol/l
Urea	19.4 mmol/l	Urea	12.1 mmol/l
Creatinine	210 μmol/l	Creatinine	129 μmol/l
Bicarbonate	16 mmol/l	Bicarbonate	26 mmol/l

Her trimethoprim dose is reduced on account of the creatinine rise. What is the most appropriate next management step?

- A

Intravenous fluids with furosemide to encourage diuresis
- B

Intravenous fluids and omit metformin, lisinopril
- C

Sodium bicarbonate
- D

Referral for dialysis
- E

Intravenous antibiotics

Explanation

The answer is - Intravenous fluids and omit metformin, lisinopril

The patient has worsening renal impairment and is developing an acidosis. Most likely this is due to contrast-induced nephropathy combined with her medications. She is at risk of metformin-induced lactic acidosis and her medications would have to be omitted. The effects of this would not be immediate and she would need hydration with intravenous fluids and close monitoring of her renal function. As this is potentially reversible she should not require dialysis at this time.



7121

Rate this question:

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	31
Responses Total:	31
Responses - % Correct:	0%

Question 44 of 203

Blood results are as follows:

Haemoglobin (Hb)	8.7 g/dl
Mean corpuscular volume (MCV)	90 fl
White cell count (WCC)	8.4 x 10 ⁹ /l-normal differential
Platelets	99 x 10 ⁹ /l
Blood film	Schistocytes, increased reticulocytes and helmet cells

Na ⁺	139 mmol/l
K ⁺	5.2 mmol/l
Urea	15 mmol/l
Creatinine	280 mmol/l
Bicarbonate	15 mmol/l

Na ⁺	137
K ⁺	5.1
Urea	9 mmol/l
Creatinine	110 mmol/l
Bicarbonate	20 mmol/l

International normalised ratio	1.2 (normal range 0.9-1.3)
Activated partial thromboplastin time (APTT)	45 s (normal range 35-45)

be.

- | | |
|---|---|
| A | Stopping the prostacyclin infusions |
| B | Cyclophosphamide |
| C | High-dose corticosteroids |
| D | Angiotensin-converting enzyme (ACE) inhibitor |
| E | Plasma exchange |

Submit

Skip Question

Question 44 of 203

Blood results are as follows:

Results from today:

Results from six weeks ago:

The most beneficial treatment over the longer term is likely to be.

Explanation

Next Question

Difficulty: Average

Peer Responses %

Conclusions

[Blog](#) [About Pastest](#) [Contact Us](#) [Help](#)

Back to Filters

Question 45 of 203

An overweight 43-year-old man with stable bipolar affective disorder treated with lithium carbonate is seen for the second time in the general medical outpatients clinic. He was originally referred from the psychiatric team with polyuria associated with excessive thirst and drinking. The psychiatric nurse who has been seeing him in the community wondered whether he might be suffering from diabetes mellitus. A capillary blood glucose measurement was normal however.

His results from the last visit are as follows:

Fasting blood glucose	6.9 mmol/l
K ⁺	3.9 mmol/l
Serum osmolality	309 mosmol/kg

The most useful management step is:

-
- A

Commencement of metformin 500 mg bd
- B

Withdrawal of lithium carbonate
- C

Admission for a water deprivation test with ADH administration
- D

Advice to the patient to reduce fluid intake, with explanation
- E

Admission for further investigation of pituitary function

7287

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 45 of 203

An overweight 43-year-old man with stable bipolar affective disorder treated with lithium carbonate is seen for the second time in the general medical outpatients clinic. He was originally referred from the psychiatric team with polyuria associated with excessive thirst and drinking. The psychiatric nurse who has been seeing him in the community wondered whether he might be suffering from diabetes mellitus. A capillary blood glucose measurement was normal however.

His results from the last visit are as follows:

Fasting blood glucose	6.9 mmol/l
K ⁺	3.9 mmol/l
Serum osmolality	309 mosmol/kg

The most useful management step is:

-
- A

Commencement of metformin 500 mg bd
- B

Withdrawal of lithium carbonate
- C

Admission for a water deprivation test with ADH administration
- D

Advice to the patient to reduce fluid intake, with explanation
- E

Admission for further investigation of pituitary function

Explanation

This man is likely to have nephrogenic diabetes insipidus as a result of lithium therapy. He has a borderline-high serum osmolality. However, an alternative cause has not been excluded by the investigations carried out so far, and a water deprivation test will clarify the cause and enable confident and accurate management. It may be appropriate to consider stopping the lithium therapy. However, in view of his serious psychiatric diagnosis which has a significant mortality associated with poor control of symptoms, it would be unwise to stop this medication without psychiatric review unless there are life-threatening and urgent indications necessitating treatment withdrawal. His fasting blood sugar suggests that he has impaired fasting glucose (IFG). This is unlikely to cause his symptoms and would be best treated at this stage with dietary advice, encouragement to lose weight and increase exercise rather than oral hypoglycaemic medication. In view of the serum osmolality, the possibility of psychogenic polydipsia, which must also be considered in this situation, is rejected. In this condition the osmolality would be normal or low and not high. It is too early to say whether investigation of pituitary function would be appropriate (for example if cranial diabetes insipidus was suspected). Again, the water deprivation test should clarify this.

7287

Rate this question:

Next Question

- Previous Question

Feedback
- Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	33
Responses Total:	33
Responses - % Correct:	0%

Back to Filters

Question 46 of 203

As the renal registrar, you attend a joint meeting with the renal and pathology departments to discuss the pathology of some of the renal patients. The histopathologist displays a slide and asks you to describe it. Your description is as follows: ‘Section of a kidney showing glomeruli, most of which appear to be filled with a pink homogenous substance. There appears to be an inflammatory infiltrate in the interstitial tissues consisting mainly of lymphocytes.’

Being rather impressed with your description, the histopathologist asks you for your diagnosis. What would be your answer?

- | | |
|---|---|
| A | Diffuse diabetic glomerulosclerosis |
| B | Toxic acute tubular necrosis (TATN) |
| C | Acute pyelonephritis |
| D | Acute diffuse proliferative glomerulonephritis (ADPG) |
| E | Chronic glomerulonephritis |

7609

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 46 of 203

As the renal registrar, you attend a joint meeting with the renal and pathology departments to discuss the pathology of some of the renal patients. The histopathologist displays a slide and asks you to describe it. Your description is as follows: ‘Section of a kidney showing glomeruli, most of which appear to be filled with a pink homogenous substance. There appears to be an inflammatory infiltrate in the interstitial tissues consisting mainly of lymphocytes.’

Being rather impressed with your description, the histopathologist asks you for your diagnosis. What would be your answer?

- A

Diffuse diabetic glomerulosclerosis
- B

Toxic acute tubular necrosis (TATN)
- C

Acute pyelonephritis
- D

Acute diffuse proliferative glomerulonephritis (ADPG)
- E

Chronic glomerulonephritis

Explanation

Chronic glomerulonephritis is an end-stage condition of nearly all forms of acute glomerulonephritis, and affects the majority of the glomeruli. This diagnosis can be made without knowing the cause.

The glomeruli will show different degrees of hyalinisation, and eventually hyalinosclerosis will occur whereby all the glomeruli and Bowman’s space are replaced by hyaline. Hyaline is a homogenous, amorphous pink material and glomeruli that are totally hyalinised are atrophic and non-functional.

An abundant inflammatory infiltrate, consisting mainly of lymphocytes, is present in the interstitial tissues. The remaining functional nephrons have dilated tubules.

In ADPG, there is proliferation of endothelial and mesangial cells but the tubules are not affected.

In acute pyleonephritis, a purulent exudate is present in the interstitium. The glomeruli are normal in the early stages. In diabetic glomerulosclerosis, a protein is deposited in the basement membrane of the capillary loops of the glomeruli, resulting in basement membrane thickening. TATN is caused by toxic damage to the proximal tubular epithelium causing necrosis. The basement membrane is intact.

7609

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Difficult

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	1
Responses Total:	1
Responses - % Correct:	0%

Question 47 of 203

Results are shown below:

Plasma osmolality	295 mOsmol/kg (275-290mOsmol/kg)
Urine osmolality	300 mOsmol/kg
Sodium	147 mmol/l
Potassium	5.0 mmol/l
Urea	7.5 mmol/l
Creatinine	105 μ mol/l
Glucose	5.8 mmol/l
9 am cortisol	400 nmol/l (280-700nmol/l)



- | | |
|---|---|
| A | Fluid restriction up to 1000 ml per day |
| B | Intravenous hydrocortisone 200 mg |
| C | Match fluid output with appropriate replacement |
| D | Immediate intranasal DDAVP |
| E | Encourage the patient to drink as much as they want |

Submit

Skip Question

Back to Filters

Question 47 of 203

A 28-year-old female presented to the emergency department with a 2-month history of headache and visual disturbance. Urgent magnetic resonance imaging of the head revealed a pituitary adenoma. Baseline pituitary function tests were normal except serum prolactin level of 15000 mU/l (<450 mU/l) Transphenoidal surgery (TSS) 2 days after admission was successful. However in the first 6 h post-surgery her urine output was 2.5 l. Otherwise she was well with no particular symptoms or complications.

Urgent biochemical test were performed.

Results are shown below:

Plasma osmolality	295 mOsmol/kg (275–290mOsmol/kg)
Urine osmolality	300 mOsmol/kg
Sodium	147 mmol/l
Potassium	5.0 mmol/l
Urea	7.5 mmol/l
Creatinine	105 μmol/l
Glucose	5.8 mmol/l
9 am cortisol	400 nmol/l (280–700nmol/l)

What is the most appropriate initial therapy?

- A

Fluid restriction up to 1000 ml per day
- B

Intravenous hydrocortisone 200 mg
- C

Match fluid output with appropriate replacement
- D

Immediate intranasal DDAVP
- E

Encourage the patient to drink as much as they want

Explanation

Diabetes insipidus (DI) post-pituitary surgery is very common. All patients should be monitored for signs of DI such as urine output >800 ml/hr over four consecutive hours. The symptom of thirst is unusual. Urgent paired osmolality and electrolytes should be requested if there is any suspicion of DI.

Plasma osmolality >285 mOsmol/kg with paired urine osmolality <500 mOsmol/kg suggests DI. This may settle a few days after surgery though, so in the initial period ensuring adequate fluid replacement is reasonable. Only some patients will require long term DDAVP replacement.

8469

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	2
Responses Total:	2
Responses - % Correct:	0%

Question 48 of 203

Venous blood results

Na ⁺	142 mmol/l
K ⁺	1.8 mmol/l
Bicarbonate	13 mmol/l
Chloride	117 mmol/l

pH	7.26
p(CO ₂)	3.0 kPa

- | | |
|---|---------------------------------|
| A | Distal renal tubular acidosis |
| B | Proximal renal tubular acidosis |
| C | Type 4 renal tubular acidosis |
| D | Bartter's syndrome |
| E | Diabetic ketoacidosis |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Question 48 of 203

Venous blood results

Na ⁺	142 mmol/l
K ⁺	1.8 mmol/l
Bicarbonate	13 mmol/l
Chloride	117 mmol/l

Arterial blood gas

pH	7.26
p(CO ₂)	3.0 kPa

The most likely diagnosis and cause for the above blood results is:

A	Distal renal tubular acidosis
B	Proximal renal tubular acidosis
C	Type 4 renal tubular acidosis
D	Bartter's syndrome
E	Diabetic ketoacidosis

Explanation



A	Distal renal tubular acidosis
---	-------------------------------

The information given is ideal to work out the anion gap:

$$(\text{Na}^+ + \text{K}^+) - (\text{Cl}^- + \text{HCO}_3^-) = (142 + 1.8) - (117 + 13) = 13.8$$

Hence we are looking at a normal anion gap acidosis with hypokalaemia. In distal (type 1) renal tubular acidosis there is failure of hydrogen excretion by the distal tubule of the nephron, with associated hypokalaemia. Of the available options this is most likely as there is an association with sickle cell anaemia, and these patients can get renal stones.

B Proximal renal tubular acidosis

In proximal (type 2) renal tubular acidosis there is failure of bicarbonate reabsorption in the proximal tubule of the nephron, usually occurring in infancy.

C Type 4 renal tubular acidosis

Type 4 renal tubular acidosis is also known as hyporeninaemic hypoaldosteronism. This condition would cause acidosis and hyperkalaemia.

D Bartter's syndrome

In this condition there is hyperplasia of the juxtaglomerular apparatus and biochemically there are hypokalaemia, alkalosis and hypercalciuria, with a normal blood pressure.

E	Diabetic ketoacidosis
---	-----------------------

This causes an elevated anion gap because of the acidic ketones in the blood

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	3
Responses Total:	3
Responses - % Correct:	0%

Question 49 of 203

Investigation results are below:

She was commenced on treatment with high-dose diuretics and immunosuppressant therapy. Several weeks later she was readmitted with abdominal pain. On examination she had a mild pyrexia and was tender in both loins.

Repeat investigation results are below:

What is the most likely cause for the acute deterioration in renal function?

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Question 49 of 203

(Hb)

She was commenced on treatment with high-dose diuretics and immunosuppressant therapy. Several weeks later she was readmitted with abdominal pain. On examination she had a mild pyrexia and was tender in both loins.

Repeat investigation. Results are below:

What is the most likely cause for the acute deterioration in renal function?

© 2011 Pearson Education, Inc. All rights reserved.

[illegible]

Explanation

D Renal

systemic lupus erythematosus

renal vein thrombosis. This occurs in 10–20% of patients with nephrotic syndrome and is more common in those with membranous glomerulonephritis. Patients with nephrotic syndrome are in a hypercoagulable state due to a combination of intravascular volume depletion (exacerbated by diuretics) and the loss of clotting factors in the urine (particularly antithrombin III), and are therefore at high risk of thromboses. Renal vein thrombosis may be silent, causing gradually worsening renal impairment, or can present acutely as in this case. Diagnosis is with Doppler ultrasound of the renal veins, computed tomography (CT)/magnetic resonance imaging (MRI) or venography. Management includes mobilisation, avoidance of volume depletion and long-term anticoagulation. The imaging results are not described here but the bilateral signs and symptoms do suggest involvement of both kidneys.

Bilateral renal vein thrombosis, although less common than the

unilateral form, can occur as an acute phenomenon and is an indication for more aggressive management including, if necessary, thrombolysis.

Pyelonephritis would tend to

dysuria and haematuria; the clinical picture of nephritic syndrome would not be typical and is much better explained here by a diagnosis of renal vein thrombosis.

Dealing with the data

erythematosus (SLE), and it does not present with abdominal pain and fever. The common presentations of renal artery stenosis would be medically resistant hypertension, flash pulmonary oedema or a gradual worsening of renal impairment.

C	Renal infarction
---	------------------

Renal infarction may give rise to acute abdominal pain and haematuria. It would be unusual to have renal infarction of both kidneys to give bilateral loin pain.

Renal calculus	
----------------	--

Although renal calculus can present with abdominal pain and, if there is associated infection, there may be fever with haematoproteinuria. However, there is no risk factor to suggest renal calculus such as dehydration, hypercalcaemia or hyperuricaemia for example. Normally the pain would be on one side rather than bilateral loin pain.

Rate this question:

100

[Previous Question](#)

Tag Question

Feedback

Extra Session

100

Responses Incorrect:	4
Responses Total:	4
Responses - % Correct:	0%

Back to Filters

Question 50 of 203

A 24-year-old woman was admitted to hospital with a 3-day history of feeling generally unwell, with fatigue, arthralgia and pruritis. She had recently finished a 5-day course of antibiotics for a urinary tract infection but there was no other significant past medical history. She had no significant findings on clinical examination except for a widespread erythematous rash.

Investigation results are below:

Hb	12.6 g/dl
WCC	13.0 × 10 ⁹ /l (eosinophilia)
PLT	390 × 10 ⁹ /l
Creatinine	720 μmol/l
Na ⁺	135 mmol/l
K ⁺	5.2 mmol/l
Urea	22.0 mmol/l
Urinalysis	Protein ++ blood +

What is the most important investigation to establish the diagnosis?



- A

Autoimmune profile
- B

Renal tract ultrasound scan
- C

Renal biopsy
- D

Urine microscopy and culture
- E

Antistreptolysin-O titre

9159

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Question 50 of 203

Investigation results are below:

Hb	12.6 g/dl
WCC	13.0 × 10 ⁹ /l (eosinophilia)
PLT	390 × 10 ⁹ /l
Creatinine	720 μmol/l
Na ⁺	135 mmol/l
K ⁺	5.2 mmol/l
Urea	22.0 mmol/l
Urinalysis	Protein ++ blood +

- | | |
|---|------------------------------|
| A | Autoimmune profile |
| B | Renal tract ultrasound scan |
| C | Renal biopsy |
| D | Urine microscopy and culture |
| E | Antistreptolysin-O titre |

- ## C Renal biopsy

A Autoimmune profile

B Renal tract ultrasound scan

D Urine microscopy and culture

E Antistreptolysin-O titre

9159

[Next Question](#)

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Section D

Responses Correct:	0
Responses Incorrect:	5
Responses Total:	5
Responses - % Correct:	0%

Question 51 of 203

His blood results are below:

Haemoglobin (Hb)	15 g/dl
White cell count	$5.0 \times 10^9/l$
Platelets	$200 \times 10^9/l$
Creatinine	90 μ mol/l
Na ⁺	136 mmol/l
K ⁺	3.8 mmol/l
Urea	5.2 mmol/l

3

- | | |
|---|---|
| A | Start angiotensin-converting enzyme inhibitor |
| B | Confirm microalbuminuria with a repeat test in 3 months |
| C | Start angiotensin II inhibitor |
| D | Commence low-protein diet |
| E | Start calcium channel antagonist |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Question 51 of 203

His blood results are below:

Haemoglobin (Hb)	15 g/dl
White cell count	$5.0 \times 10^9/l$
Platelets	$200 \times 10^9/l$
Creatinine	90 μ mol/l
Na ⁺	136 mmol/l
K ⁺	3.8 mmol/l
Urea	5.2 mmol/l

3

- | | |
|---|---|
| A | Start angiotensin-converting enzyme inhibitor |
| B | Confirm microalbuminuria with a repeat test in 3 months |
| C | Start angiotensin II inhibitor |
| D | Commence low-protein diet |
| E | Start calcium channel antagonist |

Microalbuminuria is a reliable predictive marker of progression to nephropathy in type 1 diabetes mellitus. The early urinary albumin excretion rates of 30–300 mg/24 h are too small to be detected on conventional urine dipsticks, and either special dipsticks or radioimmunoassay is used. The next stages are macroalbuminuria (>300 mg/24 h) and then persistent proteinuria, by which time end-stage renal failure is usually 5–10 years away. The optimal treatment for diabetic albuminuria is an angiotensin-converting enzyme inhibitor (or angiotensin II inhibitor) in combination with meticulous glycaemic and blood pressure control. However, the observation of microalbuminuria should be confirmed twice within a 3–6-month period before commencing treatment.

Rate this question:

Next Question

[Previous Question](#)
[Tag Question](#)

End Session

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	6
Responses Total:	6
Responses - % Correct:	0%

Back to Filters

Question 52 of 203

A 56-year-old woman with chronic renal failure secondary to chronic pyelonephritis was reviewed in the renal clinic. She was complaining of some exertional dyspnoea and fatigue but no other symptoms. She had a history of hypertension that was difficult to control, despite being on multiple agents. She has taken oral iron sulphate. On examination she was pale, blood pressure was 190/110 mmHg, jugular venous pressure not raised. She had bi-basal crepitations in her chest and mild pitting ankle odema. Her abdomen was soft, non-tender and not distended.

Blood results are below:

Hb	8.2 g/dl
WCC	5.0 × 10 ⁹ /l
PLT	220 × 10 ⁹ /l
Mean corpuscular volume (MCV)	74 fl
Ferritin	90 μg/l (20-260 μg/l)
Folate (serum)	10 μg/l (2-11 g/l)
Vitamin B12 (serum)	240 ng/l (150-675 ng/l)

What is the most appropriate management of the anaemia?

- A

Blood transfusion
- B

Intravenous iron infusion
- C

Subcutaneous erythropoietin
- D

Oral ferrous gluconate
- E

Immediate commencement of haemodialysis

9161

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 52 of 203

Blood results are below:

11

What is the most appropriate management of the anaemia?

- | | |
|---|---|
| A | Blood transfusion |
| B | Intravenous iron infusion |
| C | Subcutaneous erythropoietin |
| D | Oral ferrous gluconate |
| E | Immediate commencement of haemodialysis |

- | | |
|---|---------------------------|
| B | Intravenous iron infusion |
|---|---------------------------|

A Blood transfusion

C Subcutaneous erythropoietin

D Oral ferrous gluconate

E	Immediate commencement of haemodialysis
---	---

Rate this question:

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

[Blog](#) [About Pastest](#) [Contact Us](#) [Help](#)

Back to Filters

Question 53 of 203

A 48-year-old woman was admitted to hospital for investigation of a 2-month history of polyuria, polydipsia and increased thirst. She had bipolar affective disorder which was well controlled on lithium and no other medical problems. She smoked 5-10 cigarettes per day and did not drink alcohol. There was no relevant family history. Physical examination was unremarkable.

Investigation results are below:

Na ⁺	148 mmol/l
K ⁺	4.3 mmol/l
Urea	5.2 mmol/l
Creatinine	92 μmol/l
Ca ²⁺	2.30 mmol/l
Glucose (fasting)	5.0 mmol/l
Plasma osmolality	340 mosmol/kg (278-305)
Urine osmolality	280 mosmol/kg (350-1000)

The patient proceeded to a water deprivation test, the results of which are below:

	Urine osmolality (mosmol/kg)
After 8 h fluid deprivation	200
After desmopressin administration	240

What is the diagnosis?

- A

Cranial diabetes insipidus
- B

Psychogenic polydipsia
- C

Nephrogenic diabetes insipidus
- D

Syndrome of inappropriate antidiuretic hormone secretion
- E

Conn’s syndrome

9162

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 53 of 203

A 48-year-old woman was admitted to hospital for investigation of a 2-month history of polyuria, polydipsia and increased thirst. She had bipolar affective disorder which was well controlled on lithium and no other medical problems. She smoked 5-10 cigarettes per day and did not drink alcohol. There was no relevant family history. Physical examination was unremarkable.

Investigation results are below:

Na ⁺	148 mmol/l
K ⁺	4.3 mmol/l
Urea	5.2 mmol/l
Creatinine	92 μmol/l
Ca ²⁺	2.30 mmol/l
Glucose (fasting)	5.0 mmol/l
Plasma osmolality	340 mosmol/kg (278-305)
Urine osmolality	280 mosmol/kg (350-1000)

The patient proceeded to a water deprivation test, the results of which are below:

	Urine osmolality (mosmol/kg)
After 8 h fluid deprivation	200
After desmopressin administration	240

What is the diagnosis?



- A

Cranial diabetes insipidus
- B

Psychogenic polydipsia
- C

Nephrogenic diabetes insipidus
- D

Syndrome of inappropriate antidiuretic hormone secretion
- E

Conn’s syndrome

Explanation

Diabetes insipidus (DI) is due to either reduced vasopressin (ADH) secretion from the posterior pituitary gland (cranial DI) or renal resistance to its action (nephrogenic DI). It must be distinguished from psychogenic polydipsia, which is excessive fluid intake usually secondary to psychiatric disturbance. In DI there should be a high plasma and low urine osmolality. The diagnosis is confirmed by the water deprivation test. The patient is fluid restricted for 8 h and the blood/urine osmolality and weight are measured hourly. In cranial DI, the urine osmolality should rise to more than 660 mosmol/kg after desmopressin, whereas there will be little to no response in nephrogenic DI. The causes of nephrogenic DI include drugs (eg lithium), hypokalaemia and hypercalcaemia, or it can be inherited. Management mainly involves treating or removing the cause, although bendrofluazide can be used. Cranial DI will respond to intranasal desmopressin.

9162

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	8
Responses Total:	8
Responses - % Correct:	0%

Back to Filters

Question 54 of 203

A 48-year-old woman presents with severe left loin pain on a background of several months of malaise, thirst and polyuria. She has a long history of chronic headaches which have been extensively investigated and no pathological cause found. She also suffers from anxiety and depression, and recurrent urinary tract infections. She does not smoke or drink alcohol and has no significant family history. The only significant findings on examination were hypertension and some tenderness in the left loin.

Investigation results are below:

Hb	10.2 g/dl
WCC	10.8 x 10 ⁹ /l
Platelets	300 x 10 ⁹ /l
MCV	86 fl
Ca ²⁺	2.18 mmol/l
Phosphate	1.80 mmol/l
Urate	0.38 mmol/l
Na ⁺	130 mmol/l
K ⁺	3.6 mmol/l
Urea	15.2 mmol/l
Creatinine	380 micromol/l
Plain abdominal film	No visible calculi or calcification
Urinalysis	Protein ++
Urine microscopy	Casts, renal papillary cells and leucocytes. No organisms
Intravenous urogram	Bilateral clubbed calyces, 'ring signs', filling defect in left ureter

What is the most likely diagnosis?

- A

Reflux nephropathy
- B

Renal stone disease
- C

Medullary sponge kidney
- D

Acute interstitial nephritis
- E

Analgesic nephropathy

9164

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 54 of 203

Investigation results are below:

A	Reflux nephropathy
B	Renal stone disease
C	Medullary sponge kidney
D	Acute interstitial nephritis
E	Analgesic nephropathy

The answer is - Analgesic nephropathy

Rate this question:

[Next Question](#)

Tag Question

End Session

Difficulty: Average

Peer Responses %


Session Progress

Responses Correct:	0
Responses Incorrect:	9
Responses Total:	9
Responses - % Correct:	0%

Back to Filters

Question 55 of 203

A 44-year-old woman presented with haematuria and loin pain. Her computed tomography (CT) scan is shown below:



What is the diagnosis?

- A Adult polycystic kidney disease
- B von Hippel-Lindau syndrome
- C Medullary sponge kidney
- D Nephronophthisis
- E Hydatid disease

9168

Submit

Previous Question Skip Question


Calculator

Normal Values

Back to Filters

Question 55 of 203

A 44-year-old woman presented with haematuria and loin pain. Her computed tomography (CT) scan is shown below:



What is the diagnosis?

- A Adult polycystic kidney disease
- B von Hippel-Lindau syndrome
- C Medullary sponge kidney
- D Nephronophthisis
- E Hydatid disease

Explanation

The computed tomography (CT) shows the classic appearance of polycystic kidneys with multiple hepatic cysts (these occur in 30-70% patients). Adult polycystic kidney disease is an autosomal dominant condition and is a common cause of chronic renal failure. It can present at any age from the second decade onwards. Presenting features include loin pain, haematuria, hypertension and abdominal mass. Associated features include intracranial aneurysms (8%) and subarachnoid haemorrhage, mitral valve prolapse and renal calculi. Cysts may become infected or haemorrhage. Children and siblings of patients with this condition should be screened with ultrasound after 20 years of age.

9168

Rate this question: ⚙️ ⭐ ⭐ ⭐ ⭐ ⭐

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	10
Responses Total:	10
Responses - % Correct:	0%

Back to Filters

Question 56 of 203

A 62-year-old man is admitted to hospital with a 3-day history of dyspnoea, pleuritic chest pain and several episodes of haemoptysis. Prior to this he described several months of constant rhinorrhoea with some nasal crusting, several large epistaxes and constant pain below his eyes. More recently he had noticed some double vision and swelling of his right eye. On examination he had a low-grade pyrexia and was normotensive. There was some nasal mucosal ulceration and a right-sided proptosis. Cardiovascular examination was unremarkable. He had localised areas of crepitations throughout both lung fields. His abdomen was soft and non-tender with no masses. There were no focal neurological signs or skin lesions.

Investigation results are below:

Haemoglobin (Hb)	10.0 g/dl
White cell count	12.0 × 10 ⁹ /l
Platelets	390 × 10 ⁹ /l
Mean corpuscular volume (MCV)	90 fl
Erythrocyte sedimentation rate (ESR)	80 mm/h
Na ⁺	142 mmol/l
K ⁺	5.7 mmol/l
Urea	22.0 mmol/l
Creatinine	620 µmol/l
C-reactive protein	238 mg/L
Bilirubin	15 µmol/l
Alkaline phosphatase	88 iu/l
Aspartate transaminase	32 iu/l
Ca ²⁺	2.42 mmol/l
Gamma-glutamyl transferase (GGT)	48 iu/l
Urinalysis	Blood +++ protein ++
Chest radiograph	Multiple large cavitating nodules throughout both lung fields
Electrocardiogram (ECG)	Normal sinus rhythm, no acute changes

What would be the most useful investigation in pointing to the diagnosis if positive?

- A

Serum antiglomerular basement membrane antibodies
- B

Serum antineutrophil cytoplasmic antibodies
- C

High resolution computed tomography (CT) scan of the thorax
- D

Urine microscopy
- E

Serum antinuclear antibodies

9172

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Question 56 of 203

Investigation results are below.



A	Serum antiglomerular basement membrane antibodies
B	Serum antineutrophil cytoplasmic antibodies
C	High resolution computed tomography (CT) scan of the thorax
D	Urine microscopy
E	Serum antinuclear antibodies

This patient has the typical features of granulomatosis with polyangiitis (GPA), (also known as Wegener's), which is a necrotising granulomatous arteritis of the upper and lower respiratory tract and kidney. It is rare, with an incidence of 5-10 per million. It is a multisystem disease and may affect the eyes, skin, joints, heart and nervous system. Pulmonary involvement is seen in 95% of cases and renal involvement in 85% of cases. A positive serum anti-neutrophil cytoplasmic antibody (ANCA) is present in greater than 90% of cases of GPA, and strongly supports the diagnosis. Renal biopsy is sometimes required. Treatment is with steroids and cyclophosphamide. Untreated 1-year mortality is 80%, but with appropriate therapy, remission can be achieved in up to 90% of patients.

Rate this question:

Next Question

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	11
Responses Total:	11
Responses - % Correct:	0%

Question 57 of 203

Initial investigations were as follows:

albumin	29 g/l
urea	5.0 mmol/l
creatinine	60 μ mol/l
sodium	144 mmol/l
potassium	4.0 mmol/l
urinalysis	protein +++
24-h urine collection	3.4 g protein

11

- | | |
|---|--|
| A | Corticosteroids (high dose) |
| B | Cyclophosphamide |
| C | Antibiotics |
| D | Nothing, observation only |
| E | Loop diuretics and an albumin infusion |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 57 of 203

A 16-year-old girl presents with ankle swelling and facial puffiness. Two weeks previously she had a flu-like illness with a productive cough. Her father reports that she had suffered a similar episode of swelling 2 years ago but that things had returned to normal after a period of treatment. At this time U&E had been normal and her father does not recall her having had a rash. On examination there was no rash or fever. She had pitting oedema to the waist.

Initial investigations were as follows:

albumin	29 g/l
urea	5.0 mmol/l
creatinine	60 μ mol/l
sodium	144 mmol/l
potassium	4.0 mmol/l
urinalysis	protein +++
24-h urine collection	3.4 g protein

Given the likely diagnosis, what is the best initial treatment?

- A

Corticosteroids (high dose)
- B

Cyclophosphamide
- C

Antibiotics
- D

Nothing, observation only
- E

Loop diuretics and an albumin infusion

Explanation

The answer is - Corticosteroids (high dose)

Nephrotic syndrome is defined by urinary protein loss of >3.5 g of protein/24 h (0.05 g/kg per 24 h in children) accompanied by hypoalbuminaemia (75% of cases. Upper respiratory tract infection (URTI) commonly precedes attacks although a definite causal relationship is debated-the commonest bacterial antigens implicated are streptococcal. Minimal change disease has a good prognosis in childhood; a third have just one episode, a third have infrequent relapses while a third have frequent relapses throughout childhood that stop in adult life. Steroids are an effective treatment for minimal change disease and as this is by far the commonest cause of childhood nephrotic syndrome, a trial of high dose steroids is accepted practice, with only those patients who do not respond being investigated for other renal disease.

9853

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	12
Responses Total:	12
Responses - % Correct:	0%

Back to Filters

Question 58 of 203

A 48-year-old man was admitted for an endoscopic retrograde cholangiopancreatography (ERCP) for a common bile duct stone. He was given gentamicin and piperacillin to cover the procedure but still developed *Escherichia coli* sepsis, during which time he was systematically unwell with fever, hypotension, (BP 88/50 mmHg) and tachycardia. He was treated with iv cefuroxime. His pre-procedure bloods showed normal renal function. After 4 days he remained jaundiced but was afebrile and haemodynamically stable. There was no palpable bladder on examining his abdomen and his urine output was 40 ml/h. His bloods showed urea 26 mmol/l and creatinine 500 μmol/l with a normal full blood count.

Which of the following treatments for his acute renal failure is the best supported by clinical evidence from current research trials?

A	Low-dose dopamine infusion
B	Low-dose dopamine + high-dose furosemide + mannitol infusion
C	Low-dose dopamine + high-dose furosemide infusion
D	High-dose furosemide
E	None of the above

9854

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Question 58 of 203

Which of the following treatments for his acute renal failure is the best supported by clinical evidence from current research trials?

?

The development of acute renal failure following an episode of hypotension is a common complication of surgery and sepsis. The management of such patients has been controversial. The use of low-dose dopamine and diuretics has been established practice for many years. Evidence-based practise has however lead to a re-evaluation of such therapies based on retrospective analysis and controlled trials. The data that exists in the literature does not support the use of either loop diuretics or dopamine in the management of acute renal failure. There is some evidence to support the use of loop diuretics and mannitol in the specific situation of renal tubular damage complicating rhabdomyolysis and loop diuretics may produce a therapeutic diuresis in oliguric renal failure if the patient requires urgent off-loading and immediate dialysis is not available.

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	13
Responses Total:	13
Responses - % Correct:	0%

Back to Filters

Question 59 of 203

A 20-year-old man of Japanese descent who is studying in the UK attends the clinic for advice. He attended the optician for an eye test and was noted to have greenish-gold discoloration within the limbus of both corneas. Other past medical history of note includes polyarthritis for which he takes regular analgesia. On physical examination his blood pressure is 130/80 mmHg; chest and abdominal examination is unremarkable.

Investigations:

Hb	11.1 g/dl
K ⁺	3.4 mmol/l
Na ⁺	138 mmol/l
Creatinine	130 mol/l
Urea	4.2 mmol/l
Bicarbonate	16 mmol/l
ALT	75 IU/l
Urinary bicarbonate excretion	Increased



Which of the following diagnoses fits best with the renal dysfunction seen here?

- A

Renal tubular acidosis type IV
- B

Renal tubular acidosis type II
- C

Renal tubular acidosis type I
- D

Acute tubular necrosis
- E

Papillary necrosis

18533

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 59 of 203

HD	11.1 g/dl
K ⁺	3.4 mmol/l
Na ⁺	138 mmol/l
Creatinine	130 μmol/l
Urea	4.2 mmol/l
Bicarbonate	16 mmol/l
ALT	75 IU/l
Urinary bicarbonate excretion	Increased

- | | |
|---|--------------------------------|
| A | Renal tubular acidosis type IV |
| B | Renal tubular acidosis type II |
| C | Renal tubular acidosis type I |
| D | Acute tubular necrosis |
| E | Papillary necrosis |

- | | |
|---|--------------------------------|
| B | Renal tubular acidosis type II |
|---|--------------------------------|

A Renal tubular acidosis type IV

Type 4 is associated with a decreased response to aldosterone.

C	Renal tubular acidosis type I
---	-------------------------------

D	Acute tubular necrosis
---	------------------------

E	Papillary necrosis
---	--------------------

Rate this question:

[Next Question](#)

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	14
Responses Total:	14
Responses - % Correct:	0%

Back to Filters

Question 60 of 203

A 42-year-old man who has received long-term haemodialysis for chronic renal failure due to type 1 diabetes presents for review. He has been suffering pain and tingling in both hands, which particularly comes on during the early hours of the morning. He has also noticed shoulder pain over the past few months. He is dialysed using a conventional cellulose acetate dialysis membrane. On physical examination there is no obvious structural deformity in the hands. An X-ray of both hands is ordered and it shows a destructive arthropathy with periarticular cystic bone radiolucencies.

Which of the following diagnoses fits best with this clinical picture?

- A

Uraemic neuropathy
- B

Diabetic nephropathy
- C

AA amyloidosis
- D

AL amyloidosis
- E

Beta 2 microglobulin amyloidosis

18539

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 60 of 203

A 42-year-old man who has received long-term haemodialysis for chronic renal failure due to type 1 diabetes presents for review. He has been suffering pain and tingling in both hands, which particularly comes on during the early hours of the morning. He has also noticed shoulder pain over the past few months. He is dialysed using a conventional cellulose acetate dialysis membrane. On physical examination there is no obvious structural deformity in the hands. An X-ray of both hands is ordered and it shows a destructive arthropathy with periarticular cystic bone radiolucencies.

Which of the following diagnoses fits best with this clinical picture?

- A

Uraemic neuropathy
- B

Diabetic nephropathy
- C

AA amyloidosis
- D

AL amyloidosis
- E

Beta 2 microglobulin amyloidosis

Explanation ⚙

- E

Beta 2 microglobulin amyloidosis

Conventional dialysis membranes cannot clear substances with molecular weights of >200 dalton. These allow accumulation of beta-2 microglobulin, which has a molecular weight of 11,800 dalton. A normal functioning kidney would filter this in the glomerulus and it is catalysed in the proximal tubule. In some patients dialysed for more than 5 years on a conventional membrane, accumulation of beta-2 microglobulin amyloid leads to carpal tunnel syndrome, tenosynovitis, scapulohumeral arthropathy, bony cysts and even pathological fractures. GI or cardiovascular manifestations may occur. The mainstays of management are moving towards polyacrylnitrile or polysulphone dialysis membranes, which are higher flux and allow beta-2 microglobulin to be removed, or moving to transplant if a kidney is available. Other forms of renal amyloidosis occur in association with systemic inflammation, such as in rheumatoid arthritis, or haematological malignancies, such as myeloma. Aggressive use of secondary agents may slow deposition of amyloid in inflammatory conditions, as may chemotherapy in conditions such as myeloma.

- A

Uraemic neuropathy

Similar to diabetic neuropathy, uraemia may lead to neuropathy and hence a neuropathic arthropathy may result in dialysis patients. This is rare and there would be joint deformity with X-ray findings of joint destruction rather than bony cystic changes.

- B

Diabetic nephropathy

Diabetic nephropathy is likely to be the underlying renal disease in this case. However, it would not produce the arthropathy symptoms and signs. Diabetic patients may have diabetic neuropathy that would cause loss of sensation of peripheral nerves, which may then lead to Charcot joints. This would be more common in the feet and also there would be joint deformity, because the joint and bony structures are destroyed.

- C

AA amyloidosis

AA amyloidosis may be secondary to causes such rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis and juvenile arthritis. This is due to the chronic inflammation. These rheumatic conditions may be a cause of joint pain. However, there is no history to suggest rheumatic disease in this scenario. Also, there may be characteristic X-ray signs for these rheumatoid conditions; for instance, X-ray of the hands may show soft tissue swelling and early erosions in the proximal interphalangeal joints. It would not be the bony cystic destruction as seen here.

- D

AL amyloidosis

AL amyloidosis is due to immunoglobulin light chain production. The musculoskeletal manifestations may include muscle weakness and myopathy with arthropathy and osteopathy. There is no history in this case to suggest that this patient has myeloma, such as light chains in the urine or immunoparesis.

18539

Rate this question: ☹ ★ ★ ★ ★ ★

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Difficult

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	15
Responses Total:	15
Responses - % Correct:	0%

Back to Filters

Question 61 of 203

A 74-year-old woman is admitted by ambulance having been found by her daughter on her hall floor. It appears that she had been there for some hours. She has a past history of hypertension, which is treated with ramipril 10 mg daily, and a TIA for which she is treated with aspirin. On examination in the Emergency room her blood pressure is 120/70 mmHg, with a pulse of 105/min; she is pyrexial (38.2°C) and has evidence of right-sided pneumonia. There is a fractured left femoral neck in addition.

Investigations:

Hb	10.9 g.dl
WCC	13.1 x10 ⁹ /l
PLT	235 x10 ⁹ /l
Na ⁺	138 mmol/l
K ⁺	5.8 mmol/l
Creatinine	199 μmol/l
CK	3890 U/l (24-170)
Urine output 25ml over 2	No significant residual on catheterisation
Urine	Positive to blood on dipstick, but no cells seen on microscopy
CXR	Right sided pneumonia
ECG	Lateral ST depression

Which of the following apart from pneumonia would best explain some of the other results seen here?

- A

Anterior myocardial infarction
- B

Urinary tract infection
- C

Rhabdomyolysis
- D

Fat embolus
- E

Subendocardial myocardial infarction

18658

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Back to Filters

Question 61 of 203

A 74-year-old woman is admitted by ambulance having been found by her daughter on her hall floor. It appears that she had been there for some hours. She has a past history of hypertension, which is treated with ramipril 10 mg daily, and a TIA for which she is treated with aspirin. On examination in the Emergency room her blood pressure is 120/70 mmHg, with a pulse of 105/min; she is pyrexial (38.2°C) and has evidence of right-sided pneumonia. There is a fractured left femoral neck in addition.

Investigations:

Hb	10.9 g.dl
WCC	13.1 x10 ⁹ /l
PLT	235 x10 ⁹ /l
Na ⁺	138 mmol/l
K ⁺	5.8 mmol/l
Creatinine	199 μmol/l
CK	3890 U/l (24-170)
Urine output 25ml over 2	No significant residual on catheterisation
Urine	Positive to blood on dipstick, but no cells seen on microscopy
CXR	Right sided pneumonia
ECG	Lateral ST depression

Which of the following apart from pneumonia would best explain some of the other results seen here?

- A

Anterior myocardial infarction
- B

Urinary tract infection
- C

Rhabdomyolysis
- D

Fat embolus
- E

Subendocardial myocardial infarction

Explanation

- C

Rhabdomyolysis

This patient has a raised CK and urine testing, which is suggestive of myoglobinuria (positive dipstick but no cells seen). It is likely she has been lying on the floor for some time, which has precipitated muscle breakdown and the picture consistent with rhabdomyolysis seen here. Her creatinine of 199 μmol/l is likely to represent a degree of acute on chronic renal impairment. Vigorous rehydration with isotonic crystalloid is the mainstay of therapy for rhabdomyolysis. Severe or refractory cases may require haemodialysis.

- A

Anterior myocardial infarction

An anterior myocardial infarction would result from an occlusion of the left anterior descending artery. This carries the worst prognosis of all infarct locations, mainly because it affects a large area. The ECG finding would be ST segment elevation with Q wave formation in the precordial leads (V1-6) and possible the high lateral leads (I and aVL). There would be reciprocal ST depression in the inferior leads (III and aVF). In this scenario there is lateral ST depression, which is not consistent with anterior myocardial infarction.

- B

Urinary tract infection

Although urinary tract infection may result in blood and protein in the urine with fever, this does not produce a raised creatinine kinase which, in this case, is the key to diagnosis.

- D

Fat embolus

A fat embolism is a clot that consists of fatty material. This may result from physical trauma such as a fracture of a long bone, soft tissue trauma or burns. If fat embolism is severe it may cause respiratory failure or myocardial infarction. There may be purpura that appears on the upper anterior body including chest, neck and upper arm - this is pathognomonic sign of fat embolism syndrome. This is not present in this case.

- E

Subendocardial myocardial infarction

Subendocardial myocardial infarction involves infarction of the subendocardial wall of the left ventricle, ventricular septum or papillary muscles. There would be Q wave in myocardial infarction. In this case there is subendocardial ischaemia because of myocardial strain from anaemia and pneumonia. Myocardial strain occurs with inadequate oxygenation to the heart. This is evidenced by the lateral ST depression.

18658

Rate this question: ⚙️☆☆☆☆☆

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	16
Responses Total:	16
Responses - % Correct:	0%

Back to Filters

Question 62 of 203

A 39-year-old man presents to the renal clinic with hypertension and worsening renal function. His GP has tried to manage his blood pressure for many years and he is currently treated with ramipril and amlodipine. He does not know his family medical history as he is adopted. On examination his BP is 155/92 mmHg and you can feel bilateral flank masses; you also feel an irregularly enlarged liver.

Investigations:

Hb	10.9 g/dl
WCC	4.8 x 10 ⁹ /l
PLT	294 x 10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	5.4 mmol/l
Creatinine	290 μmol/l
Abdominal USS - Multiple cysts in both kidneys, cysts also visualised in the liver	
Urine dipstick	blood +

Which of the following is the most important problem associated with his likely underlying diagnosis?

- A

Male infertility
- B

Cerebral aneurysms
- C

Colonic carcinoma
- D

Type 1 diabetes
- E

Mitral stenosis

18659

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 62 of 203

A 39-year-old man presents to the renal clinic with hypertension and worsening renal function. His GP has tried to manage his blood pressure for many years and he is currently treated with ramipril and amlodipine. He does not know his family medical history as he is adopted. On examination his BP is 155/92 mmHg and you can feel bilateral flank masses; you also feel an irregularly enlarged liver.

Investigations:

Hb	10.9 g/dl
WCC	4.8 x 10 ⁹ /l
PLT	294 x 10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	5.4 mmol/l
Creatinine	290 μmol/l
Abdominal USS – Multiple cysts in both kidneys, cysts also visualised in the liver	
Urine dipstick	blood +

Which of the following is the most important problem associated with his likely underlying diagnosis?

- A Male infertility
- B Cerebral aneurysms
- C Colonic carcinoma
- D Type 1 diabetes
- E Mitral stenosis

Explanation ⚙

- B Cerebral aneurysms

This patient has a clinical picture which is consistent with polycystic kidney disease. The condition is autosomal dominant, but we are told that he is adopted, so there is no clue as to his family history. Cerebral aneurysms are associated with greatly increased risk of haemorrhagic stroke and, as such, should be managed by a specialist neurosurgeon. Other associations include pancreatic and hepatic cysts, colonic diverticula and mitral valve prolapse.

- A Male infertility

Patients with polycystic kidney disease have normal fertility. Because the condition is autosomal dominant it is important to give patients genetic counselling as to the likelihood of any offspring having polycystic kidney disease and the need to have screening if they are adults.

- C Colonic carcinoma

Colonic diverticulum is associated with polycystic kidney disease rather than colonic carcinoma. Malignancy may be associated with other renal disease such as membranous nephropathy.

- D Type 1 diabetes

Diabetes is not associated with polycystic kidney disease. The most common renal abnormality with type 1 diabetes is diabetic nephropathy.

- E Mitral stenosis

Mitral valve prolapse rather than mitral stenosis is associated with polycystic kidney disease. The valve no longer closes properly and blood leaks backwards in mitral valve prolapse.

18659

Rate this question: ⚙ ⭐⭐⭐⭐⭐

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	17
Responses Total:	17
Responses - % Correct:	0%

Question 63 of 203

Investigations;

Hb	11.5 g/dl
WCC	5.6 x10 ⁹ /l
PLT	295 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	5.0 mmol/l
Creatinine	656 μ mol/l
Glucose	3.9 mmol/l
CT head	no focal lesion identified

3

- | | |
|---|---------------------------------|
| A | Bacterial meningitis |
| B | Viral meningitis |
| C | Uraemic encephalopathy |
| D | Erythropoietin induced epilepsy |
| E | Ischaemic stroke |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 63 of 203

A 49-year-old man with a history of Type 1 diabetes is admitted to the Emergency room with an epileptic seizure. He has chronic renal failure and has been receiving dialysis for the past 3 months. Medication includes ramipril, amlodipine, aspirin, calcium and vitamin D, and periodic injections of recombinant erythropoietin which were begun recently. On examination his BP is 145/85 mmHg, with a temp of 37.8°C, his GCS is 13.

Investigations;

Hb	11.5 g/dl
WCC	5.6 x10 ⁹ /l
PLT	295 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	5.0 mmol/l
Creatinine	656 μmol/l
Glucose	3.9 mmol/l
CT head	no focal lesion identified

Which of the following is the most likely cause of his seizure?

- A

Bacterial meningitis
- B

Viral meningitis
- C

Uraemic encephalopathy
- D

Erythropoietin induced epilepsy
- E

Ischaemic stroke

Explanation

This is erythropoietin induced epilepsy. This rare adverse event is seen within 90 days of erythropoietin initiation. His white blood cell count is normal, making sepsis less likely and the elevated temperature here is just likely to be the result of his fit. The creatinine of 656 μmol/l is most likely to be due to the interval between his dialysis sessions. The exact mechanism related to the seizures is unclear, but may be related to veno-occlusive disease or the rise in haemoglobin associated with EPO use.

18668

Rate this question:

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	18
Responses Total:	18
Responses - % Correct:	0%

Back to Filters

Question 64 of 203

A 72-year-old man with a history of hypertension presents to the Emergency department in extremis. He awoke in the early hours of the morning with severe shortness of breath as if he was going to gasp his last breath. You understand that his GP has commenced him on ramipril 5mg daily a short time ago. On examination he has a BP of 175/85 mmHg, a pulse of 100/min and is in left ventricular failure. You manage him with a nitrate infusion and furosemide and he improves.

Investigations:

Hb	13.1 g/dl
WCC	5.0 x10 ⁹ /l
PLT	212 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.5 mmol/l
Creatinine	204 μmol/l (134 one month earlier)

Echo shows mild LVH

What is the most likely cause of his presentation with pulmonary oedema?

- A

Coronary artery disease
- B

Hypertensive heart disease
- C

Hypertrophic cardiomyopathy
- D

Aortic stenosis
- E

Renal artery stenosis

20553

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 64 of 203

A 72-year-old man with a history of hypertension presents to the Emergency department in extremis. He awoke in the early hours of the morning with severe shortness of breath as if he was going to gasp his last breath. You understand that his GP has commenced him on ramipril 5mg daily a short time ago. On examination he has a BP of 175/85 mmHg, a pulse of 100/min and is in left ventricular failure. You manage him with a nitrate infusion and furosemide and he improves.

Investigations:

Hb	13.1 g/dl
WCC	5.0 x10 ⁹ /l
PLT	212 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.5 mmol/l
Creatinine	204 µmol/l (134 one month earlier)

Echo shows mild LVH

What is the most likely cause of his presentation with pulmonary oedema?

- A

Coronary artery disease
- B

Hypertensive heart disease
- C

Hypertrophic cardiomyopathy
- D

Aortic stenosis
- E

Renal artery stenosis

Explanation

- E

Renal artery stenosis

Patients with renal artery stenosis may present in this manner with rapidly worsening renal function, an increase in blood pressure and so-called flash pulmonary oedema. In this case, the presentation has almost certainly been accelerated by the starting of an ACE inhibitor. Investigations to confirm the diagnosis include ultrasound, which may show differential kidney size, and magnetic resonance angiography. Management centres on non-ACE inhibitor control of blood pressure. Large-scale studies have suggested little or no benefit of angioplasty in renal artery stenosis, although it may be appropriate in patients with recurrent episodes of pulmonary oedema.

- A

Coronary artery disease

Myocardial infarction may be a cause of pulmonary oedema and renal impairment. However, the start of ACEI recently would not result in coronary artery disease. The echo would show left ventricular failure rather than just mild LVH.

- B

Hypertensive heart disease

Hypertensive heart disease can cause left ventricular failure for patients who have hypertension over a long period of time. There would be severe left ventricular hypertrophy and left ventricular failure on the echo.

- C

Hypertrophic cardiomyopathy

Hypertrophic cardiomyopathy is the leading cause of death in young athletes. This condition can produce symptoms that mimic congestive heart failure, such as activity intolerance and dyspnoea. On the echo there is systolic anterior motion of the anterior leaflet of the mitral valve. In this scenario, the echo findings do not show this.

- D

Aortic stenosis

Aortic stenosis may be a cause of flash pulmonary oedema. However, you would expect severe left ventricular hypertrophy if there is sufficiently severe aortic stenosis to result in renal impairment and pulmonary oedema. Furthermore ACEI should be helpful in aortic stenosis for cardiac remodelling and blood pressure control. In this case, the ACEI resulting in acceleration of renal impairment suggests renal artery stenosis.

20553

Rate this question:

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	19
Responses Total:	19
Responses - % Correct:	0%

Question 65 of 203

Investigations;

Hb	10.4 g/dl
WCC	$6.1 \times 10^9/\text{l}$
PLT	$180 \times 10^9/\text{l}$
Na ⁺	139 mmol/l
K ⁺	4.9 mmol/l
Creatinine	276 $\mu\text{mol/l}$
Urine	Blood +, Protein +

17

- | | |
|---|--|
| A | IV hydrocortisone |
| B | Co-amoxiclav |
| C | IV methylprednisolone and cyclophosphamide |
| D | IV normal saline |
| E | IV sodium bicarbonate |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 65 of 203

A 32-year-old man is being treated with an indinavir-containing HAART regime for HIV infection. Other medication of note includes co-trimoxazole which was recently started as prophylaxis against PCP. He has noticed increasing nausea and tiredness over the past few weeks.

Investigations;

Hb	10.4 g/dl
WCC	6.1 x10 ⁹ /l
PLT	180 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.9 mmol/l
Creatinine	276 µmol/l
Urine	Blood +, Protein +

Given the most likely renal diagnosis, which of the following is the most appropriate therapy for this patient?

- A

IV hydrocortisone
- B

Co-amoxiclav
- C

IV methylprednisolone and cyclophosphamide
- D

IV normal saline
- E

IV sodium bicarbonate

Explanation ⚙

- D

IV normal saline

This patient has crystal nephropathy, a condition known to be associated with indinavir use, which is a common component of HAART. Certain drug-drug interactions lead to increased indinavir concentration, one of which is co-trimoxazole. Adequate rehydration is crucial to avoid crystal nephropathy, and patients taking indinavir are recommended to take at least 2-3 litres of fluid per day. IV rehydration is the treatment of choice for crystal nephropathy and, given adequate fluid replacement, indinavir therapy can often continue.

- A

IV hydrocortisone

IV hydrocortisone is not a treatment for crystal nephropathy. Hydrocortisone may be used for treatment of adrenal failure as a replacement for corticosteroids.

- B

Co-amoxiclav

Co-amoxiclav would treat any bacterial urinary tract infection. Although blood and protein in the urine may be mistaken for a urinary tract infection (UTI), you would expect other symptoms such as dysuria and urgency of urination. In addition, the white cell count may be raised in a UTI.

- C

IV methylprednisolone and cyclophosphamide

IV methylprednisolone and cyclophosphamide is not used for treatment of renal stones, but it may be used in this combination to treat renal vasculitis. In this condition there is a rapidly progressive renal dysfunction with blood and protein in the urine.

- E

IV sodium bicarbonate

Sodium bicarbonate therapy may be used in some renal stone disease but not for indinavir crystals. An example is that in one short-term study it has been reported that sodium bicarbonate is an effective treatment for hypocitraturic calcium oxalate stones.¹

20636

Rate this question: ⊖ ★ ★ ★ ★ ★

Next Question

- Previous Question

Tag Question
- Feedback

End Session

Difficulty: Average

Peer Responses %

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

0%

Back to Filters

Question 66 of 203

A 42-year-old man has completed 2 years of haemodialysis and is suffering increasing symptoms of tiredness and lethargy. His past medical history includes hypertension and type II diabetes mellitus. He is frustrated with waiting for a cadaveric transplant and wants to discuss options for securing a new kidney. On physical examination his blood pressure is 140/80 mmHg; chest and abdominal examination is unremarkable.

Investigations:

Hb	10.9 g/dl
WCC	5.4 x10 ⁹ /l
PLT	167 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	5.4 mmol/l
Creatinine	456 μ mol/l

Ejection fraction 48%

Which of the following is the best advice?

- A

Convince him to take his willing 13-year-old son’s kidney
- B

Suggest that he will have to wait his turn on cadaveric transplant list
- C

His wife is highly likely to be a match and should be approached
- D

Refer for urgent cadaveric transplant
- E

Urge him to contact an overseas supplier to source a kidney

20639

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 66 of 203

Investigations:

Hb	10.9 g/dl
WCC	5.4 x10 ⁹ /l
PLT	167 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	5.4 mmol/l
Creatinine	456 μmol/l

Ejection fraction 48%

Which of the following is the best advice?

- | | |
|---|---|
| A | Convince him to take his willing 13-year-old son's kidney |
| B | Suggest that he will have to wait his turn on cadaveric transplant list |
| C | His wife is highly likely to be a match and should be approached |
| D | Refer for urgent cadaveric transplant |
| E | Urge him to contact an overseas supplier to source a kidney |

- | | |
|---|---|
| B | Suggest that he will have to wait his turn on cadaveric transplant list |
|---|---|

A Convince him to take his willing 13-year-old son's kidney

C His wife is highly likely to be a match and should be approached

D Refer for urgent cadaveric transplant

E	Urge him to contact an overseas supplier to source a kidney
---	---

Rate this question:

End Session

Difficulty: Average

Peer Responses %

Section D

Responses Correct:	0
Responses Incorrect:	21
Responses Total:	21
Responses - % Correct:	0%

Question 67 of 203

Investigations;

Hb	11.8 g/dl
WCC	5.4 x10 ⁹ /l
PLT	145 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.5 mmol/l
Creatinine	132 mol/l
Urine	blood ++, protein +

■
 ■
 ■

- | | |
|---|------------------|
| A | Urine culture |
| B | Serum IgA levels |
| C | ASOT |
| D | Renal biopsy |
| E | Cystoscopy |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Question 67 of 203

investigations,

Hb	11.8 g/dl
WCC	5.4 x10 ⁹ /l
PLT	145 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.5 mmol/l
Creatinine	132 mol/l
Urine	blood ++, protein +

Which of the following investigations is most likely to confirm the diagnosis?

- | | |
|---|------------------|
| A | Urine culture |
| B | Serum IgA levels |
| C | ASOT |
| D | Renal biopsy |
| E | Cystoscopy |

- | | |
|---|--------------|
| D | Renal biopsy |
|---|--------------|

A	Urine culture
---	---------------

B Serum IgA levels

C	ASOT
---	------

E	Cystoscopy
---	------------

Rate this question:

End Session

Peer Responses %

Session Progress	
Responses Correct:	0
Responses Incorrect:	22
Responses Total:	22
Responses - % Correct:	0%

Back to Filters

Question 68 of 203

A 19-year-old man presents to the clinic with raised BP of 152/82 mmHg. He has been previously fit and well as a child, with no developmental problems. On examination, apart from the raised blood pressure, his chest and abdominal examination is unremarkable. You are concerned that he may have serious underlying pathology and arrange a series of tests.

Investigations:

Hb	12.1 g/dl
WCC	5.0 x10 ⁹ /l
PLT	204 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	3.0 mmol/l
Creatinine	105 μmol/l
Bicarbonate	35 mmol/l
Aldosterone erect	150 pmol/l (200-1000)
Renin erect	2.2 pmol/ml/hr (2.8-4.5)

Which of the following is the most likely diagnosis?

- A

Renal artery stenosis
- B

Bartter syndrome
- C

Gitelman syndrome
- D

Conn syndrome
- E

Liddle syndrome

20985

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 68 of 203

Investigations:

Hb	12.1 g/dl
WCC	5.0 x10 ⁹ /l
PLT	204 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	3.0 mmol/l
Creatinine	105 μ mol/l
Bicarbonate	35 mmol/l
Aldosterone erect	150 pmol/l (200-1000)
Renin erect	2.2 pmol/ml/hr (2.8-4.5)

- | | |
|---|-----------------------|
| A | Renal artery stenosis |
| B | Bartter syndrome |
| C | Gitelman syndrome |
| D | Conn syndrome |
| E | Liddle syndrome |

- | | |
|---|-----------------|
| E | Liddle syndrome |
|---|-----------------|

A Renal artery stenosis

- | | |
|---|-----------------------|
| A | Renal artery stenosis |
|---|-----------------------|
- Renal artery stenosis would be associated with elevated renin and

B	Bartter syndrome
---	------------------

- B Bartter syndrome

C	Gitelman syndrome
---	-------------------

- C Gitelman syndrome

D Conn syndrome

- | | |
|---|---------------|
| D | Conn syndrome |
|---|---------------|

Rate this question:      

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	23
Responses Total:	23
Responses - % Correct:	0%

Back to Filters

Question 69 of 203

A 39-year-old man with a history of treated hypertension treated with ramipril 10mg daily comes to see the GP complaining of tiredness and lethargy. He runs some routine bloods which are given below:

Investigations:

Hb	12.0 g/dl
WCC	5.0 x10 ⁹ /l
PLT	170 x10 ⁹ /l
Na ⁺	133 mmol/l
K ⁺	5.6 mmol/l
Bicarbonate	16 mmol/l
Creatinine	180 μmol/l

Which of the following is the most likely diagnosis?

- A

Renal tubular acidosis type I
- B

Renal tubular acidosis type II
- C

Renal tubular acidosis type IV
- D

Addison’s disease
- E

Conn syndrome

21009

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 69 of 203

A 39-year-old man with a history of treated hypertension treated with ramipril 10mg daily comes to see the GP complaining of tiredness and lethargy. He runs some routine bloods which are given below:

Investigations:

Hb	12.0 g/dl
WCC	5.0 x10 ⁹ /l
PLT	170 x10 ⁹ /l
Na ⁺	133 mmol/l
K ⁺	5.6 mmol/l
Bicarbonate	16 mmol/l
Creatinine	180 μmol/l

Which of the following is the most likely diagnosis?



- A

Renal tubular acidosis type I
- B

Renal tubular acidosis type II
- C

Renal tubular acidosis type IV
- D

Addison’s disease
- E

Conn syndrome

Explanation



- C

Renal tubular acidosis type IV

Type IV RTA, also known as hyperkalaemic RTA occurs when the kidney fails to respond to aldosterone. This leads to hyperkalaemic metabolic acidosis. Pharmacological causes of type IV RTA include spironolactone, ACE inhibitors and ARBS, trimethoprim, heparin, pentamidine and NSAIDs. Structural kidney diseases such as SLE and amyloidosis can also affect response to aldosterone and lead to type IV RTA. Where there is a causative agent that can be discontinued, alternative therapy should be provided, otherwise patients may be offered bicarbonate supplements and agents to lower serum potassium.

- A

Renal tubular acidosis type I

There is disordered excretion of acid (H⁺) in this condition from the collecting ducts, resulting in acidosis. Therefore the clinical picture will be a hyperchloraemic, hypokalaemic metabolic acidosis with hypophosphataemic metabolic bone disease, renal stones or diffuse nephrocalcinosis. It is rarely inherited, or it may be secondary to conditions such as chronic tubular-interstitial disease and dysproteinaemias.

- B

Renal tubular acidosis type II

There is impaired bicarbonate retention in the proximal tubule leading to bicarbonate wasting and a systemic acidosis. This will present with a hyperchloraemic metabolic acidosis, with other features of proximal tubular dysfunction such as phosphate wasting and reduced serum urate. There would be glycosuria and proteinuria with loss of amino acids.

- D

Addison’s disease

Patients with Addison’s disease have low blood pressure rather than hypertension. There may be hyperpigmentation of the skin, even in areas that are not sun-exposed. Characteristic areas of darkening include skin creases, nipple and buccal mucosa. This is because melanocyte-stimulating hormone and ACTH share the same precursor (pro-opiomelanocortin). Investigations review a low sodium, high potassium, high calcium and low glucose.

- E

Conn syndrome

Conn syndrome is associated with elevated aldosterone and therefore the serum potassium would be low rather than high in this case. Patients with Conn syndrome often have few or no symptoms, but some may get occasional muscle weakness, muscle spasms, paraesthesiae or polyuria. High blood pressure and muscle cramps occur due to low serum calcium.

21009

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	24
Responses Total:	24
Responses - % Correct:	0%

Back to Filters

Question 70 of 203

A 62-year-old man with a long history of type 2 diabetes and chronic renal failure comes to the clinic complaining of tiredness. Current therapy includes Mixtard insulin, ramipril, amlodipine, atorvastatin and aspirin. On examination, his blood pressure is 13,080 mmHg, he looks pale and his chest and abdominal examination is unremarkable.

Investigations:

Hb	9.7 g/dl
WCC	4.9 x10 ⁹ /l
PLT	201 x10 ⁹ /l
Na ⁺	141 mmol/l
K ⁺	5.1 mmol/l
Creatinine	210 μmol/l
Ferritin	260 μg/l

Which of the following is the most appropriate management for him?

- A

Commence oral ferrous sulphate
- B

Treat with EPO to a target Hb of 13.5g/dl
- C

Arrange IV iron replacement
- D

Treat with EPO to a target Hb of 11 g/dl
- E

No current intervention, but plan for review in 3 months

21014

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 70 of 203

A 62-year-old man with a long history of type 2 diabetes and chronic renal failure comes to the clinic complaining of tiredness. Current therapy includes Mixtard insulin, ramipril, amlodipine, atorvastatin and aspirin. On examination, his blood pressure is 13,080 mmHg, he looks pale and his chest and abdominal examination is unremarkable.

Investigations:

Hb	9.7 g/dl
WCC	4.9 x10 ⁹ /l
PLT	201 x10 ⁹ /l
Na ⁺	141 mmol/l
K ⁺	5.1 mmol/l
Creatinine	210 μmol/l
Ferritin	260 μg/l

Which of the following is the most appropriate management for him?

- A

Commence oral ferrous sulphate
- B

Treat with EPO to a target Hb of 13.5g/dl
- C

Arrange IV iron replacement
- D

Treat with EPO to a target Hb of 11 g/dl
- E

No current intervention, but plan for review in 3 months

Explanation

- D

Treat with EPO to a target Hb of 11 g/dl

Further ferritin replacement will not improve his Hb levels further. A Cochrane meta-analysis suggested that treating to the higher Hb target of 13.5 g/dl may result in increased risk of hypertensive crises. European renal guidelines therefore recommend a lower target haemoglobin of 11 g/dl.

- A

Commence oral ferrous sulphate

This patient is iron replete. The recommendation from the UK Renal Association is for the ferritin to be >100mg/L and TSAT >20% (or <10% hypochromic red cells). Therefore, there is no need for further iron replacement.

- B

Treat with EPO to a target Hb of 13.5g/dl

As explained above, a Cochrane meta-analysis suggested that treating to the higher Hb target of 13.5 g/dl may result in increased risk of hypertensive crises. European renal guidelines therefore recommend a lower target haemoglobin of 11 g/dl.

- C

Arrange IV iron replacement

The recommendation from the UK Renal Association is for the ferritin to be >100 mg/l and TSAT >20% (or <10% hypochromic red cells). Therefore, this patient is iron replete and there is no need for additional iron replacement.

- E

No current intervention, but plan for review in 3 months

Given his symptoms of lethargy, it would be unacceptable to leave his Hb at 9.7 g/dl. It is important to ensure that he is iron replete, which he is, then to increase haemoglobin by administering erythropoietin.

21014

Rate this question: ⓪☆☆☆☆☆

Next Question

- Previous Question

Feedback
- Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	25
Responses Total:	25
Responses - % Correct:	0%

Back to Filters

Question 71 of 203

A 22-year-old man with a history of nephrotic syndrome due to minimal change disease comes to the clinic because he feels more poorly than usual. He has noticed that despite his usual dose of furosemide, his leg oedema is worsening and for the past few days he has suffered from some vague left flank pain. On examination his BP is 155/92 mmHg and you confirm the left flank pain.

Investigations;

Hb	11.1 g/dl
WCC	7.1 x10 ⁹ /l
PLT	232 x10 ⁹ /l
Na ⁺	140 mmol/l
K ⁺	5.4 mmol/l
Creatinine	280 μmol/l (190 μmol/l at clinic a month earlier)
24h urinary protein	3g (up from 1.6g last time it was assessed)
Cholesterol	6.9 mmol/l

Which of the following is the most likely diagnosis?

- A

Renal artery thrombosis
- B

Renal vein thrombosis
- C

Acute pyelonephritis
- D

Nephrolithiasis
- E

Left sided renal cysts

21164

Submit

Previous Question

Skip Question

Calculator



Normal Values



Back to Filters

Question 71 of 203

A 22-year-old man with a history of nephrotic syndrome due to minimal change disease comes to the clinic because he feels more poorly than usual. He has noticed that despite his usual dose of furosemide, his leg oedema is worsening and for the past few days he has suffered from some vague left flank pain. On examination his BP is 155/92 mmHg and you confirm the left flank pain.

Investigations;

Hb	11.1 g/dl
WCC	7.1 x10 ⁹ /l
PLT	232 x10 ⁹ /l
Na ⁺	140 mmol/l
K ⁺	5.4 mmol/l
Creatinine	280 µmol/l (190 µmol/l at clinic a month earlier)
24h urinary protein	3g (up from 1.6g last time it was assessed)
Cholesterol	6.9 mmol/l

Which of the following is the most likely diagnosis?



- A

Renal artery thrombosis
- B

Renal vein thrombosis
- C

Acute pyelonephritis
- D

Nephrolithiasis
- E

Left sided renal cysts

Explanation



- B

Renal vein thrombosis

Patients with nephrotic syndrome are known to suffer from a hypercoagulable state and, as such, are more prone to suffering from renal vein thrombosis. This picture of vague flank pain, coupled with worsening creatinine and proteinuria, is fairly typical. Initial investigation with ultrasound is likely to reveal a swollen, oedematous kidney on the left when compared to the other side. Long-term anticoagulation may be considered in this patient for prophylaxis against further thromboembolic events.

- A

Renal artery thrombosis

Renal artery thrombosis causes complete blockage to the blood flow through the main arteries. This is a rare cause of acute kidney injury. The clinical manifestation depends on the amount of renal tissue being supplied by the renal artery being blocked by the embolism. It may range from mild increase in serum creatinine to complete anuria. Most patients experience a sudden onset of abdominal, flank or back pain. Renal artery thrombosis is not associated with minimal change disease/nephritic syndrome. The major causes of renal artery stenosis include thromboemboli (systemic clot emboli) and atheroemboli (cholesterol emboli).

- C

Acute pyelonephritis

Pyelonephritis may give left flank pain, but there would be other symptoms such as fever and chills or rigors. You would expect the white cell count to be raised. There is no other history to suggest acute pyelonephritis.

- D

Nephrolithiasis

Nephrolithiasis typically causes renal colic when passed. Patients are usually male and in their 4th decade. There would be renal angle tenderness and haematuria, which is often macroscopic with urinary symptoms such as dysuria and frequency of urination. Predisposing factors include dehydration, exercise and high protein load. There is no history to support renal stones in this case.

- E

Left sided renal cysts

Renal cysts may give rise to left flank pain, especially with haemorrhage or rupture of the cyst. There is no increase in incidence of renal cysts in minimal change disease and there is no history in this case to suggest that renal cysts are present.

21164

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	26
Responses Total:	26
Responses - % Correct:	0%

Question 72 of 203

Investigations:

Hb	11.4 g/dl
WCC	5.9 x10 ⁹ /l
PLT	201 x10 ⁹ /l
Na ⁺	138 mmol/l
K ⁺	4.9 mmol/l
Creatinine	175 μmol/l
Ca ⁺⁺	2.1 mmol/l
IgA	0.7 g/l (0.8-3.0)
IgG	8.9 g/l (6.0-13.0)
IgM	1.0 g/l (0.4-2.5)
24hr urinary protein	3.5g

3

- | | |
|---|------------------------------------|
| A | Myeloma kidney |
| B | Membranous nephropathy |
| C | AL amyloidosis |
| D | Minimal change disease |
| E | Focal segmental glomerulosclerosis |

Submit

Skip Question

Back to Filters

Question 72 of 203

A 62-year-old man is referred to the renal clinic with swollen ankles. His GP is managing him for hypertension and he is currently managed with ramipril, indapamide and amlodipine. The GP is concerned as he noticed 3+ of proteinuria on dipstick testing. On examination in the clinic is BP is 149/91 mmHg, with pulse is 88/min and regular. He has bilateral pitting oedema.

Investigations:

Hb	11.4 g/dl
WCC	5.9 x10 ⁹ /l
PLT	201 x10 ⁹ /l
Na ⁺	138 mmol/l
K ⁺	4.9 mmol/l
Creatinine	175 mol/l
Ca ⁺⁺	2.1 mmol/l
IgA	0.7 g/l (0.8-3.0)
IgG	8.9 g/l (6.0-13.0)
IgM	1.0 g/l (0.4-2.5)
24hr urinary protein	3.5g

Which of the following is the most likely primary diagnosis?

- A

Myeloma kidney
- B

Membranous nephropathy
- C

AL amyloidosis
- D

Minimal change disease
- E

Focal segmental glomerulosclerosis

Explanation



- B

Membranous nephropathy

In this particular age group, membranous nephropathy is the most likely diagnosis (as opposed to minimal change disease, which presents in younger individuals), with the insidious onset as seen here typical of the presentation. Whilst membranous nephropathy may be associated with malignancy, in fact only 5-10% of cases are associated with an underlying cancer. Autoimmune pathology, infectious diseases and drugs may also be associated with the development of the condition.

- A

Myeloma kidney

In the presence of near-normal immunoglobulins, (IgA is only very slightly reduced), myeloma kidney is not likely to be the underlying diagnosis.

- C

AL amyloidosis

Given the lack of associated chronic disease, amyloidosis is unlikely. AL amyloidosis is caused by monoclonal light chains (usually lambda) or light-chain fragments produced by a plasma cell dyscrasia that forms amyloid sheets. It is more common in men than women, by a ratio of 2:1, and most frequently in patients >50 years of age.

- D

Minimal change disease

Minimal change disease is a more common cause of nephrotic syndrome in young children. The renal biopsy is normal on light microscopy and immunofluorescence is negative. Electron microscopy reveals diffuse effacement of podocyte foot processes.

- E

Focal segmental glomerulosclerosis

Focal segmental glomerulosclerosis (FSGS) could be a differential for proteinuria, but it is less common than membranous disease. The cause is unknown and presentation is proteinuria with microscopic haematuria and hypertension. Impaired renal function is common. It is more common in young black males. There is often profound hypoalbuminaemia (albumin <20 g/l) and renal impairment running a severe course, leading to end-stage renal failure.

21222

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	27
Responses Total:	27
Responses - % Correct:	0%

Back to Filters

Question 73 of 203

A 45-year-old man with a history of IgA nephropathy comes to the renal clinic. He has a history of microscopic haematuria and proteinuria which has been under follow-up for 10 years, and a recent renal biopsy has suggested IgA nephropathy with extensive sclerosis. On examination he has a BP of 158/92 mmHg and has bilateral ankle oedema.

Investigations:

Hb	11.3 g/dl
WCC	6.1 x10 ⁹ /l
PLT	201 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.8 mmol/l
Creatinine	280 μmol/l
PTH	10 pmol/l (0.9-5.4)
24h urinary protein	4.7g

Which of the following is the most important additional therapy for him?

- A

Vitamin D
- B

Sevelamer
- C

Ramipril
- D

Atenolol
- E

Prednisolone and azathioprine

21227

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 73 of 203

investigations.

Which of the following is the most important additional therapy for him?

- Explanation ⚙

This man has a well-established glomerulosclerosis related to his IgA nephropathy with impaired renal function; as such, aggressive immunosuppressive therapy is less likely than improved blood pressure control to have a significant impact on his prognosis. Among available blood pressure agents, ACE inhibitors are thought to have the greatest effect in reducing progression of proteinuria and worsening of creatinine.

- vitamin D would not be a treatment in IgA nephropathy. Many patients with renal disease are vitamin D deficient, and therefore it would be useful for measurement and replacement with vitamin D if levels are low. This is especially if patients have symptoms of lethargy.

- Sevelamer is a phosphate binder, and we aren't given information on calcium/phosphate status here, but the PTH is within twice the upper limit of normal. Guidelines only recommend treating PTH when it is above twice the upper limit of normal because of the risk of adynamic bone disease.

- Atenolol would be useful to control blood pressure. However, the evidence is for use of ACEI for the reduction of proteinuria and worsening of creatinine in IgA nephropathy.

- Because glomerulosclerosis is established, aggressive immunosuppressive therapy is less likely than improved blood pressure control to have a significant impact on his prognosis. Prednisolone and azathioprine may be considered if there are crescents in the biopsy or if there is active disease with rapidly

21227

End Session

Peer Responses %

0.001

[Blog](#) [About Pastest](#) [Contact Us](#) [Help](#)

Back to Filters

Question 74 of 203

A patient with chronic renal failure related to diabetes mellitus comes to the clinic. She is managed with a number of anti-hypertensive medications and BD insulin to control her blood sugar. She also takes atorvastatin and aspirin. On examination her BP is 142/82 mmHg, pulse is 85/min and regular. Her chest and abdominal examination is unremarkable.

Investigations:

Hb	10.9 g/dl
WCC	5.9 x 10 ⁹ /l
PLT	201 x 10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	5.0 mmol/l
Creatinine	230 μmol/l
Ca ⁺⁺	2.1 mmol/l
PO ₄ ³⁻	0.85 mmol/l
PTH	16.1 pmol/l (0.9-5.4)

Which of the following is the most appropriate treatment?

- A

Ergocalciferol
- B

Alphacalcidol
- C

Calcitonin
- D

Sevelamer
- E

Cinacalcet

21228

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Question 74 of 203

Investigations:

Hb	10.9 g/dl
WCC	$5.9 \times 10^9/\text{l}$
PLT	$201 \times 10^9/\text{l}$
Na ⁺	139 mmol/l
K ⁺	5.0 mmol/l
Creatinine	230 μmol/l
Ca ⁺⁺	2.1 mmol/l
PO ₄ ³⁻	0.85 mmol/l
PTH	16.1 pmol/l (0.9-5.4)

- | | |
|---|----------------|
| A | Ergocalciferol |
| B | Alphacalcidol |
| C | Calcitonin |
| D | Sevelamer |
| E | Cinacalcet |

- | | |
|---|---------------|
| B | Alphacalcidol |
|---|---------------|

A	Ergocalciferol
---	----------------

C	Calcitonin
---	------------

D Sevelamer

E	Cinacalcet
---	------------

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	29
Responses Total:	29
Responses - % Correct:	0%

Back to Filters

Question 75 of 203

A 54-year-old man with chronic renal failure managed with peritoneal dialysis. He has been on dialysis for the past 3 years and has a history of long-standing Type 1 diabetes. On examination in the clinic his BP is 152/84 mmHg, pulse 70/min and regular. There are no unexpected findings.

Investigations;

Hb	10.4 g/dl
WCC	5.4 x10 ⁹ /l
PLT	180 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.5 mmol/l
Creatinine	480 mol/l
Ca ⁺⁺	2.85 mmol/l
PO ₄ ³⁻	1.8 mmol/l
PTH	18.2 pmol/l (0.9-5.4)

Which of the following best treats the range of abnormalities associated with tertiary hyperparathyroidism?

- A

Calcium and vitamin D
- B

Vitamin D
- C

Sevelamer
- D

Cinacalcet
- E

Low phosphate diet

21229

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Question 75 of 203

Investigations;

Hb	10.4 g/dl
WCC	5.4 x10 ⁹ /l
PLT	180 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.5 mmol/l
Creatinine	480 μmol/l
Ca ⁺⁺	2.85 mmol/l
PO ₄ ³⁻	1.8 mmol/l
PTH	18.2 pmol/l (0.9-5.4)

- | | |
|---|-----------------------|
| A | Calcium and vitamin D |
| B | Vitamin D |
| C | Sevelamer |
| D | Cinacalcet |
| E | Low phosphate diet |

Explanation

Vitamin D (1-alpha-calcidol) is used in the early stages of hyperparathyroidism related to renal failure, when calcium levels are low-normal, and PTH is above twice the upper limit of the normal range. In this case we have both a raised serum phosphate and marked hyperparathyroidism. In this respect the PTH antagonist cinacalcet may be used in the treatment of hyperparathyroidism, in conjunction with other measures such as a low phosphate diet and a suitable phosphate binder such as Sevelamer, but only if the patient is unable to undergo surgery.

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	30
Responses Total:	30
Responses - % Correct:	0%

Back to Filters

Question 76 of 203

A 32-year-old man attends the clinic as an urgent referral from his GP, he has been suffering from a cough, fevers and increasing lethargy over the past few days, but most recently has begun to suffer from haemoptysis. When he makes it to the clinic he looks quite unwell, with a raised respiratory rate of 31/min. His BP is 155/82 mmHg and he has crackles at both lung bases on auscultation.

Investigations:

Hb	10.9 g/dl
WCC	12.1 × 10 ⁹ /l
PLT	185 × 10 ⁹ /l
Na ⁺	141 mmol/l
K ⁺	5.9 mmol/l
Creatinine	194 μmol/l
c-ANCA	Negative
ESR	74 mm/h
CXR	Patchy perihilar and basal infiltration
PFTS	Transfer factor increased



Which of the following tests would you do next?

- A

Anti-nuclear antibodies
- B

Anti-smooth muscle antibodies
- C

P-ANCA
- D

Anti-GBM antibodies
- E

Anti-Ro antibodies

21236

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 76 of 203

Investigations:

Hb	10.9 g/dl
WCC	$12.1 \times 10^9/\text{l}$
PLT	$185 \times 10^9/\text{l}$
Na ⁺	141 mmol/l
K ⁺	5.9 mmol/l
Creatinine	194 $\mu\text{mol/l}$
c-ANCA	Negative
ESR	74 mm/h
CXR	Patchy perihilar and basal infiltration
PFTS	Transfer factor increased

Which of the following tests would you do next?

- | | |
|---|-------------------------------|
| A | Anti-nuclear antibodies |
| B | Anti-smooth muscle antibodies |
| C | P-ANCA |
| D | Anti-GBM antibodies |
| E | Anti-Ro antibodies |

Explanation



- | | |
|---|---------------------|
| D | Anti-GBM antibodies |
|---|---------------------|

A Anti-nuclear antibodies

- The anti-nuclear antibodies would be negative in this case. This test is used to identify connective tissue diseases such as rheumatoid arthritis, systemic lupus erythematosus (SLE), polymyositis and scleroderma. It is an auto-antibody that binds to contents of the cell nucleus.

- B Anti-smooth muscle antibodies

C	P-ANCA
---	--------

- P-ANCA shows a perinuclear staining pattern and the most common p-ANCA target is myeloperoxidase, a neutrophil granule

protein: P-ANCA's associated with ulcerative colitis, eosinophilic granulomatosis with polyangiitis, microscopic polyangiitis and focal necrotising and crescentic glomerulonephritis.

- | | |
|---|--------------------|
| E | Anti-Ro antibodies |
|---|--------------------|

Anti-Ro antibodies are associated with many autoimmune diseases such as SLE, neonatal lupus and primary biliary cirrhosis. The presence of anti-Ro antibodies in pregnant women with SLE is associated with heart block in the fetus.

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	31
Responses Total:	31
Responses - % Correct:	0%

Back to Filters

Question 77 of 203

A 55-year-old man presents to the clinic for review. He has suffered one episode of transient loss of vision in his right eye, and most recently has suffered a mononeuritis affecting his left common peroneal nerve. He also has a history of hypertension for which he currently takes ramipril and amlodipine. On examination his BP is 155/85 mmHg, he has left foot drop consistent with his nerve injury. You notice that he also has livedo reticularis.

Investigations;

Hb	12.2 g/dl
WCC	12.3 x10 ⁹ /l (increased neutrophils)
PLT	192 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.8 mmol/l
Creatinine	194 mol/l
P-ANCA	Positive
ESR	61 mm/hr
Renal angiogram	Multiple microaneurysms

Which of the following is the most likely diagnosis?



- A

Wegener’s granulomatosis
- B

SLE
- C

Polyarteritis nodosa
- D

Berger’s disease
- E

Goodpasture’s syndrome

21237

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 77 of 203

A 55-year-old man presents to the clinic for review. He has suffered one episode of transient loss of vision in his right eye, and most recently has suffered a mononeuritis affecting his left common peroneal nerve. He also has a history of hypertension for which he currently takes ramipril and amlodipine. On examination his BP is 155/85 mmHg, he has left foot drop consistent with his nerve injury. You notice that he also has livedo reticularis.

Investigations;

Hb	12.2 g/dl
WCC	12.3 x10 ⁹ /l (increased neutrophils)
PLT	192 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.8 mmol/l
Creatinine	194 mol/l
P-ANCA	Positive
ESR	61 mm/hr
Renal angiogram	Multiple microaneurysms

Which of the following is the most likely diagnosis?

- A

Wegener’s granulomatosis
- B

SLE
- C

Polyarteritis nodosa
- D

Berger’s disease
- E

Goodpasture’s syndrome

Explanation



- C

Polyarteritis nodosa

Both CNS and PNS symptoms are common presenting features of polyarteritis nodosa (PAN), in which pANCA may be positive but not necessarily specific to the condition. Angiography revealing small to medium-sized artery renal aneurysms is very typical of the condition. Hepatitis B is associated with an increased risk of PAN, and for this reason prevalence is increased in populations where Hep B is endemic. Histology of lesions reveals microscopic polyangitis, which is also seen in Wegener’s granulomatosis. Corticosteroids and cyclophosphamide form the mainstay of therapy for idiopathic PAN. Multiple renal microaneurysms are very typical of PAN, and this pushes us away from a diagnosis of Wegeners, SLE, etc.

- A

Wegener’s granulomatosis

Wegener’s granulomatosis is a small vessel vasculitis characterised by the presence of anti-neutrophil cytoplasmic antibodies (ANCA). The presentation includes fever, weight loss, malaise, myalgia and flitting polyarthralgia. There are usually ear, nose and throat symptoms such as nasal discharge, epistaxis, sinusitis, oral or nasal ulcers and otitis media.

- B

SLE

SLE is a heterogenous multi-system disease characterised by auto-antibody production and impaired immune complex clearance. It is more common in Afro-Caribbean, Hispanic or Asian females in youth or young adulthood. The American College of Rheumatology has established clinical criteria for diagnosis. These include malar rash, photosensitivity, oral ulcers, arthritis, serositis and renal disease.

- D

Berger’s disease

Berger’s disease is otherwise known as IgA nephropathy. It is the most common cause of haematuria in the developed world. The clinical presentation is asymptomatic, with patients being investigated for incidental haematuria on dipstick. The haematuria may occasionally be classically timed with the onset of upper respiratory illnesses (synpharyngeal haematuria). Associated proteinuria is common and it may present with rapidly progressive glomerulonephritis with acute renal failure.

- E

Goodpasture’s syndrome

Goodpasture, also known as anti-GBM disease, occurs because of anti-basement membrane autoantibodies directed against type-IV collagen. HLA-DR2 and HLA-B7 are thought to occur with increased frequency in patients who have Goodpasture. Renal biopsy demonstrates features of crescentic necrotising glomerulonephritis. Worsening haemoptysis due to pulmonary haemorrhage, and acute renal failure, is typical of Goodpasture syndrome.

21237

Rate this question: ⚙️ ⭐ ⭐ ⭐ ⭐ ⭐

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	32
Responses Total:	32
Responses - % Correct:	0%

Back to Filters

Question 78 of 203

A 29-year-old man with a history of polycystic kidney disease presents to the Emergency room with 24hrs of left loin pain. He is managed with ramipril and amlodipine for his BP and has suffered a progressive rise in his creatinine over the past few years. On examination his BP is 142/82 mmHg. The pain gradually resolves in the Emergency room with simple analgesia.

Investigations;

Hb	11.4 g/dl
WCC	5.2 x10 ⁹ /l
PLT	201 x10 ⁹ /l
Na ⁺	141 mmol/l
K ⁺	4.9 mmol/l
Creatinine	156 μmol/l (153 3 months earlier in the clinic)
CRP	12 mg/l
Urine	blood +, protein -

Which of the following is the most appropriate next step?

- A

Admit for observation
- B

Discharge with ciprofloxacin
- C

Arrange an abdominal CT scan
- D

Arrange an abdominal ultrasound scan
- E

Arrange review in 3 days for urine culture results

21239

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 78 of 203

Investigations;

Hb	11.4 g/dl
WCC	5.2 x10 ⁹ /l
PLT	201 x10 ⁹ /l
Na ⁺	141 mmol/l
K ⁺	4.9 mmol/l
Creatinine	156 μ mol/l (153 3 months earlier in the clinic)
CRP	12 mg/l
Urine	blood +, protein -

- **Prevalence** – the proportion of the population with a disease at a particular point in time
- **Incidence** – the number of new cases of a disease in a particular population over a period of time
- **Prevalence** = **Incidence** x **Duration**

- | | |
|---|--|
| A | Admit for observation |
| B | Discharge with ciprofloxacin |
| C | Arrange an abdominal CT scan |
| D | Arrange an abdominal ultrasound scan |
| E | Arrange review in 3 days for urine culture results |

- | | |
|---|--------------------------------------|
| D | Arrange an abdominal ultrasound scan |
|---|--------------------------------------|

A Admit for observation

- Haematuria is usually self-limiting, and as the pain has resolved it seems likely that there is no further bleeding. Therefore, there is no need to keep the patient in for observation. Any investigations can

be carried out as an outpatient.

- B Discharge with ciprofloxacin

As explained above, this man is unlikely to have a urinary tract infection as a cause of the pain because he did not have any urinary symptoms such as dysuria and frequency of urination. You may expect some white cells in the urine if there is a urinary tract infection. As a result, treatment with ciprofloxacin is unlikely to be helpful.

- C Arrange an abdominal CT scan

An abdominal CT will show haemorrhage into the renal cyst. However, the treatment would still be conservation, i.e. analgesia for any pain and rest. Therefore, there is no indication at present to subject the patient to high doses of radiation from a CT scan.

- | | |
|---|--|
| E | Arrange review in 3 days for urine culture results |
|---|--|

This patient did not have any urinary symptoms such as dysuria and frequency of urination. You may expect some white cells in the urine if there is a urinary tract infection. Therefore, this man is more likely to have haemorrhage into a renal cyst for which a review with

Rate this question:

NOTES

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	33
Responses Total:	33
Responses - % Correct:	0%

Back to Filters

Question 79 of 203

A 62-year-old man who has been managed with long-term haemodialysis for end stage renal failure as a result of Type 1 diabetes presents to the clinic complaining of pain and tingling in his hands, particularly in the early hours of the morning. He also has trouble with dysphagia and indigestion, and echocardiography performed for decreased LV function revealed a suspicion of early constrictive pericarditis. On further questioning he also admits to drinking a glass of whisky each evening. On examination he has weakness of thumb abduction, apposition and flexion and there is suspicion of some sensory loss.

Investigations;

Hb	10.1 g/dl
WCC	4.3 x10 ⁹ /l
PLT	192 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.7 mmol/l
Creatinine	421 μmol/l
Urate	0.54 mmol/l
Glucose	9.2 mmol/l

Which of the following is the most likely cause of his upper limb neurological symptoms?

- A

Uraemic neuropathy
- B

Carpal tunnel syndrome
- C

B₁₂ deficiency
- D

Diabetic neuropathy
- E

Alcoholic neuropathy

21274

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 79 of 203

Investigations:

Which of the following is the most likely cause of his upper limb neurological symptoms?

- ### Explanation

Rate this question:

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

[Blog](#) [About Pastest](#) [Contact Us](#) [Help](#)

Back to Filters

Question 80 of 203

A 54-year-old man who is on continuous peritoneal ambulatory dialysis, presents to the Emergency room with dull abdominal pain. He has a history of hypertension for which he takes a number of agents, and Type 2 diabetes for which he takes BD insulin. On examination, he is pyrexial 37.8°C; the dialysate fluid is cloudy, but the site of the catheter looks clean.

Investigations;

Hb	10.2 g/dl
WCC	12.1 x10 ⁹ /l
PLT	204 x10 ⁹ /l
Na ⁺	140 mmol/l
K ⁺	4.9 mmol/l
Creatinine	321 mol/l
Dialysate fluid	120 white cells/ml

Which of the following is the most appropriate initial treatment?

- A

Oral ciprofloxacin
- B

IV gentamicin and vancomycin
- C

Intraperitoneal vancomycin and gentamicin
- D

Oral vancomycin
- E

IV ciprofloxacin

21487

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Question 80 of 203

Investigations;

A	Oral ciprofloxacin
B	IV gentamicin and vancomycin
C	Intraperitoneal vancomycin and gentamicin
D	Oral vancomycin
E	IV ciprofloxacin

This man has symptoms and signs consistent with spontaneous bacterial peritonitis associated with continuous ambulatory peritoneal dialysis (CAPD). Standard therapy involves instilling IP vancomycin and gentamicin, as coagulase negative staphylococcus is a very common pathogen. Once culture results are available antibiotics can then be tailored appropriately. Persistent or relapsing peritonitis, erosion or protrusion of the cuff around the dialysis catheter may unfortunately necessitate removal.

Rate this question:

Previous Question

Tag Question

Feedback

End Session

Peer Responses %

Responses Correct:	0
Responses Incorrect:	35
Responses Total:	35
Responses - % Correct:	0%

Question 81 of 203

Investigations:

Hb	9.9 g/dl
WCC	5.6 x10 ⁹ /l
PLT	184 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	5.7 mmol/l
Urea	12.2 mmol/l
Creatinine	305 μmol/l
Ferritin	130 μg/l (15-300 μg/l)

A	10.0 g/dl
B	11.0 g/dl
C	13.0 g/dl
D	14.0 g/dl
E	14.5 g/dl

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 81 of 203

A 68-year-old man comes to the renal clinic as a GP referral. He has a long history of hypertension which is managed with ramipril 10 mg daily and amlodipine 10 mg, but most recently he has complained of increasing tiredness. On physical examination his blood pressure was 140/80 mmHg, with chest and abdominal examination unremarkable. The GP ran some blood tests and was very concerned about the results.

Investigations:

Hb	9.9 g/dl
WCC	5.6 x10 ⁹ /l
PLT	184 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	5.7 mmol/l
Urea	12.2 mmol/l
Creatinine	305 μmol/l
Ferritin	130 μg/l (15-300 μg/l)

You plan to treat him with EPO once his ferritin is maintained above 200. Which of the following is the most appropriate Hb target to get above?

- A

10.0 g/dl
- B

11.0 g/dl
- C

13.0 g/dl
- D

14.0 g/dl
- E

14.5 g/dl

Explanation

- B

11.0 g/dl

Guidelines from the European Renal Association support a target for Hb of >11.0 g/dl or haematocrit of >33%. Additionally, they recommend reaching this target within 4 months of commencing treatment. Renal patients are often additionally ferritin depleted, and assessment of ferritin level is crucial before starting EPO. Patients who are ferritin depleted should first be given an IV infusion. Some controversy exists as to whether a higher target would lead to lower CV events, but a study sponsored by Roche, which targeted EPO use to achieving an Hb in the normal range, showed no CV benefit.

- A

10.0 g/dl

This target is too low, and therefore if patients were treated to this target there may still be some who experience anaemia symptoms such as shortness of breath and lethargy. Guidelines from the European Renal Association support a target for Hb of >11.0 g/dl or haematocrit of >33%.

- C

13.0 g/dl

This level of haemoglobin is a little too high, with some concern for risk.

- D

14.0 g/dl

The Correction of Haemoglobin and Outcomes in Renal Insufficiency (CHOIR) investigators concluded that a haemoglobin target of 13.5 g/dl was associated with significantly greater risk for the primary composite outcome (death, MI, hospitalization for congestive heart failure without renal replacement therapy, or stroke) compared with a lower target of 11.3 g/dl. Therefore there should be a caution in haemoglobins >13.5g/dl.

- E

14.5 g/dl

As above, the CHOIR investigators concluded that a haemoglobin target of 13.5 g/dl was associated with significantly greater risk for the primary composite outcome (death, MI, hospitalization for congestive heart failure without renal replacement therapy, or stroke) compared with a lower target of 11.3 g/dl. Therefore there should be a caution in haemoglobins greater than 13.5 g/dl.

21529

Rate this question: ⚙️☆☆☆☆☆

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	36
Responses Total:	36
Responses - % Correct:	0%

Question 82 of 203

Investigations:

Hb	10.0 g/dl
WCC	4.8 x10 ⁹ /l
PLT	178 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.9 mmol/l
Creatinine	593 μ mol/l
Ca ⁺⁺ corr	2.61 mmol/l (2.2-2.6 mmol/l)
ALP	295 U/l (45-105 U/l)
PTH	14 pmol/l (1-6 pmol/l)
PO ₄ ³⁻	0.70 mmol/l
X-ray left femur	lucency in mid-shaft
T score	-3.0

10

- | | |
|---|--------------------------|
| A | Dialysis related amyloid |
| B | Multiple myeloma |
| C | Hyperparathyroidism |
| D | Osteomalacia |
| E | Malabsorption |

Submit

Skip Question

Back to Filters

Question 82 of 203

A 50-year-old woman who has a 10-year history of end-stage renal failure and is on haemodialysis is admitted with a pathological fracture of her left femur. She is taking ramipril 10 mg, amlodipine 10 mg, atorvastatin 10 mg, calcium and vitamin D. On examination she has hepatosplenomegaly. Her BP is 155/92 mmHg and she has peripheral oedema. Previously when she was still passing reasonable amounts of urine, her protein excretion was measured at 3.2 g/24 h.

Investigations:

Hb	10.0 g/dl
WCC	4.8 x10 ⁹ /l
PLT	178 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.9 mmol/l
Creatinine	593 μmol/l
Ca ⁺⁺ corr	2.61 mmol/l (2.2-2.6 mmol/l)
ALP	295 U/l (45-105 U/l)
PTH	14 pmol/l (1-6 pmol/l)
PO ₄ ³⁻	0.70 mmol/l
X-ray left femur	lucency in mid-shaft
T score	-3.0

Which of the following is the most likely cause of the fracture?

- A Dialysis related amyloid
- B Multiple myeloma
- C Hyperparathyroidism
- D Osteomalacia
- E Malabsorption

Explanation



- C Hyperparathyroidism

This person has familial renal amyloidosis, as evidenced by hepatosplenomegaly, nephrotic syndrome and progressive renal failure. The fracture is most likely to be related to hyperparathyroidism and loss of bone density. The raised alkaline phosphatase is likely to be as a result of the recent fracture, and the calcium level to slight over-replacement with calcium and vitamin D. Cinacalcet would be the obvious next therapy for this patient.

- A Dialysis related amyloid

Features of amyloid arthropathy on radiography will include juxta-articular soft tissue swelling, mild periarticular osteoporosis, subchondral cystic lesions with well-defined sclerotic margins and normal joint space. These lesions were not apparent in the radiograph in this case.

- B Multiple myeloma

Fractures and skeletal complications are found in up to 80% of patients with myeloma due to the plasma cell dyscrasias. Radiographically there are multiple destructive lytic lesions in the skeleton, due to severe demineralization that characterises multiple myeloma. This is not the case in this scenario.

- D Osteomalacia

Osteomalacia is due to insufficient mineralisation of the osteoid due to vitamin D deficiency or defects in phosphate metabolism. The radiograph shows diffuse demineralisation with the characteristic smudgy, ‘erased’ type of demineralisation with pseudo-fractures (Looser zones).

- E Malabsorption

Malabsorption may lead to pathological fractures because of reduced absorption of vitamin D and calcium essential for bone formation. There is no history of malabsorption, such as diarrhoea or weight loss, in this case. We may expect the calcium level to be low.

21532

Rate this question:

Next Question

- Previous Question
- Tag Question
- Feedback
- End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	37
Responses Total:	37
Responses - % Correct:	0%

Back to Filters

Question 83 of 203

A 60-year-old man presents to the GP with tiredness and stress at work. He is due to retire in 5 years and wants to work out his time, but is being forced to retire early. Past medical history of note includes hypertension which is managed with ramipril. Clinical examination reveals a BP of 142/78 mmHg, otherwise there are no other abnormal findings.

Investigations;

Hb	12.9 g/dl
WCC	7.8 x10 ⁹ /l
PLT	192 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.5 mmol/l
Creatinine	130 mol/l (127 μ mol/l some 2 years earlier)
Plasma Viscosity	1.78 mPa/s (1.50-1.72)
Immunoglobulins	IgG paraprotein band, total IgG elevated at 27g
Bone marrow	plasma cells<10%

Which of the following is the most likely diagnosis?

- A

Multiple myeloma
- B

Monoclonal gammopathy of unknown significance (MGUS)
- C

Hodgkin’s lymphoma
- D

Non-hodgkin’s lymphoma
- E

Waldenstrom’s macrogammaglobulinaema

21540

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 83 of 203

A 60-year-old man presents to the GP with tiredness and stress at work. He is due to retire in 5 years and wants to work out his time, but is being forced to retire early. Past medical history of note includes hypertension which is managed with ramipril. Clinical examination reveals a BP of 142/78 mmHg, otherwise there are no other abnormal findings.

Investigations;

Hb	12.9 g/dl
WCC	7.8 x10 ⁹ /l
PLT	192 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.5 mmol/l
Creatinine	130 mol/l (127 μ mol/l some 2 years earlier)
Plasma Viscosity	1.78 mPa/s (1.50-1.72)
Immunoglobulins	IgG paraprotein band, total IgG elevated at 27g
Bone marrow	plasma cells<10%

Which of the following is the most likely diagnosis?

- A

Multiple myeloma
- B

Monoclonal gammopathy of unknown significance (MGUS)
- C

Hodgkin’s lymphoma
- D

Non-hodgkin’s lymphoma
- E

Waldenstrom’s macrogammaglobulinaema

Explanation

The incidental finding of a paraprotein band, without evidence of end-organ damage, and less than 10% plasma cell involvement of bone marrow is typical of MGUS. Progression to myeloma is around 1% per year, and the vast majority of patients can be managed conservatively. Evidence of deteriorating renal function or impaired immunity may imdicate that transformation to myeloma has occurred.

21540

Rate this question:

⊖

★

★

★

★

★

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	38
Responses Total:	38
Responses - % Correct:	0%

Back to Filters

Question 84 of 203

A 24-year-old man presents with facial swelling. He has recently recovered from an upper respiratory tract infection for which his GP had prescribed a course of amoxicillin 3 days earlier. He was diagnosed with minimal change nephropathy 2 years ago, which was confirmed on biopsy. He isn't currently taking corticosteroids. On examination his BP is 150/88 mmHg and he also has pitting ankle oedema.

Investigations;

Hb	12.6 g/dl
WCC	7.8 x10 ⁹ /l
PLT	195 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.8 mmol/l
Creatinine	130 mol/l
Albumin	25 g/l
Urine protein	+++

Which of the following is the most likely diagnosis?

- A

Post streptococcal glomerulonephritis
- B

Relapse of minimal change disease
- C

Membranous nephropathy
- D

IgA nephropathy
- E

Focal segmental glomerulosclerosis

21578

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 84 of 203

Investigations:

Hb	12.6 g/dl
WCC	7.8 x10 ⁹ /l
PLT	195 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.8 mmol/l
Creatinine	130 μmol/l
Albumin	25 g/l
Urine protein	+++

11

- | | |
|---|---------------------------------------|
| A | Post streptococcal glomerulonephritis |
| B | Relapse of minimal change disease |
| C | Membranous nephropathy |
| D | IgA nephropathy |
| E | Focal segmental glomerulosclerosis |

Explanation

Given the previous diagnosis, a relapse of minimal change disease is the most likely underlying diagnosis here. The interval between streptococcal infection and appearance of symptoms is too short to indicate that this is post-streptococcal glomerulonephritis. Minimal change disease is usually very responsive to corticosteroids, and hypertension should be managed aggressively with ACE inhibitors the first choice therapy.

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	39
Responses Total:	39
Responses - % Correct:	0%

Back to Filters

Question 85 of 203

A 75-year-old woman with a history of Type 2 diabetes presents with symptoms of a urinary tract infection. She takes a mix of medications including gliclazide MR, ramipril, amlodipine and atorvastatin. On examination she has a temperature of 37.8°C, her BP is 125/80 mmHg, her pulse is 75/min. She has suprapubic and left loin tenderness consistent with a UTI.

Investigations reveal;

Hb	12.3 g/dl
WCC	14.1 x10 ⁹ /l
PLT	202 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.9 mmol/l
Creatinine	182 mol/l
Urine	white cells ++, protein ++

Which of the following is the most appropriate therapy according to SIGN?

- A

Oral amoxicillin
- B

IV amoxicillin
- C

Oral ciprofloxacin
- D

Oral trimethoprim
- E

Oral nitrofurantoin

23328

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Question 85 of 203

Investigations reveal;

Hb	12.3 g/dl
WCC	14.1 x10 ⁹ /l
PLT	202 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	4.9 mmol/l
Creatinine	182 μmol/l
Urine	white cells ++, protein ++



- | | |
|---|---------------------|
| A | Oral amoxicillin |
| B | IV amoxicillin |
| C | Oral ciprofloxacin |
| D | Oral trimethoprim |
| E | Oral nitrofurantoin |

Explanation

Significant resistance to amoxicillin exists with respect to urinary tract pathogens, being present in up to 50% of isolates in some areas. Co-amoxiclav effectively overcomes this resistance problem but we are not presented with co-amoxiclav as an option here. Nitrofurantoin may be associated with peripheral neuropathy and is not a first line choice, and fluoroquinolone resistant *E. coli* make up more than 10% of those tested in some areas, although ciprofloxacin has still been recommended as first line therapy by SIGN for symptoms of upper UTI as seen here. Trimethoprim resistance is apparently as high as 40% but it is still recommended as a 3 day course for uncomplicated lower urinary tract infection.

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	40
Responses Total:	40
Responses - % Correct:	0%

Back to Filters

Question 86 of 203

A 65-year-old man with no significant previous medical history presents to his GP with symptoms of an upper respiratory tract infection for which he is prescribed a 10 day course of amoxicillin. Towards the end of the course of antibiotics he comes back to the GP complaining of increasing nausea, anorexia and shortness of breath. He has a BP of 145/95 mmHg and a diffuse rash over his groin and thighs.

Investigations;

Hb	13.1 g/dl
WCC	9.2 x10 ⁹ /l (elevated eosinophil count)
PLT	187 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	6.8 mmol/l
Creatinine	430 mol/l
Urine	Blood +, protein +, WBC +

Which of the following is the most likely cause?

- A

IgA nephropathy
- B

Acute interstitial nephritis
- C

Acute tubular necrosis
- D

Henoch Schonlein purpura
- E

Goodpasture’s syndrome

23362

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Back to Filters

Question 86 of 203

A 65-year-old man with no significant previous medical history presents to his GP with symptoms of an upper respiratory tract infection for which he is prescribed a 10 day course of amoxicillin. Towards the end of the course of antibiotics he comes back to the GP complaining of increasing nausea, anorexia and shortness of breath. He has a BP of 145/95 mmHg and a diffuse rash over his groin and thighs.

Investigations;

Hb	13.1 g/dl
WCC	9.2 x10 ⁹ /l (elevated eosinophil count)
PLT	187 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	6.8 mmol/l
Creatinine	430 mol/l
Urine	Blood +, protein +, WBC +



Which of the following is the most likely cause?

Your answer was correct

- A

IgA nephropathy
- B

Acute interstitial nephritis
- C

Acute tubular necrosis
- D

Henoch Schonlein purpura
- E

Goodpasture’s syndrome

Explanation

AIN is an important cause of renal failure and may be the cause of renal failure in up to 15% of patients undergoing renal biopsy evaluation. Sulphonamides, penicillins, cephalosporins, diuretics and NSAIDS are all recognised causes. Discontinuation of the possible causative agent is the treatment of choice, and slow recovery is possible over the course of around 6 weeks. It is possible however that up to 40% of patients may progress to chronic renal failure. High dose prednisolone may improve the outcome when used in the acute situation. The findings of elevated creatinine, peripheral blood eosinophilia and red cells, protein and white cells in the urine all support the underlying diagnosis.

IgA nephropathy usually follows a benign course, with no significant rise in BP or creatinine in the majority of cases, out of keeping with what is seen here. Acute tubular necrosis is associated with a significant period of renal hypoperfusion, which may for example be related to sepsis or haemorrhage. Henoch Schonlein Purpura does not fit with the eosinophilia seen here, and Goodpasture’s is associated with pulmonary haemorrhage.

23362

Rate this question:

⊖

★

★

★

★

★

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	1
Responses Incorrect:	40
Responses Total:	41
Responses - % Correct:	2%

Question 87 of 203

Investigations show:

Creatinine	1200 mol/l
ECG	left ventricular hypertrophy. No other abnormality
Hb	9.5 g/dl
Potassium	5.8 mmol/l
Urea	60 mmol/l
WCC	$25 \times 10^9/l$

-
-
-

- | | |
|---|----------------------|
| A | Rapid IV Fluids |
| B | Fluid restriction |
| C | Dialysis |
| D | Blood transfusion |
| E | Insulin and dextrose |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 87 of 203

A 56-year-old woman with known chronic renal failure (usual creatinine 400 μmol/l) secondary to adult polycystic kidney disease presents with a 2-day history of dysuria, left loin pain, fevers and vomiting. On examination she is pyrexial (temperature 38.5° C) with left loin tenderness and a pericardial rub. Mucosal surfaces and skin turgor look normal. Her blood pressure is elevated at 162/90 mmHg. You commence her on IV antibiotics.

Investigations show:

Creatinine	1200 μmol/l
ECG	left ventricular hypertrophy. No other abnormality
Hb	9.5 g/dl
Potassium	5.8 mmol/l
Urea	60 mmol/l
WCC	25 x 10 ⁹ /l

Which one of the following is the most appropriate intervention with respect to her renal impairment and circulatory status?

- A

Rapid IV Fluids
- B

Fluid restriction
- C

Dialysis
- D

Blood transfusion
- E

Insulin and dextrose

Explanation

This woman demonstrates a case of acute-on-chronic renal failure (CRF) secondary to likely pyelonephritis/infected cyst. She needs dialysis urgently as she has a pericardial rub, which is most likely due to uraemic pericarditis. She also has severe uraemia with a urea and creatinine value as high as this. Although cautious IV fluids would not be inappropriate, she is only mildly dehydrated and they will not correct this degree of renal failure. Insulin and dextrose will lower the potassium but this is not dangerously elevated, and dialysis, required for the severe renal impairment and uraemic pericarditis will result in lowering of potassium anyway. A blood transfusion is not needed as many patients with CRF are anaemic secondary to erythropoietin deficiency and this is usually worse in the setting of acute infection.

32381

Rate this question:      

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	1
Responses Incorrect:	41
Responses Total:	42
Responses - % Correct:	2%

Back to Filters

Question 88 of 203

A 50-year-old retired bank manager attended a ‘well man’ clinic. He has always enjoyed good health, but has recently developed a dry cough. He takes daily over-the-counter multivitamin tablets. On examination he is extremely well with no abnormality detected. Screening test results are normal with the exception of corrected calcium of 2.82 mmol/l. A repeat uncuffed sample with simultaneous plasma parathyroid hormone (PTH) reveals corrected calcium of 2.86 mmol/l and PTH of 5.4 pmol/l (0.9–5.4).

Which of the following is the most likely diagnosis?

- A

Primary hyperparathyroidism
- B

Multiple myeloma
- C

Prolonged tourniquet application
- D

Vitamin D excess
- E

Sarcoidosis

32382

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 88 of 203

A 50-year-old retired bank manager attended a ‘well man’ clinic. He has always enjoyed good health, but has recently developed a dry cough. He takes daily over-the-counter multivitamin tablets. On examination he is extremely well with no abnormality detected. Screening test results are normal with the exception of corrected calcium of 2.82 mmol/l. A repeat uncuffed sample with simultaneous plasma parathyroid hormone (PTH) reveals corrected calcium of 2.86 mmol/l and PTH of 5.4 pmol/l (0.9–5.4).

Which of the following is the most likely diagnosis?

A	Primary hyperparathyroidism
B	Multiple myeloma
C	Prolonged tourniquet application
D	Vitamin D excess
E	Sarcoidosis

Explanation

The plasma parathyroid hormone (PTH) is the best single test for differentiating between hypercalcaemia due to parathyroid over activity and that due to non-parathyroid causes. Primary hyperparathyroidism can be associated with a plasma PTH within the normal range. The normal response to hypercalcaemia is suppression of the plasma PTH below the lower reference limit. Levels within the reference limit are inappropriately high and suggestive of parathyroid disease. Familial hypocalciuric hypercalcaemia is a hereditary disease with autosomal dominant transmission, and is characterised by moderate chronic hypercalcaemia and normal or moderately elevated plasma PTH concentrations. The fractional excretion of calcium is lower than in primary hyperparathyroidism and the urine calcium/creatinine ratio is usually less than 0.01. The importance of this condition is that the hypercalcaemia rarely leads to significant clinical signs and parathyroidectomy is unsuccessful as a therapy.

32382

Rate this question:      

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	1
Responses Incorrect:	42
Responses Total:	43
Responses - % Correct:	2%

Back to Filters

Question 89 of 203

A 27-year-old dancer comes to the emergency department because of severe pain in her left side and dark urine. She denies fever but states that her general practitioner (GP) had seen her several days ago for urinary frequency and burning. She had been treated with trimethoprim at that time and the dysuria had resolved. The pain she now feels comes on very suddenly and the paroxysms are so severe that she cannot walk. There is some nausea and she had vomited twice. The pain radiates to her groin. Her vital signs are temperature of 37.2°C, blood pressure 110/70 mmHg, pulse 95/min and respiration 14/min. Physical examination of the chest and heart is normal. The abdomen is soft and non-tender with no guarding. Her white count and ESR are in the normal range. She is given analgesia.

Which of the following is the most appropriate early treatment?

- A

Intravenous (IV) broad-spectrum antibiotics
- B

IV Furosemide
- C

Percutaneous removal of stones
- D

Lithotripsy
- E

Hydration

32383

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 89 of 203

A 27-year-old dancer comes to the emergency department because of severe pain in her left side and dark urine. She denies fever but states that her general practitioner (GP) had seen her several days ago for urinary frequency and burning. She had been treated with trimethoprim at that time and the dysuria had resolved. The pain she now feels comes on very suddenly and the paroxysms are so severe that she cannot walk. There is some nausea and she had vomited twice. The pain radiates to her groin. Her vital signs are temperature of 37.2°C, blood pressure 110/70 mmHg, pulse 95/min and respiration 14/min. Physical examination of the chest and heart is normal. The abdomen is soft and non-tender with no guarding. Her white count and ESR are in the normal range. She is given analgesia.

Which of the following is the most appropriate early treatment?

- A

Intravenous (IV) broad-spectrum antibiotics
- B

IV Furosemide
- C

Percutaneous removal of stones
- D

Lithotripsy
- E

Hydration

Explanation

Kidney stones form when there is a high level of calcium (hypercalciuria), oxalate (hyperoxaluria), or uric acid (hyperuricosuria) in the urine; a lack of citrate in the urine; or insufficient water in the kidneys to dissolve waste products. The kidneys must maintain an adequate amount of water in the body to remove waste products. If dehydration occurs, high levels of substances that do not dissolve completely (e.g. calcium, oxalate, uric acid) may form crystals that slowly build up into kidney stones. Urine normally contains chemicals (including citrate, magnesium and pyrophosphate) that prevent the formation of crystals. Low levels of these inhibitors can contribute to the formation of kidney stones. Of these, citrate is thought to be the most important. Approximately 80-85% of stones pass spontaneously. Approximately 20% of patients require hospital admission because of unrelenting pain, inability to retain enteral fluids, proximal urinary tract infection (UTI), or inability to pass the stone. A ureteral stone associated with obstruction and upper UTI is a true urological emergency. Complications include perinephric abscess, urosepsis, and death. Immediate involvement of the urologist is essential. Diagnosis Includes: blood tests, urine analysis, X-rays, intravenous urogram, and ultrasound scan. Treatment depends on the type and cause of the stone. The usual initial treatment is hydration and analgesia which allows spontaneous passage of stones in the majority of patients. For those patients whose stones do not pass, methods of stone removal including percutaneous and lithotripsy approaches may be considered.

32383

Rate this question: ⚙️ ⭐️ ⭐️ ⭐️ ⭐️ ⭐️

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	1
Responses Incorrect:	43
Responses Total:	44
Responses - % Correct:	2%

Back to Filters

Question 90 of 203

A 19-year-old man comes to your office referred by his rugby coach. He is usually in robust, excellent health and is in the middle of 3 weeks of intensive training before a University tour to Australia and New Zealand. He has been passing dark urine for the past few days and he is worried that he might have a serious illness. He describes the urine as cola-coloured. He denies fever and has no complaint or discomfort other than the dark urine. He is a large muscular young man who is in no obvious distress. His temperature is 37.4°C, his blood pressure is 120/70 mmHg and pulse is 65/min. His extremities are non oedematous and non-tender. The remainder of the physical examination is unremarkable. Full blood count is normal.

Which test would be the most appropriate for initial work up?

- A

ESR
- B

Urinalysis
- C

Kidney, ureter and bladder (KUB) x-ray
- D

Renal ultrasound
- E

Computed tomography (CT) of the abdomen

32384

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 90 of 203

A 19-year-old man comes to your office referred by his rugby coach. He is usually in robust, excellent health and is in the middle of 3 weeks of intensive training before a University tour to Australia and New Zealand. He has been passing dark urine for the past few days and he is worried that he might have a serious illness. He describes the urine as cola-coloured. He denies fever and has no complaint or discomfort other than the dark urine. He is a large muscular young man who is in no obvious distress. His temperature is 37.4°C, his blood pressure is 120/70 mmHg and pulse is 65/min. His extremities are non oedematous and non-tender. The remainder of the physical examination is unremarkable. Full blood count is normal.

Which test would be the most appropriate for initial work up?

A	ESR
B	Urinalysis
C	Kidney, ureter and bladder (KUB) x-ray
D	Renal ultrasound
E	Computed tomography (CT) of the abdomen

Explanation

Urinalysis

The differential diagnosis of haematuria includes stones, hematological disorders such as thrombocytopenia, infection, tumours, trauma and treatment, like cyclophosphamide. In this case the patient is at risk for rhabdomyolysis from the vigorous exercise he has been undergoing. Rhabdomyolysis would give haemoglobin on stick urinalysis but no red blood cells would be seen on microscopy. Other causes of haematuria in a young man without pain, who is fit and healthy, such as renal stones or urological malignancy, are highly unlikely. Creatinine kinase may be raised. No specific treatment is required apart from maintaining adequate hydration.

32384

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	1
Responses Incorrect:	44
Responses Total:	45
Responses - % Correct:	2%

Question 91 of 203

His blood results are:

Hb	11.6 g/dl
ESR	78 mm/hr
Albumin	23 g/l
Ca ²⁺ (corrected)	3.24 mmol/l
Na ⁺	153 mmol/l

A	Corticosteroids
B	Thiazide diuretics
C	Hydration with normal saline/5% dextrose alternating bags
D	Furosemide IV
E	Calcitonin nitrate intravenously

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 91 of 203

A 68-year-old man complained of weight loss, polyuria, chronic cough and left shoulder pain. Over the past few days he has suffered from increasing nausea and vomiting. He is an ex-smoker who smoked 15 cigarettes/day for 20 years. On examination he looks dehydrated, his BP is 112/72 mmHg, pulse is 85/min and regular. There are coarse crackles on auscultation of the chest consistent with COPD. He is thin, with a BMI of 21.

His blood results are:

Hb	11.6 g/dl
ESR	78 mm/hr
Albumin	23 g/l
Ca ²⁺ (corrected)	3.24 mmol/l
Na ⁺	153 mmol/l



Which initial step is most appropriate to begin to correct his hypercalcaemia?

- A

Corticosteroids
- B

Thiazide diuretics
- C

Hydration with normal saline/5% dextrose alternating bags
- D

Furosemide IV
- E

Calcitonin nitrate intravenously

Explanation

Hydration with normal saline/5% dextrose alternating bags

Hypercalcaemia is due to PTH related peptide secreted by the tumour cells. The calcium can be reduced in a relatively short time by rehydrating with IV fluids. Furosemide is calciuric, but should initially be avoided until he is rehydrated adequately. IV bisphosphonates are very effective, but work over a few hours. Corticosteroids have variable efficacy, but may be added to IV fluid and bisphosphonates. Calcitonin is rarely given and not usually as initial therapy.

32385

Rate this question:

Next Question

- Previous Question

Feedback
- Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	1
Responses Incorrect:	45
Responses Total:	46
Responses - % Correct:	2%

Question 92 of 203

Her blood results showed:

Na ⁺	136 mmol/l
K ⁺	2.8 mmol/l
Urea	5 mmol/l
Bicarbonate	32 mmol/l
pH	7.5

11

- | | |
|---|--------------------------|
| A | Addison's disease |
| B | Eating disorder |
| C | Long-term diuretic abuse |
| D | Conn's syndrome |
| E | Cushing's syndrome |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 92 of 203

An 18-year-old A-level student comes to the clinic complaining of amenorrhea. There is no history of weight loss. She has been recently referred to the dentist for assessment of badly eroded teeth. On examination, her parotid glands are enlarged and scratch marks are noticed over the dorsum of the knuckles. The rest of the examination is normal. She is an active girl, who runs five miles daily. Her BMI is 17. Her BP is 105/70 mmHg.

Her blood results showed:

Na ⁺	136 mmol/l
K ⁺	2.8 mmol/l
Urea	5 mmol/l
Bicarbonate	32 mmol/l
pH	7.5

Which of the following conditions is the most likely diagnosis?

≡

- A

Addison’s disease
- B

Eating disorder
- C

Long-term diuretic abuse
- D

Conn’s syndrome
- E

Cushing’s syndrome

Explanation

The acid-base disturbance shown is a metabolic alkalosis. Recurrent vomiting +/- laxative abuse may well have lead to the hypokalaemia and raised bicarbonate seen here, the knuckle callusing also supporting the diagnosis. Long-term diuretic abuse may coexist, but given her BMI is only 17, it is vomiting that is the likely predominant factor. Hypokalaemia is not a feature of Addison’s and Cushing’s is effectively ruled out by her BMI. Conn’s is associated with hypertension.

32386

Rate this question:      

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	1
Responses Incorrect:	46
Responses Total:	47
Responses - % Correct:	2%

Question 93 of 203

Electrolyte results are as follows:

Na+	138 mmol/l
K+	5.5 mmol/l
Urea	6.0 mmol/l
Creatinine	150 μmol/l
HbA1c	54.10 mmol/mol (7.1%) (9.29-53.0 mmol/mol, 3-7%)

g

- | | |
|---|--|
| A | A non-progressive rise in creatinine of 20-30% is acceptable |
| B | Lisinopril should be withdrawn immediately given the creatinine is 150 |
| C | Cough is much less likely to occur versus angiotensin receptor blockers |
| D | Renal artery stenosis is likely if creatinine rises 15% and is then stable |
| E | Creatinine should be first checked 3 months after initiation |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 93 of 203

A 50-year-old female with a 4-year history of type 2 diabetes was referred to the renal clinic. She was commenced on 5 mg of lisinopril 5 weeks ago for hypertension. Otherwise she is well, without any symptoms. Her diabetes is controlled with BD mixed insulin. On examination her BP is 142/81 mmHg, pulse is 75/min and regular. There is minor loss of sensation affecting both feet.

Electrolyte results are as follows:

Na+	138 mmol/l
K+	5.5 mmol/l
Urea	6.0 mmol/l
Creatinine	150 mol/l
HbA1c	54.10 mmol/mol (7.1%) (9.29-53.0 mmol/mol, 3-7%)

Which of the following fits best with her treatment after starting angiotensin-converting enzyme (ACE) inhibitor therapy?

- A

A non-progressive rise in creatinine of 20-30% is acceptable
- B

Lisinopril should be withdrawn immediately given the creatinine is 150
- C

Cough is much less likely to occur versus angiotensin receptor blockers
- D

Renal artery stenosis is likely if creatinine rises 15% and is then stable
- E

Creatinine should be first checked 3 months after initiation

Explanation

A non-progressive rise in creatinine of 20-30% is acceptable

With regards to angiotensin-converting enzyme (ACE) inhibitors, a non-progressive rise in creatinine of 20-30% is acceptable. This is observed in almost all intervention trials with respect to the renin angiotensin system, and is associated with an initial reduction in BP, it has no prognostic significance. Diabetes is the main cause of chronic kidney disease, and a significant proportion of patients with Type 2 diabetes actually have evidence of complications at the time of first presentation with hyperglycaemia. A progressive rise in creatinine should prompt consideration of renal artery stenosis. Guidelines generally suggest checking serum creatinine 1 week, 1 month and then periodically after starting ACE inhibition.

32387

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	1
Responses Incorrect:	47
Responses Total:	48
Responses - % Correct:	2%

Back to Filters

Question 94 of 203

A 29-year-old female of Jamaican origin presented to her general practitioner with 4 months of amenorrhoea and increasing tiredness. Routine examination reveals a blood pressure of 210/135 mmHg and a palpable mass in the abdomen. She was sent for a routine blood test and ultrasound of the abdomen and pelvis. She was adopted as a child and recalls that her immediate brother died of an uncertain kidney disease a few years ago in Jamaica.

Investigation results are as follows:

Ultrasound scan	Enlarged kidneys with multiple cysts, consistent with polycystic kidneys
Prolactin	1200 mU/l (<360 mU/l)
Hb	14.5 g/dl
WCC	9×10^9 /l
Platelets	350×10^9 /l
Na ⁺	140 mmol/l
K ⁺	6.2 mmol/l
Urea	16.8 mmol/l
Creatinine	750 μmol/l

Which of the following is the most likely diagnosis?

- A

Autosomal dominant polycystic kidney disease
- B

Chronic ureteric reflux
- C

Renal cell carcinoma
- D

Renovascular disease
- E

Transitional cell carcinoma

32388

Submit

Previous Question

Skip Question

Calculator

✔

Normal Values

✔

Back to Filters

Question 94 of 203

A 29-year-old female of Jamaican origin presented to her general practitioner with 4 months of amenorrhoea and increasing tiredness. Routine examination reveals a blood pressure of 210/135 mmHg and a palpable mass in the abdomen. She was sent for a routine blood test and ultrasound of the abdomen and pelvis. She was adopted as a child and recalls that her immediate brother died of an uncertain kidney disease a few years ago in Jamaica.

Investigation results are as follows:

Ultrasound scan	Enlarged kidneys with multiple cysts, consistent with polycystic kidneys
Prolactin	1200 mU/l (<360 mU/l)
Hb	14.5 g/dl
WCC	9 × 10 ⁹ /l
Platelets	350 × 10 ⁹ /l
Na ⁺	140 mmol/l
K ⁺	6.2 mmol/l
Urea	16.8 mmol/l
Creatinine	750 mol/l

Which of the following is the most likely diagnosis?

- A

Autosomal dominant polycystic kidney disease
- B

Chronic ureteric reflux
- C

Renal cell carcinoma
- D

Renovascular disease
- E

Transitional cell carcinoma

Explanation

Autosomal dominant polycystic kidney disease (APKD) presents with hypertension and haematuria, and later in middle age with chronic renal failure. Acute loin pain or haematuria due to haemorrhage into the cysts are common. Cysts may be found in other organs such as the liver and ovaries, but liver cysts are more commonly associated with childhood autosomal recessive polycystic kidney disease. Hypertension develops in over 75% of cases. Subarachnoid haemorrhage from berry aneurysm is reported to occur in about 9% of patients, and 8% of patients with APKD are thought to have asymptomatic intracranial aneurysms. Early aggressive management of blood pressure is the cornerstone of therapy.

32388

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	1
Responses Incorrect:	48
Responses Total:	49
Responses - % Correct:	2%

Back to Filters

Question 95 of 203

A-50-year old female received a cadaveric renal transplant 8 weeks ago for diabetic nephropathy. She attended the transplant follow-up clinic where it was discovered that the serum creatinine had risen to 150 μ mol/l from having been within the normal range. You understand that her GP prescribed an antibiotic for a respiratory tract infection during the last 2 weeks and she is penicillin allergic. Her BP is 135/72 mmHg, and her transplant scar is healing well.

Which one of the following is most likely to explain the rise in creatinine?

- A

Acute ciclosporin toxicity
- B

Chronic graft rejection
- C

Acute graft rejection
- D

Vascular thrombosis of the transplant vessels
- E

Cytomegalovirus infection

32389

Submit

Previous Question

Skip Question

Calculator

✔

Normal Values

✔

Back to Filters

Question 95 of 203

A-50-year old female received a cadaveric renal transplant 8 weeks ago for diabetic nephropathy. She attended the transplant follow-up clinic where it was discovered that the serum creatinine had risen to 150 μmol/l from having been within the normal range. You understand that her GP prescribed an antibiotic for a respiratory tract infection during the last 2 weeks and she is penicillin allergic. Her BP is 135/72 mmHg, and her transplant scar is healing well.

Which one of the following is most likely to explain the rise in creatinine?

A	Acute ciclosporin toxicity
B	Chronic graft rejection
C	Acute graft rejection
D	Vascular thrombosis of the transplant vessels
E	Cytomegalovirus infection

Explanation

Acute ciclosporin toxicity

Causes of graft dysfunction up to 4 months after transplantation are:

- acute rejection: the risk is greater up to 2weeks post-operatively, it occurs in about 30-50% of all transplants
- ciclosporin toxicity: occurs either as acute or chronic in otherwise well patients
- acute tubular necrosis (ATN) of the graft
- vascular thrombosis
- ureteric leakage from the anastomosis

Given the recent history of antibiotic use for respiratory tract infection in a patient who is penicillin allergic, it’s likely she was exposed to erythromycin or clarithromycin. These two macrolides are potent inhibitors of CYP3A4, and as such lead to elevated levels of ciclosporin. Given its narrow therapeutic window, a relatively small elevation in ciclosporin can result in renal toxicity. Erythromycin and clarithromycin should be avoided in patients treated with ciclosporin if at all possible. Azithromycin is associated with a lesser degree of 3A4 inhibition.

32389

Rate this question: ⚙️ ⭐️ ⭐️ ⭐️ ⭐️ ⭐️

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	1
Responses Incorrect:	49
Responses Total:	50
Responses - % Correct:	2%

Back to Filters

Question 96 of 203

A 28-year-old patient was admitted with a 3-day history of oliguria, dyspnoea, nausea and vomiting. On examination she is pyrexial 37.6° C, BP is 165/92 mmHg, pulse is 95/min and regular. There are scattered crackles on auscultation of her chest. Abdomen is soft but generally tender as a result of the vomiting. Initial investigations showed acute renal failure with urea of 22 mmol/l and creatinine of 400 μmol/l. An ultrasound and renal biopsy were ordered.

Two days later the following results were obtained:

Na ⁺	136 mmol/l
K ⁺	5.8 mmol/l
Urea	28 mmol/l
Creatinine	560 μmol/l
Hb	8.0 g/dl
WCC	12 × 10 ⁹ /l
Platelets	160 × 10 ⁹ /l
Renal biopsy	Focal and segmental necrosis with epithelial cell crescents

What possible underlying diagnosis is most likely?

≡

- A

Polyarteritis nodosa
- B

Immunoglobulin A (IgA) nephropathy
- C

Hypertensive nephropathy
- D

Anti-GBM (glomerular basement membrane) glomerulonephritis
- E

Membranous nephropathy

32390

Submit

Previous Question

Skip Question

Calculator

✔

Normal Values

✔

Back to Filters

Question 96 of 203

A 28-year-old patient was admitted with a 3-day history of oliguria, dyspnoea, nausea and vomiting. On examination she is pyrexial 37.6° C, BP is 165/92 mmHg, pulse is 95/min and regular. There are scattered crackles on auscultation of her chest. Abdomen is soft but generally tender as a result of the vomiting. Initial investigations showed acute renal failure with urea of 22 mmol/l and creatinine of 400 μmol/l. An ultrasound and renal biopsy were ordered.

Two days later the following results were obtained:

Na ⁺	136 mmol/l
K ⁺	5.8 mmol/l
Urea	28 mmol/l
Creatinine	560 μmol/l
Hb	8.0 g/dl
WCC	12 × 10 ⁹ /l
Platelets	160 × 10 ⁹ /l
Renal biopsy	Focal and segmental necrosis with epithelial cell crescents

What possible underlying diagnosis is most likely?



- A

Polyarteritis nodosa
- B

Immunoglobulin A (IgA) nephropathy
- C

Hypertensive nephropathy
- D

Anti-GBM (glomerular basement membrane) glomerulonephritis
- E

Membranous nephropathy

Explanation

Anti-GBM (glomerular basement membrane) glomerulonephritis

The clinical picture along with the results represents a classical rapidly progressive glomerulonephritis (RPGN). The disease is characterised by fulminant renal failure and pathologically by focal and segmental necrosis of the glomeruli with epithelial cell proliferation (crescents).

The clinical presentation may be similar to other glomerulonephritis, but the progression of renal failure is rapid.

Classification of RPGN is as follows:

- pauci-immune RPGN: microscopic polyangiitis, Wegener’s granulomatosis, idiopathic crescentic glomerulonephritis
- immune complex RPGN: systemic lupus erythematosus, glomerulonephritis, cryoglobulinaemia, infective endocarditis, hepatitis B
- anti-GBM mediated RPGN: anti-GBM glomerulonephritis (without lung haemorrhage), Goodpasture’s disease (with lung haemorrhage)

Management is with high dose intravenous steroids and a second line agent such as cyclophosphamide.

32390

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	1
Responses Incorrect:	50
Responses Total:	51
Responses - % Correct:	2%

Back to Filters

Question 97 of 203

A 36-year-old farmer was referred by her general practitioner (GP) to the General Medical Clinic. She originally presented to the emergency department a few weeks ago with frank transient haematuria. This settled soon after presentation but recurred some days later for a short period. She was seen following this by her GP. She has been complaining of some fatigue and also occasional dizziness occurring over the last 6 months. In addition she complains of some unsteadiness and has lost her balance a number of times recently leading to some falls. She is otherwise well although she had some visual problems around 5 years ago for which she was briefly under the care of an ophthalmologist. She was told that she had ‘cysts’ in her eyes and was offered further investigations at this point which she declined. She is an only child and her mother died aged 40 years, from widespread cancer although she does not know the primary site. She has one child aged four and is trying to conceive at present. She has a normal cardiovascular and respiratory examination. Abdominal examination reveals no masses. Neurological examination reveals mild ataxia of gait and on finger nose testing. Fundoscopy demonstrates several abnormal accumulations of vessels on the retina but no papilloedema.

Blood tests from her Emergency Department admission are as follows:

Hb	17.4 g/dl
WCC	11.9 × 10 ⁹ /l
Platelets	590 × 10 ⁹ /l
MCV	95 fl
Na ⁺	139 mmol/l
K ⁺	4.0 mmol/l
Urea	4.5 mmol/l
Creatinine	65 mol/l
Albumin	36 g/l
ALT	34 IU/l
Bilirubin	14 mol/l
ALP	105 IU/l

Her urine is clear with no frank haematuria, however urinalysis demonstrates:

Blood	3+
Protein	1+
Nitrites	Negative
Glucose	Negative

Which of the following would be the next most important step in investigating her?

- A

24-h urinary vanillylmandelic acid (VMA) measurement
- B

Urinary 5-hydroxyindole-3-acetic acid (5-HIAA) measurement
- C

Ultra-violet light (Wood lamp) examination of her skin
- D

Magnetic resonance imaging (MRI) of the brain
- E

TSC1 and TSC2 gene analysis for tuberous sclerosis

32391

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 97 of 203

Fundoscopy demonstrates several abnormal accumulations of vessels on the retina but no papilloedema.

Blood tests from her Emergency Department admission are as

demonstrates:

[illegible]

- ### Explanation

Magnetic resonance imaging (MRI) of the brain

This lady has symptoms and signs suggestive of urothelial malignancy. It is therefore essential in her initial management that this is investigated further and treated accordingly. Computed tomography (CT) of the abdomen and pelvis with contrast is likely to form part of these investigations. Contraceptive advice is important as this lady is trying to conceive and may have a malignant disease as well as an underlying genetic condition. This lady has features suggestive of von Hippel-Lindau (VHL) syndrome and in view of this and her ataxia definitive imaging of the central nervous system (CNS) is required. In view of the common occurrence of posterior fossa haemangioblastomas a magnetic resonance imaging scan (MRI) of the brain will be most useful in order to reach diagnosis and to decide on any management that needs to follow on from this. VHL syndrome is an autosomal dominant condition resulting from a mutation in the VHL gene, a tumour suppression gene on chromosome 3 leading to retinal haemangiomas, CNS haemangioblastomas, pheochromocytoma, renal cell carcinoma, renal and pancreatic cysts. She has a family history of renal tract carcinoma with a history of ophthalmological lesions. In addition she has features that might relate to a posterior fossa space-occupying brain lesion. She has a polycythaemia which may be secondary to a renal cell carcinoma. She will also require as part of her work up: VHL gene analysis, 24-h urinary vanillylmandelic acid (VMA) (in view of the possibility of pheochromocytoma), and ophthalmology review, but this can wait until the diagnosis is confirmed.

Tuberous sclerosis is another condition that can be inherited in an autosomal dominant fashion, although up to 50 % of cases are thought to be the result of new mutations. It shares some similarity to VHL syndrome in that it has intracranial, ophthalmological and renal manifestations. Patients often have learning difficulties and seizures from an early age with intracranial calcification. Retinal hamartomas can develop. They are predisposed to the development of giant intracranial astrocytomas. They have characteristic skin manifestations with facial angiofibroma, subungual fibromas and ash leaf patches (which can be identified by use of UV light). They are also predisposed to renal cysts and angioliomata, as well as cardiac and pulmonary tumours.

100

Peer Responses %

Session Progress

Back to Filters

Question 98 of 203

A 73-year-old lady has been admitted from clinic for investigation of suspected intra-abdominal malignancy. She has diabetes, congestive cardiac failure and mild renal impairment (serum creatinine 130 μ mol/l). She is taking gliclazide 80 mg/day, furosemide 20 mg od and diclofenac 75 mg bd for back pain thought to be related to neoplastic disease. An abdominal ultrasound has been performed which has shown no definite lesions. An abdominal and pelvic computed tomography (CT) scan with contrast is planned.

Which one of the following is the most important step to reduce the chance of contrast-induced nephrotoxicity?

- A

Oral sodium bicarbonate 325 mg/day for a day prior to the procedure and for at least two days following

≡
- B

Use of large volume contrast media
- C

Oral acetylcysteine 600 mg bd from the day before the procedure and for two days following
- D

Use of high-osmolality contrast media
- E

Stopping the furosemide and diclofenac at least 48 h prior to the planned procedure

32392

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Question 98 of 203

Which one of the following is the most important step to reduce the chance of contrast-induced nephrotoxicity?

A	Oral sodium bicarbonate 325 mg/day for a day prior to the procedure and for at least two days following
B	Use of large volume contrast media
C	Oral acetylcysteine 600 mg bd from the day before the procedure and for two days following
D	Use of high-osmolality contrast media
E	Stopping the furosemide and diclofenac at least 48 h prior to the planned procedure

Stopping the furosemide and diclofenac at least 48 h prior to the planned procedure

Stopping nephrotoxic medications including diclofenac at least 48 h prior to the procedure should reduce the risk the risk of contrast nephropathy. In addition furosemide may lead to fluid depletion, which increases the chances of nephrotoxicity. The patient will have to be monitored closely for signs of worsening heart failure if this medication is stopped, and monitoring of urea and electrolyte concentrations will be very important. Although intravenous hydration does reduce the risk of contrast nephropathy, its use in this lady and the risk of precipitating an acute exacerbation of her cardiac failure probably make the risk/benefit analysis favour caution. There is no evidence that sodium bicarbonate reduces the chance of nephrotoxicity. There is nothing to suggest that changing to an insulin sliding scale to improve glycaemic control can reduce the risk, and indeed such a move risks precipitating hypoglycaemia and wide blood sugar variations if it is not monitored closely.

Rate this question:

Next Question

[Previous Question](#)

[Tag Question](#)

[Feedback](#)

[End Session](#)

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	1
Responses Incorrect:	52
Responses Total:	53
Responses - % Correct:	2%

Question 99 of 203

Urgent blood results are as follows:

- **Stressors** are external factors that trigger a stress response.

In his subsequent management which one of the following steps is most likely to improve his outcome?

A	Urinary acidification
B	Further intravenous calcium supplementation
C	Continuous or intermittent renal replacement therapy (haemodialysis) if urine output does not remain adequate
D	'renal dose' dopamine infusion if oliguric in spite of rehydration
E	Loop diuretic use to maintain urine output once rehydrated

Submit

Skip Question

Question 99 of 203

Urgent blood results are as follows:

In his subsequent management which one of the following steps is most likely to improve his outcome?

The answer is Continuous or intermittent renal replacement therapy (haemodialysis) if urine output does not remain adequate -

Rate this question:

[Next Question](#)

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	1
Responses Incorrect:	53
Responses Total:	54
Responses - % Correct:	2%

Back to Filters

Question 100 of 203

A 45-year-old man presents to the emergency department with excruciating unilateral abdominal pain radiating from his left loin to his perineum. The pain is waxing and waning in character and when it is at its most severe he is calling out and screaming. He is managed with intravenous fluids and analgesia after it is established that his ureters are not dilated. His symptoms settle. However, during investigations he is noted to have impaired renal function and bilateral renal medullary calcification.

Which of the following is a recognised cause of renal medullary calcification?

- A

Primary hyperparathyroidism
- B

Hypercalcaemia secondary to metastatic malignancy
- C

Renal tubular acidosis type 4
- D

Chronic glomerulonephritis
- E

Hypothyroidism

32394

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Question 100 of 203

Which of the following is a recognised cause of renal medullary calcification?

A	Primary hyperparathyroidism
B	Hypercalcaemia secondary to metastatic malignancy
C	Renal tubular acidosis type 4
D	Chronic glomerulonephritis
E	Hypothyroidism

This man has presented with renal colic presumably secondary to his medullary renal calcification. Primary hyperparathyroidism (along with other causes of chronic hypercalcaemic states) is a recognised cause of medullary calcification. PTH, calcium and phosphate, along with renal function are important laboratory tests in his workup.

Type 1 renal tubular acidosis leads to poorly acidified urine and predisposes to renal tract calcification. The medullary sponge kidney is also a recognised cause of renal medullary calcification. Hypercalcaemia in the context of metastatic malignancy is usually too short-lived to lead to renal medullary calcification. Chronic glomerulonephritis does predispose to cortical but not to medullary nephrocalcinosis, as well as renal cortical necrosis. Hyper- but not hypothyroidism is associated with this condition. Vitamin D excess but not deficiency can lead to medullary calcinosis.

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	1
Responses Incorrect:	54
Responses Total:	55
Responses - % Correct:	2%

Back to Filters

Question 101 of 203

A 72-year-old woman who has undergone haemodialysis for 8 years presents with numbness and tingling in both hands which wakes her from sleep in the early hours. In addition she has had trouble holding items between her fingers and thumb recently. She also complains of dysphagia. On examination her BP is 155/82 mmHg, pulse is 75/min, BMI is 23. There is wasting of the thenar eminence on both hands and numbness over the palmar aspect of the thumb index and middle fingers.

Which one of the following measures is most appropriate in this patient?

- A

Corticosteroid therapy
- B

Non-steroidal anti-inflammatory drugs
- C

Incision of the flexor retinaculae
- D

Switch to low-flux dialysis membranes
- E

Ciclosporin therapy

32395

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 101 of 203

Which one of the following measures is most appropriate in this patient?

Explanation

End Session

Peer Responses %

Responses Correct:	0
Responses Incorrect:	1
Responses Total:	1
Responses - % Correct:	0%

Back to Filters

Question 102 of 203

A 71-year-old man is referred to the renal clinic with a creatinine concentration of 195 μ mol/l. There is a past medical history of renal colic on two previous occasions. Additional history includes three presentations to his general practitioner with an acute arthritis affecting his first MTP joint on the right foot. He takes ramipril 10 mg daily for his hypertension. On examination, a number of tophi are noted, particularly around the cartilaginous portion of the external auditory meatus. His blood pressure is 155/75 mmHg.

Which one of the following would be the most appropriate intervention with respect to his blood pressure control?

- A

Bendroflumethiazide 2.5 mg
- B

Amlodipine 5 mg
- C

Allopurinol 600 mg
- D

Furosemide 40 mg
- E

Diclofenac 50 mg

32396

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Back to Filters

Question 102 of 203

A 71-year-old man is referred to the renal clinic with a creatinine concentration of 195 μmol/l. There is a past medical history of renal colic on two previous occasions. Additional history includes three presentations to his general practitioner with an acute arthritis affecting his first MTP joint on the right foot. He takes ramipril 10 mg daily for his hypertension. On examination, a number of tophi are noted, particularly around the cartilaginous portion of the external auditory meatus. His blood pressure is 155/75 mmHg.

Which one of the following would be the most appropriate intervention with respect to his blood pressure control?

- A

Bendroflumethiazide 2.5 mg
- B

Amlodipine 5 mg
- C

Allopurinol 600 mg
- D

Furosemide 40 mg
- E

Diclofenac 50 mg

Explanation

Amlodipine 5 mg

This patient is most likely to be suffering from urate nephropathy as evidenced by the attacks of gout and his presentations with renal colic. Addition of diuretics is likely to worsen his serum urate levels; therefore the most appropriate additional antihypertensive from the choices above is amlodipine 5 mg. A dose of allopurinol as high as 600 mg would not be appropriate until an estimate of creatinine clearance is made. A starting dose of 200 mg would be correct. Other appropriate measures include increasing fluid intake and considering restriction of dietary purines (chiefly animal in origin). Urate nephropathy is around four times more common in men, and may occur acutely in patients undergoing chemotherapy for haematological malignancy, for which rasburicase has emerged as an effective newer therapy.

32396

Rate this question: ⚙️☆☆☆☆☆

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	2
Responses Total:	2
Responses - % Correct:	0%

Back to Filters

Question 103 of 203

A 72-year-old woman with a long history of chronic back pain presents for review. Past medical history of note includes osteoporosis of the lumbar spine and she takes paracetamol, tramadol and diclofenac for pain relief. Review of her notes reveals a number of abnormally high BP recordings.

Investigations reveal:

Blood pressure	155/90 mmHg
Hb	10.2g/dl
Creatinine	195 mol/l
24 h urinary protein	1.4 g
Urine culture	White cells but no growth

Which one of the following is the most appropriate intervention in this case?

- A

Stop diclofenac
- B

Prednisolone 40 mg daily
- C

Cyclophosphamide
- D

Stop paracetamol
- E

Bendroflumethiazide 2.5 mg daily

32397

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Question 103 of 203

Investigations reveal:

Blood pressure	155/90 mmHg
Hb	10.2g/dl
Creatinine	195 mol/l
24 h urinary protein	1.4 g
Urine culture	White cells but no growth

Which one of the following is the most appropriate intervention in this case?

- | | |
|---|----------------------------------|
| A | Stop diclofenac |
| B | Prednisolone 40 mg daily |
| C | Cyclophosphamide |
| D | Stop paracetamol |
| E | Bendroflumethiazide 2.5 mg daily |

Explanation

This patient is most likely to be suffering from chronic interstitial nephritis as a result of long-term use of non-steroidal anti-inflammatory agents. Diagnosis is made clinically with a history of heavy analgesic use and supported by the presence of papillary microcalcification in computed tomography (CT) scan. Treatment involves withdrawal of the offending analgesic agent and aggressive control of blood pressure. Corticosteroids and other immune-modulating agents are not indicated in this condition. Long-term follow-up of these patients shows progression to renal replacement therapy in some cases and an increased risk of urothelial malignancy. An ACE inhibitor or ARB would be the initial treatment of choice for blood pressure control.

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	3
Responses Total:	3
Responses - % Correct:	0%

Back to Filters

Question 104 of 203

A 34-year-old man signs on with a new general practitioner and is noted to be hypertensive at 160/90 mmHg. Proteinuria in the microalbuminuric range is noted and one plus of haematuria; there is no past history of diabetes mellitus. His father died of a brain haemorrhage at the age of 42 years. Other past medical history of note includes two presentations with loin pain. On examination there are bilateral irregularly enlarged kidneys, the right larger than the left.

Which one of the following would be the most appropriate initial investigation in this case?

- A

Renal tract ultrasound scan
- B

CT of the abdomen
- C

Intravenous urography
- D

24 h urinary protein collection
- E

Cardiac echo

32398

Submit

Previous Question

Skip Question

Calculator

✔

Normal Values

✔

Back to Filters

Question 104 of 203

A 34-year-old man signs on with a new general practitioner and is noted to be hypertensive at 160/90 mmHg. Proteinuria in the microalbuminuric range is noted and one plus of haematuria; there is no past history of diabetes mellitus. His father died of a brain haemorrhage at the age of 42 years. Other past medical history of note includes two presentations with loin pain. On examination there are bilateral irregularly enlarged kidneys, the right larger than the left.

Which one of the following would be the most appropriate initial investigation in this case?

- A

Renal tract ultrasound scan
- B

CT of the abdomen
- C

Intravenous urography
- D

24 h urinary protein collection
- E

Cardiac echo

Explanation

Renal tract ultrasound scan

This history, examination findings and investigations are highly suggestive of autosomal dominant polycystic kidney disease. Ultrasound scanning is highly effective in demonstrating renal cysts. The family history of probable subarachnoid haemorrhage means that cerebral circulation aneurysms should be excluded with a magnetic resonance angiogram of the cerebral circulation. Other co-existent conditions may be colonic diverticulae, demonstrable with barium enema, and mitral valve prolapse, seen on cardiac echo. Medical management centres on control of blood pressure, pain from enlarged kidneys and treatment of urinary tract infections which occur with increased frequency in patients with adult polycystic kidney disease.

32398

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	4
Responses Total:	4
Responses - % Correct:	0%

Back to Filters

Question 105 of 203

A 65-year-old woman is admitted as an emergency, following her renal function having been assessed one month after starting ramipril therapy for hypertension. Examination in the clinic reveals abdominal and carotid bruits; she is clinically euvolaemic. Her blood pressure is 156/88 mmHg. ECG on admission is unremarkable.

Investigations reveal:

On admission:

Na ⁺	142 mmol/l
K ⁺	6.0 mmol/l
Creatinine	340 mol/l

One month previously:

Na ⁺	140 mmol/l
K ⁺	4.2 mmol/l
Creatinine	175 mol/l

Which one of the following represents the most appropriate initial step in the management of this patient?

- A

Start calcium resonium
- B

Observe renal function over the next week
- C

Add bendroflumethiazide 2.5 mg po daily
- D

Stop ramipril
- E

Advise a low potassium diet

32454

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 105 of 203

A 65-year-old woman is admitted as an emergency, following her renal function having been assessed one month after starting ramipril therapy for hypertension. Examination in the clinic reveals abdominal and carotid bruits; she is clinically euvolaemic. Her blood pressure is 156/88 mmHg. ECG on admission is unremarkable.

Investigations reveal:

On admission:

Na ⁺	142 mmol/l
K ⁺	6.0 mmol/l
Creatinine	340 mol/l

One month previously:

Na ⁺	140 mmol/l
K ⁺	4.2 mmol/l
Creatinine	175 mol/l

Which one of the following represents the most appropriate initial step in the management of this patient?

- A

Start calcium resonium
- B

Observe renal function over the next week
- C

Add bendroflumethiazide 2.5 mg po daily
- D

Stop ramipril
- E

Advise a low potassium diet

Explanation

This clinical scenario is highly suggestive of underlying renal artery stenosis. Angiotensin-converting enzyme (ACE) inhibitors result in a decrease in efferent renal arteriolar blood pressure and a deterioration in serum creatinine in patients with renal artery stenosis. Magnetic resonance angiography is the definitive investigation. The key intervention for this patient is to stop the ramipril and substitute another antihypertensive drug, such as a calcium channel antagonist if required. Depending on local protocol, most physicians check renal function 1 week and 1 month after initiating ACE inhibitors or angiotensin II receptor blockade. Her potassium is likely to fall within a day or two of stopping the ramipril therapy; given the fact that her ECG on admission is unremarkable no acute intervention for her potassium is warranted here.

32454

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	5
Responses Total:	5
Responses - % Correct:	0%

Back to Filters

Question 106 of 203

A 58-year-old woman presents with lethargy, confusion and vomiting. She has not presented to health services for many years, although she has been known to use paracetamol/dextropropoxyphene (coproxamol) tablets for arthritis in the past and takes a diuretic for management of hypertension. On examination her blood pressure is 165/85 mmHg and she is confused, with evidence of nystagmus, hyper-reflexia and clonus. General examination reveals evidence of vaginal bleeding.

Blood tests reveal:

Urea	48.2 mmol/l
Creatinine	845 mol/l
Potassium	6.2 mmol/l

Which one of the following diagnoses fit best with this clinical picture?

- A

Hypertensive encephalopathy
- B

Uraemic encephalopathy
- C

Intracranial metastases
- D

Opiate intoxication
- E

Subdural haematoma

32455

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Question 106 of 203

Blood tests reveal:

Urea	48.2 mmol/l
Creatinine	845 μmol/l
Potassium	6.2 mmol/l

三

- | | |
|---|-----------------------------|
| A | Hypertensive encephalopathy |
| B | Uraemic encephalopathy |
| C | Intracranial metastases |
| D | Opiate intoxication |
| E | Subdural haematoma |

Explanation

Uraemic encephalopathy

The evidence of vaginal bleeding is suggestive of uterine or cervical cancer. Cervical cancer is much more likely to lead to ureteric obstruction and this type of presentation with renal failure. The neurological findings are suggestive of uraemic encephalopathy, which progresses to coma, seizures and death if left untreated. The standard therapy for uraemic encephalopathy is initiation of dialysis, although a long-term commitment to renal replacement therapy should be considered carefully in the light of possible underlying advanced cervical carcinoma. Conventionally in this case urgent imaging would be considered to elucidate the extent of any underlying tumour and consequent obstruction before necessarily committing to dialysis.

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	6
Responses Total:	6
Responses - % Correct:	0%

Back to Filters

Question 107 of 203

A 42-year-old homosexual male presents for review. He admits to unprotected sex with a number of males and has suffered a dry cough, shortness of breath, and weight loss over the past few months. His major current complaint is that has had progressive lower limb swelling which has worsened over the past few weeks. Blood pressure on examination is 135/75 mmHg and there is peripheral oedema. His BMI is 21.

Investigations reveal:

24 h urinary protein excretion	5.2 g
Haemoglobin	10.9 g/dl
CD4 count	175/ l (>500)
Albumin	27 g
Na ⁺	138 mmol/l
K ⁺	4.2 mmol/l
Creatinine	195 mol/l
Urea	12.2 mmol/l



Given the likeliest diagnosis from these data, which one of the following would represent the most appropriate treatment for this patient?

- A

Initiate antiretroviral drugs
- B

Initiate corticosteroids
- C

Initiate ciclosporin
- D

Intiate cyclophosphamide
- E

Initiate furosemide

32456

Submit

Previous QuestionSkip Question

Calculator



Normal Values



Back to Filters

Question 107 of 203

A 42-year-old homosexual male presents for review. He admits to unprotected sex with a number of males and has suffered a dry cough, shortness of breath, and weight loss over the past few months. His major current complaint is that has had progressive lower limb swelling which has worsened over the past few weeks. Blood pressure on examination is 135/75 mmHg and there is peripheral oedema. His BMI is 21.

Investigations reveal:

24 h urinary protein excretion	5.2 g
Haemoglobin	10.9 g/dl
CD4 count	175/ l (>500)
Albumin	27 g
Na ⁺	138 mmol/l
K ⁺	4.2 mmol/l
Creatinine	195 mol/l
Urea	12.2 mmol/l



Given the likeliest diagnosis from these data, which one of the following would represent the most appropriate treatment for this patient?

- A

Initiate antiretroviral drugs
- B

Initiate corticosteroids
- C

Initiate ciclosporin
- D

Intiate cyclophosphamide
- E

Initiate furosemide

Explanation

This patient has HIV-associated nephropathy (HIVAN), characterised by nephrotic range proteinuria, normal blood pressure, normal or increased kidney size on ultrasound scan and focal segmental glomerulosclerosis on renal biopsy. Treatment includes aggressive antiretroviral therapy and angiotensin-converting enzyme (ACE) inhibitor therapy for patients where this is not contraindicated due to hyperkalaemia. No completed randomised controlled trials exist for the use of second line agents such as ciclosporin or corticosteroids, although observational studies suggest that a benefit may exist.

32456

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	7
Responses Total:	7
Responses - % Correct:	0%

Back to Filters

Question 108 of 203

A 31-year-old woman with proven diabetic nephropathy as demonstrated by microalbuminuria and elevated creatinine presents to the clinic 12 weeks pregnant. HbA1c has been relatively satisfactory in the past few years, averaging 55.19 mmol/mol (7.2%), and the last recorded creatinine was 145 μmol/l. She has already stopped her ACE inhibitor but wants to know about the prognosis with respect to her kidney disease.

Which one of the following best describes the prognosis with respect to her diabetic nephropathy during pregnancy?



- | | |
|---|--|
| A | She has a 10% chance of worsening renal function |
| B | There is little chance of a worsening of her renal function |
| C | She has an 80% chance of worsening renal function |
| D | She has a 45% chance of worsening renal function |
| E | She should be advised to undergo a termination as she is likely to develop acute renal failure |

32457

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Question 108 of 203

Which one of the following best describes the prognosis with respect to her diabetic nephropathy during pregnancy?

- | | |
|---|--|
| A | She has a 10% chance of worsening renal function |
| B | There is little chance of a worsening of her renal function |
| C | She has an 80% chance of worsening renal function |
| D | She has a 45% chance of worsening renal function |
| E | She should be advised to undergo a termination as she is likely to develop acute renal failure |

A recent case series demonstrated that there is worsening of renal function in up to 45% of patients who get pregnant with pre-existing diabetic nephropathy. In a general case series, Katz et al demonstrated worsening renal function to end-stage renal disease in 16% of patients with mild renal impairment (creatinine < 150 μmol/l), worsening to end-stage renal disease in 20% with creatinine 150-240 μmol/l, and worsening to end-stage renal disease in 45% where creatinine is above 240 μmol/l. Worsening hypertension is strongly associated with the degree of renal progression. A generally poorer prognosis during pregnancy is associated with lupus as the underlying cause of renal impairment.

Rate this question:      

End Session

Peer Responses %

Responses Correct:	0
Responses Incorrect:	8
Responses Total:	8
Responses - % Correct:	0%

Back to Filters

Question 109 of 203

A 71-year-old woman is admitted with general deterioration and ‘off-legs’. On examination in the emergency department she looks dry and uraemic. There are peaked T waves on the 12 lead ECG.

Investigations reveal:

K ⁺	7.2 mmol/l
Na ⁺	145 mmol/l
Creatinine	512 mol/l
Urea	36.8 mmol/l



She is immediately put on a cardiac monitor.

Which one of the following measures is next most important in her immediate management?

- A

IV Insulin and dextrose infusion
- B

Salbutamol nebuliser
- C

IV Calcium gluconate
- D

Calcium resonium
- E

IV Saline infusion

32458

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 109 of 203

Investigations reveal:

3

Which one of the following measures is next most important in her immediate management?

A	IV Insulin and dextrose infusion
B	Salbutamol nebuliser
C	IV Calcium gluconate
D	Calcium resonium
E	IV Saline infusion

IV Calcium gluconate

Rate this question:

Next Question

[Previous Question](#)

[Tag Question](#)

[Feedback](#)

[End Session](#)

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	9
Responses Total:	9
Responses - % Correct:	0%

Back to Filters

Question 110 of 203

A 17-year-old boy presents to his general practitioner complaining of macroscopic haematuria which began 48 hours after he started with a recent respiratory tract infection. On examination his urine is negative to culture, and contains large amounts of red blood cells and some protein but not in the macroalbuminuric range. Microscopy also identifies leucocytes and red cell casts. Renal biopsy reveals diffuse mesangial proliferation and extracellular matrix expansion. A few necrotising lesions with crescent formation are noted. Serum creatinine is 155 μ mol/l and blood pressure is 142/86 mmHg.

Which one of the following statements fits best with the management of this condition?

- A

Prednisolone therapy is indicated as initial therapy

≡
- B

Ciclosporin should be added early to a corticosteroid regime
- C

Two or three agents are required as chronic therapy to control blood pressure in the majority of patients
- D

Calcium channel antagonists are the preferred agents for treating hypertension
- E

Low-protein diets should be used to reduce proteinuria

32459

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Question 110 of 203

Which one of the following statements fits best with the management of this condition?

A	Prednisolone therapy is indicated as initial therapy
B	Ciclosporin should be added early to a corticosteroid regime
C	Two or three agents are required as chronic therapy to control blood pressure in the majority of patients
D	Calcium channel antagonists are the preferred agents for treating hypertension
E	Low-protein diets should be used to reduce proteinuria

This young man has immunoglobulin A (IgA) nephropathy as illustrated by the presentation with haematuria post-respiratory tract infection and the histological findings on renal biopsy. Gross proteinuria is rare, but haematuria is the commonest form of presentation. The disease is commoner in males, and presents most often in the second and third decades. Management includes control of hypertension, with angiotensin-converting enzyme (ACE) inhibitors the preferred agent, and use of corticosteroids. Where there is crescentic nephritis on biopsy, intravenous pulsed corticosteroids are employed, with the addition of cyclophosphamide. Thankfully, progression to end-stage renal disease is uncommon in this condition. Where end-stage renal disease does occur, renal transplantation carries a high chance of success.

Rate this question:

End Session

Peer Responses %

Responses Correct:	0
Responses Incorrect:	10
Responses Total:	10
Responses - % Correct:	0%

Back to Filters

Question 111 of 203

A 78-year-old man presents with tiredness and lethargy accompanied by weight loss of 1 stone over the past 4 months, and severe lower back pain. He has a history of hypertension which is managed with ramipril and amlodipine, and an inferior myocardial infarction some 7 years earlier. On examination his BP is 142/72 mmHg, pulse is 82/min and regular, his BMI is 21. He looks pale and emaciated.

Investigations reveal:

Hb	8.9 g/dl (normocytic)
Platelets	125 × 10 ⁹ /l
WCC	8.5 × 10 ⁹ /l
Na ⁺	135 mmol/l
K ⁺	4.6 mmol/l
Creatinine	245 μmol/l
Ca ²⁺	2.95 mmol/l
Albumin	31 g/l
ESR	85 mm/h
Renal biopsy	Cast nephropathy



Given the likely underlying diagnosis, which one of the following treatments is likely to be an initial therapy in this case?

- A

Methotrexate
- B

Interferon gamma
- C

Non-steroidal anti-inflammatory agents
- D

Interferon alpha
- E

Melphalan

32460

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 111 of 203

Investigations reveal:

Hb	8.9 g/dl (normocytic)
Platelets	125 × 10 ⁹ /l
WCC	8.5 × 10 ⁹ /l
Na ⁺	135 mmol/l
K ⁺	4.6 mmol/l
Creatinine	245 μmol/l
Ca ²⁺	2.95 mmol/l
Albumin	31 g/l
ESR	85 mm/h
Renal biopsy	Cast nephropathy

A	Methotrexate
B	Interferon gamma
C	Non-steroidal anti-inflammatory agents
D	Interferon alpha
E	Melphalan

This man has a clinical picture which is highly suggestive of myeloma with renal involvement. Diagnosis of myeloma relies on plasma electrophoresis and bone marrow biopsy to assess plasma cell infiltration. Cast nephropathy is classically associated with myeloma of the kidney and is characterised by eosinophilic, dense, homogeneous casts that are often fractured or laminated and are partially surrounded by multinucleated foreign body-type giant cells. Congo red-positive casts have been reported in a few cases. Intratubular light chains apparently may undergo alteration in situ, resulting in amyloid formation. Conventional treatment of myeloma typically involves a regime including melphalan and corticosteroids. An aggressive regime involving the VAMP combination of agents, or CVAMP may be considered in younger patients with better initial functional status.

Rate this question:

End Session

Peer Responses %

Responses Correct:	0
Responses Incorrect:	11
Responses Total:	11
Responses - % Correct:	0%

Question 112 of 203

Investigations show:

Diuretic treatment is commenced; which of the following is the other most important step?

Skip Question

Back to Filters

Question 112 of 203

A 63-year-old male patient is admitted to the ward with generalised oedema. He has a background of bronchial malignancy, treated with chemotherapy, and chronic obstructive pulmonary disease and tells you that the swelling has gradually increased over the past few weeks. Examination reveals oedema of the legs, arms and face.

Investigations show:

Blood pressure	145/75 mmHg
Hb	13.6 g/dl
WCC	$4.7 \times 10^9/l$
PLT	$315 \times 10^9/l$
Na ⁺	146 mmol/l
K ⁺	4.3 mmol/l
Urea	7.2 mmol/l
Creatinine	63 mol/l
INR	1.1
Bilirubin	15 mol/l
AST	21 U/l
ALP	70 U/l
Albumin	24 g/l
24-h urinary protein collection	3.6 g/24 h (<0.2)
Renal biopsy	shows membranous glomerulonephritis

Diuretic treatment is commenced; which of the following is the other most important step?



- A

High protein diet
- B

Intravenous albumin
- C

High dose prednisolone
- D

Prophylactic antibiotics
- E

Prophylactic LMW heparin

Explanation

The answer is Prophylactic LMW heparin -

This patient has nephrotic syndrome secondary to membranous glomerulonephritis, which is associated with bronchial malignancies. The best initial management strategies are to restrict dietary sodium and commence diuretics, which is likely to be useful in controlling oedema and blood pressure. With nephrotic syndrome there is an increased risk of thromboembolism (related to loss of anti-thrombin III), and this patient also has a malignancy, hence anticoagulation is indicated. High-protein diet is not beneficial and albumin intravenously produces only a transient rise in serum levels. High-dose prednisolone is useful in minimal change disease but is not usually effective here. Although nephrotic syndrome confers an increased risk of infection, prophylactic antibiotics are not recommended, however, vaccinations (e.g. the pneumococcal vaccine) can be used. Immunosuppressants are not indicated, at least not initially in this patient.

32461

Rate this question:

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	12
Responses Total:	12
Responses - % Correct:	0%

Back to Filters

Question 113 of 203

A 42-year-old patient presents with generalised oedema. He has a background history of intravenous drug use and excess alcohol intake. He smokes 20 cigarettes/day. He suffers from depression and mild asthma and is currently on fluoxetine, and has progressively lost weight and energy over the past few weeks and months.

Initial investigations reveal:

Na ⁺	146 mmol/l
K ⁺	4.1 mmol/l
Urea	3.4 mmol/l
Creatinine	103 mol/l
Albumin	23 g/l
24-h urinary protein collection	3.6 g/24h (<0.2)
Renal biopsy	shows focal segmental glomerulosclerosis

In this patient what is the most likely associations with this disease?



- A

Hepatitis A
- B

Hepatitis C
- C

HIV
- D

Alcohol abuse
- E

Cigarette smoking

32462

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 113 of 203

A 42-year-old patient presents with generalised oedema. He has a background history of intravenous drug use and excess alcohol intake. He smokes 20 cigarettes/day. He suffers from depression and mild asthma and is currently on fluoxetine, and has progressively lost weight and energy over the past few weeks and months.

Initial investigations reveal:

Na ⁺	146 mmol/l
K ⁺	4.1 mmol/l
Urea	3.4 mmol/l
Creatinine	103 mol/l
Albumin	23 g/l
24-h urinary protein collection	3.6 g/24h (<0.2)
Renal biopsy	shows focal segmental glomerulosclerosis

In this patient what is the most likely associations with this disease?



- A

Hepatitis A
- B

Hepatitis C
- C

HIV
- D

Alcohol abuse
- E

Cigarette smoking

Explanation

Focal segmental glomerulosclerosis (FSGC) can present with nephrotic syndrome. It has been associated with HIV, Hepatitis B, intravenous heroin use and massive obesity. The outcome varies and some patients will progress to end-stage renal disease.

With nephrotic syndrome he is at risk of thromboembolism, infection (especially pneumococcus), and hyperlipidaemia. In HIV-related FSGS, highly active retroviral therapy (HAART) is associated with a reduction in proteinuria.

Other measures include following a low salt diet and use of ACE inhibitors. Where heroin addiction persists, discontinuation of heroin may also result in a reduction in proteinuria.

32462

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	13
Responses Total:	13
Responses - % Correct:	0%

Back to Filters

Question 114 of 203

A 52-year-old gentleman is referred by the general practitioner with painless macroscopic haematuria. He is a smoker and is on bendroflumethiazide for hypertension. He takes no other medication and has no other significant past medical history. On examination his BP is 138/78 mmHg, pulse is 75/min and regular. There are changes consistent COPD on auscultation of his chest. His BMI is 23 and there are no masses on palpation of the abdomen. Initial investigations reveal:

Hb	12.2 g/dl
WCC	7 × 10 ⁹ /l
PLT	310 × 10 ⁹ /l
Na ⁺	138 mmol/l
K ⁺	3.8 mmol/l
Urea	8.2 mmol/l
Creatinine	109 mol/l
Urine dipstick	Blood +++
INR	1.0



What one of the following is the most appropriate next step to determine the diagnosis?

- A

Urine culture
- B

Computerised tomography of the renal tract with contrast
- C

Flexible cystoscopy
- D

Retrograde urography
- E

USS renal tract

32463

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 114 of 203

Initial investigations reveal:

Hb	12.2 g/dl
WCC	$7 \times 10^9/l$
PLT	$310 \times 10^9/l$
Na ⁺	138 mmol/l
K ⁺	3.8 mmol/l
Urea	8.2 mmol/l
Creatinine	109 μmol/l
Urine dipstick	Blood +++
INR	1.0

- | | |
|---|--|
| A | Urine culture |
| B | Computerised tomography of the renal tract with contrast |
| C | Flexible cystoscopy |
| D | Retrograde urography |
| E | USS renal tract |

Computed tomography of the renal tract with contrast

With painless macroscopic haematuria in a 52-year-old patient, a neoplastic process has to be ruled out. Initially the best investigations are urine microscopy to look for malignant cells and contrast CT to rule out upper urinary tract pathology. If there is no evidence of upper urinary tract pathology then cystoscopy and further imaging are required to elucidate the cause. Smoking is of course associated with increased risk of transitional cell carcinoma, as are occupational activities such as work in the plastics and dye industries.

Rate this question:

Next Question

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	14
Responses Total:	14
Responses - % Correct:	0%

Question 115 of 203

Investigations reveal:



A	Anti-double stranded DNA antibodies
B	Serial blood cultures
C	Rheumatoid factor
D	Anticardiolipin antibodies
E	Computerised tomography of the thorax and abdomen

Submit

Skip Question

Question 115 of 203

Investigations reveal:



A	Anti-double stranded DNA antibodies
B	Serial blood cultures
C	Rheumatoid factor
D	Anticardiolipin antibodies
E	Computerised tomography of the thorax and abdomen

This patient has hypocomplementaemia and glomerulonephritis with features in the history and examination suggesting infective endocarditis as the aetiology. The blood results suggest an underlying infection with the raised white cell count and C-reactive protein. The erythrocyte sedimentation rate, an indicator for inflammation, is also raised. Serial blood cultures from different sites and an echocardiogram are the next investigations of choice. She should be managed with IV flucloxacillin and gentamicin and there is no need to wait for investigations before commencing therapy.

Rate this question:

Previous Question

Tag Question

Feedback

End Session

Peer Responses %

Responses Correct:	0
Responses Incorrect:	15
Responses Total:	15
Responses - % Correct:	0%

Back to Filters

Question 116 of 203

A 36-year-old woman, a mother of three, presents with a history of generalised weakness of 1 month’s duration as well as cough productive of blood-tinged sputum. A 7-day course of erythromycin has not resulted in any improvement.

On examination she appears pale and mildly dyspnoeic. The respiratory rate is 25/min; BP 165/95 mmHg; pulse 90/min. There are fine inspiratory crackles up to the mid-zones in both lung fields. The jugular venous pressure (JVP) is not elevated and there is a trace of pedal oedema. The remainder of physical examination is unremarkable.

Investigations:

Hb	7.2 g/dl
MCV	56 fl
MCH	19 pg
WCC	10.1 × 10 ⁹ /litre with 75% neutrophils
PLT	223 × 10 ⁹ /litre
Urea	14.2 mmol/litre
Creatinine	165 mol/litre
Na ⁺	135 mmol/litre
K ⁺	4.3 mmol/litre
Cl ⁻	99 mmol/litre
HCO ₃ ⁻	26 mmol/litre
Urine	Protein ++; RBC ++ with RBC casts
CXR	Alveolar shadows up to mid-zones in both lung fields
Renal ultrasound	normal-sized kidneys
Renal biopsy	crescents in up to 50% of glomeruli, and linear deposition of IgG along basement membrane



Which of the following is an essential treatment?

- A

IV furosemide
- B

IV immunoglobulins
- C

IV methylprednisolone and cyclophosphamide
- D

Chloroquine
- E

Infliximab

32474

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Back to Filters

Question 116 of 203

A 36-year-old woman, a mother of three, presents with a history of generalised weakness of 1 month’s duration as well as cough productive of blood-tinged sputum. A 7-day course of erythromycin has not resulted in any improvement.

On examination she appears pale and mildly dyspnoeic. The respiratory rate is 25/min; BP 165/95 mmHg; pulse 90/min. There are fine inspiratory crackles up to the mid-zones in both lung fields. The jugular venous pressure (JVP) is not elevated and there is a trace of pedal oedema. The remainder of physical examination is unremarkable.

Investigations:

Hb	7.2 g/dl
MCV	56 fl
MCH	19 pg
WCC	10.1 × 10 ⁹ /litre with 75% neutrophils
PLT	223 × 10 ⁹ /litre
Urea	14.2 mmol/litre
Creatinine	165 mol/litre
Na ⁺	135 mmol/litre
K ⁺	4.3 mmol/litre
Cl ⁻	99 mmol/litre
HCO ₃ ⁻	26 mmol/litre
Urine	Protein ++; RBC ++ with RBC casts
CXR	Alveolar shadows up to mid-zones in both lung fields
Renal ultrasound	normal-sized kidneys
Renal biopsy	crescents in up to 50% of glomeruli, and linear deposition of IgG along basement membrane

Which of the following is an essential treatment?

- A

IV furosemide
- B

IV immunoglobulins
- C

IV methylprednisolone and cyclophosphamide
- D

Chloroquine
- E

Infliximab

Explanation

This patient has acute glomerulonephritis secondary to Goodpasture’s syndrome, as evident by the presence of RBC casts, crescents on renal biopsy and linear IgG staining. The haemoptysis and pulmonary infiltrates imply the presence of pulmonary haemorrhage.

The treatment of choice in these patients is a combination of plasmapheresis, which removes the circulating autoantibodies and inflammatory cytokines such as complement, and the combination of methylprednisolone and cyclophosphamide to prevent new antibody formation. Prognosis is poor if timely therapy is not instituted. Case reports of successful use of rituximab in refractory Goodpasture’s indicate that it may have a role, although randomised controlled trial evidence is as yet unavailable. On the other hand infliximab does not appear to be useful in this condition.

32474

Rate this question: ⓪☆☆☆☆☆

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	16
Responses Total:	16
Responses - % Correct:	0%

Question 117 of 203

Investigation results are below:

What of the following is the most likely diagnosis?

Skip Question

Back to Filters

Question 117 of 203

A 68-year-old woman is admitted with a 4-day history of profuse bloody diarrhoea, vomiting and lower abdominal pain. She had a past medical history of well-controlled hypertension and mild asthma, but her health had been good until the onset of the presenting illness. She had no recent foreign travel and did not smoke or drink. On examination she was drowsy, but oriented when roused. Her temperature was 39°C, pulse 110/min and regular, blood pressure 102/60 mmHg, and she was clinically dehydrated. Cardiorespiratory examination was unremarkable. Her abdomen was soft and diffusely tender with scanty bowel sounds and no palpable masses. There were no focal neurological signs or rashes. Investigation results are below:

Hb	8.0 g/dl
WCC	26.0 × 10 ⁹ /l
Platelets	48 × 10 ⁹ /l
MCV	84 fl
Na ⁺	147 mmol/l
K ⁺	5.8 mmol/l
Urea	25 mmol/l
Creatinine	780 mmol/l
Bilirubin	32 μmol/l
AST	34 IU/l
ALP	80 IU/l
GGT	54 U/l
Urinalysis	Protein ++ blood ++
Abdominal radiograph	No dilated bowel loops
Chest radiograph	Clear lung fields, no subdiaphragmatic air

What of the following is the most likely diagnosis?



- A

Campylobacter
- B

E coli
- C

Norovirus
- D

Salmonellosis
- E

Staphylococcus aureus gastroenteritis

Explanation

E coli infection

The diagnosis is haemolytic uraemic syndrome (HUS), which causes a microangiopathic haemolysis with thrombocytopenia. HUS is similar to thrombotic thrombocytopaenic purpura (TTP), but there are a number of differences between the two conditions. HUS usually presents in childhood or old age, with acute renal failure following a diarrhoeal illness, whereas TTP is more common in young adults with predominantly neurological features, such as fits and confusion. There is a strong association with verotoxin producing E. coli 0157 and this organism should be looked for specifically in the stool culture. A normal coagulation screen will exclude disseminated intravascular coagulation which can present similarly. The blood film will show fragmented and deformed red cells. The mortality of HUS with good supportive care is less than 10%. Treatment includes fresh frozen plasma and plasma exchange, but antibiotics are not seen to positively affect prognosis.

32475

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	17
Responses Total:	17
Responses - % Correct:	0%

Back to Filters

Question 118 of 203

A 64-year-old man is referred urgently to the hospital by his general practitioner due to severe renal impairment picked up on routine blood tests. He had consulted his doctor because he had been feeling unwell for the previous few months, with malaise, anorexia, back pain and weight loss. He had only been passing very small amounts of urine over the past few weeks, despite an adequate fluid intake. He had no significant past medical history, was on no regular medication and did not smoke or drink alcohol. His renal function was last checked 8 years ago and was normal. On examination he was thin and dehydrated. He was apyrexial with a regular pulse of 98/min and blood pressure 126/72 mmHg. Cardiorespiratory examination was unremarkable. His abdomen was soft and non-tender with no palpable masses. Digital rectal examination revealed an enlarged smooth prostate. There were no focal neurological signs.

Investigation results are below:

Hb	8.2 g/dl
WCC	$4.2 \times 10^9/l$
Platelets	$156 \times 10^9/l$
MCV	98 fl
ESR	90mm/h
Urate	0.52 mmol/l
Na ⁺	136 mmol/l
K ⁺	5.2 mmol/l
Urea	32.0 mmol/l
Creatinine	820 μ mol/l
Ca ²⁺	3.2 mmol/l
GGT	48 IU/l
Total protein	102 g/l
Albumin	28 g/l
ALP	130 IU/l
Bilirubin	14 μ mol/l
AST	28 IU/l
Urinalysis	Protein ++

Which of the following investigations is most likely to point towards the underlying diagnosis?



- A

Autoimmune profile
- B

Chest radiograph
- C

Serum creatine kinase
- D

Serum electrophoresis
- E

Serum complement levels

32476

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 118 of 203

A 64-year-old man is referred urgently to the hospital by his general practitioner due to severe renal impairment picked up on routine blood tests. He had consulted his doctor because he had been feeling unwell for the previous few months, with malaise, anorexia, back pain and weight loss. He had only been passing very small amounts of urine over the past few weeks, despite an adequate fluid intake. He had no significant past medical history, was on no regular medication and did not smoke or drink alcohol. His renal function was last checked 8 years ago and was normal. On examination he was thin and dehydrated. He was afebrile with a regular pulse of 98/min and blood pressure 126/72 mmHg. Cardiorespiratory examination was unremarkable. His abdomen was soft and non-tender with no palpable masses. Digital rectal examination revealed an enlarged smooth prostate. There were no focal neurological signs.

Investigation results are below:

Hb	8.2 g/dl
WCC	4.2 × 10 ⁹ /l
Platelets	156 × 10 ⁹ /l
MCV	98 fl
ESR	90mm/h
Urate	0.52 mmol/l
Na ⁺	136 mmol/l
K ⁺	5.2 mmol/l
Urea	32.0 mmol/l
Creatinine	820 μmol/l
Ca ²⁺	3.2 mmol/l
GGT	48 IU/l
Total protein	102 g/l
Albumin	28 g/l
ALP	130 IU/l
Bilirubin	14 μmol/l
AST	28 IU/l
Urinalysis	Protein ++

Which of the following investigations is most likely to point towards the underlying diagnosis?



- A

Autoimmune profile
- B

Chest radiograph
- C

Serum creatine kinase
- D

Serum electrophoresis
- E

Serum complement levels

Explanation

The most likely diagnosis is myeloma, and serum protein electrophoresis is used to determine the type of each protein present and may indicate a characteristic curve (i.e. where the spike is observed). Urine protein electrophoresis may also be used to identify the presence of the Bence Jones protein in urine. Urinary tract obstruction due to prostatic hypertrophy or carcinoma should also be excluded. An ultrasound must be performed within 24 h for all patients presenting with new-onset renal failure. If it shows hydronephrosis and an obstructed urinary system, urgent decompression may be indicated. Based on the history and blood results in this case, it is most likely that the renal failure is chronic. Glomerulonephritis and autoimmune causes for the renal failure are less likely in this patient.

32476

Rate this question:

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	18
Responses Total:	18
Responses - % Correct:	0%

Back to Filters

Question 119 of 203

A 72-year-old man is admitted to hospital after being found lying on his bedroom floor by his son, who was unsure how long he had been there for. Apart from several falls recently, he was otherwise well, with no significant past medical history and he was on no regular medication. A BM blood glucose checked by the ambulance team was normal. On examination he was drowsy and clinically dehydrated with Kussmaul respiration. He was apyrexial, blood pressure 90/50 mmHg, pulse 120/min, oxygen saturations 85% on air. Apart from extensive bruising, there were no specific findings on examination.

Investigation results are below:

Hb	15 g/dl
WCC	12 × 10 ⁹ /l
Platelets	420 × 10 ⁹ /l
INR	1.1
Prothrombin time	13 s
Creatine kinase	30 000 IU/l
Ca ²⁺	2.6 mmol/l
Na ⁺	132 mmol/l
K ⁺	7.2 mmol/l
Urea	42 mmol/l
Creatinine	980 μmol/l
Phosphate	1.4 mmol/l
Urate	0.78 mmol/l
Bilirubin	16 μmol/l
AST	56 IU/l
ALP	80 IU/l
GGT	54 IU/l

Arterial blood gases:

pH	7.05
(on air) pa(O ₂)	7.2 kPa
Pa(CO ₂)	2.0 kPa
Bicarbonate	8 mmol/l
Electrocardiogram (ECG)	Tented T waves, normal QRS complex
Chest radiograph	Basal atelectasis, no focal lung lesion

Intravenous access was obtained and he was given high-flow oxygen and connect to continuous ECG monitoring.

What should be the next step in management?

- A

Intravenous normal saline
- B

Intravenous sodium bicarbonate 8.4%
- C

Oral calcium resonium
- D

Intravenous calcium gluconate
- E

Insulin/dextrose infusion

32477

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 119 of 203

Investigation results are below:

Arterial blood gases:



What should be the next step in management?

Explanation

Date this question:

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

[Blog](#) [About Pastest](#) [Contact Us](#) [Help](#)

Back to Filters

Question 120 of 203

A 78-year-old man is brought to the hospital by his daughter who is concerned about a significant change in his personality over the last 3 weeks. He has become increasingly disoriented, irritable and confused, as well as complaining of persistent nausea and headache. She noticed that he had lost some weight and had also had several falls. He had been depressed since the death of his wife 6 months ago and had recently been started on fluoxetine by his general practitioner. He smokes 5 cigarettes per day and drinks no alcohol. On examination he is afebrile with a pulse of 50/min regular and blood pressure 148/62 mmHg. Cardiorespiratory and abdominal examination is unremarkable. He is confused but fully alert with no focal neurological signs

Investigation results are below:

Hb	15 g/dl
WCC	$7.0 \times 10^9/l$
Platelets	$160 \times 10^9/l$
MCV	96 fl
ESR	10 mm/h
Glucose (random)	7.2 mmol/l
K ⁺	3.7 mmol/l
Na ⁺	118 mmol/l
Urea	2.3 mmol/l
Creatinine	120 μ mol/l
ALP	80 IU/l
Bilirubin	10 μ mol/l
Aspartate transaminase	32 IU/l
GGT	42 IU/l
Plasma osmolality	260 mosmol/kg (278-305)
Urine osmolality	430 mosmol/kg (350-1000)
ECG	Normal sinus rhythm
Urine culture	No growth

Which one of the following investigations would be the most useful next step?



- A

Thyroid function tests
- B

MRI of the pituitary fossa
- C

Short synacthen test
- D

Serum and urine electrophoresis
- E

Dexamethasone suppression test

32478

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Question 120 of 203

Investigation results are below.

Which one of the following investigations would be the most useful next step?

- ### Explanation

32478

[Next Question](#)

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

[Blog](#) [About Pastest](#) [Contact Us](#) [Help](#)

Question 121 of 203

Her urea and electrolytes are below:

Na ⁺	136 mmol/l
K ⁺	3.7 mmol/l
Urea	7.2 mmol/l
Creatinine	146 μmol/l

Which of the following is the next radiological investigation of choice?

- | | |
|---|--|
| A | CT abdomen |
| B | Doppler ultrasound scan of the transplant site |
| C | Intravenous urography |
| D | Renal angiography |
| E | Plain abdominal film |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 121 of 203

A 38-year-old woman who underwent a live related renal transplant for end-stage renal failure secondary to chronic pyelonephritis 12 weeks previously, attends the clinic for routine follow up. She is taking tacrolimus and mycophenolate mofetil (MMF).

Her urea and electrolytes are below:

Na ⁺	136 mmol/l
K ⁺	3.7 mmol/l
Urea	7.2 mmol/l
Creatinine	146 μmol/l

She had last been seen in clinic 2 weeks previously, when her urea was 4.2 mmol/l and creatinine 98 μmol/l. She is clinically well and asymptomatic. On examination she was afebrile and normotensive. Her transplant site was non-tender with no swelling and there were no other signs to be found.

Which of the following is the next radiological investigation of choice?

- A

CT abdomen
- B

Doppler ultrasound scan of the transplant site
- C

Intravenous urography
- D

Renal angiography
- E

Plain abdominal film

Explanation

Doppler ultrasound scan of the transplant site

Graft dysfunction is an early complication of renal transplantation and can present asymptotically with a rising serum creatinine. It is important to exclude sepsis in these patients, even if they are asymptomatic and afebrile, and cultures should always be checked. Tacrolimus levels should also be measured, as this drug can be directly nephrotoxic. MMF does not need levels monitoring. Doppler ultrasound of the transplant site should be arranged to look for any obstruction, arterial or venous occlusion and is the initial radiological investigation of choice. Renal angiography or biopsy may be needed in due course, but they are not first line investigations.

32479

Rate this question:      

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	21
Responses Total:	21
Responses - % Correct:	0%

Back to Filters

Question 122 of 203

A 45-year-old woman with type 2 diabetes presents for review. She currently takes metformin and gliclazide and has an HbA1c of 50.82 mmol/mol (6.8%). Blood pressure is 142/82 mmHg on no antihypertensive medication. Total cholesterol is 5.2 mmol/l, but high-density lipoprotein (HDL) level is only 0.8 mmol/l; serum creatinine is 92 μ mol/l. Urinary protein is in the microalbuminuric range, and was at the previous appointment 3 months earlier.

Which one of the following is most likely to be renoprotective in this case?

- | | |
|---|----------------------------|
| A | Atorvastatin 10 mg |
| B | Ramipril 10mg |
| C | Bendroflumethiazide 2.5 mg |
| D | Atenolol 50 mg |
| E | Amlodipine 5 mg |

32480

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Question 122 of 203

Which one of the following is most likely to be renoprotective in this case?

Explanation

End Session

Peer Responses %

0%

Back to Filters

Question 123 of 203

A 74-year-old woman presents with lethargy on a background of depression and Type 2 Diabetes mellitus. Her blood pressure is 130/70 mmHg and she is clinically euvolaemic. Routine biochemical investigations reveal Na^+ 120mmol/l, K^+ 3.8mmol/l, creatinine 76 mol/l and normal thyroid hormone levels.

Which biochemical test would be most useful in determining her state of water balance?

- A 24-hour urinary protein and creatinine
- B Erect / supine renin and aldosterone estimation
- C Fractional excretion of Na^+
- D Plasma ADH
- E Serum and urine osmolality

32482

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Question 123 of 203

Which biochemical test would be most useful in determining her state of water balance?

A	24-hour urinary protein and creatinine
B	Erect / supine renin and aldosterone estimation
C	Fractional excretion of Na^+
D	Plasma ADH
E	Serum and urine osmolality

Urine osmolality relative to serum osmolality gives an accurate assessment of the state of the ADH axis. In the absence of other causes (eg CCF), a high urine osmolality relative to serum osmolality in the setting of hyponatraemia is inappropriate. Other tests, such as plasma ADH and detailed urine electrolyte studies, can give similar information, but at much greater expense and difficulty. Sulphonylureas are recognised as a rare cause of SIADH, the condition is seen most commonly with chlorpropamide, whilst this has been replaced by more modern agents such as gliclazide, it is still occasionally seen.

Rate this question:      

End Session

Peer Responses %

Responses Correct:	0
Responses Incorrect:	23
Responses Total:	23
Responses - % Correct:	0%

Back to Filters

Question 124 of 203

A 70-year-old male presented with fever, productive cough and dyspnoea of 3 days’ duration. He has been gradually unwell for 4 months with lower back pain, lethargy, reduced appetite and has lost 2 kg of weight. He is hypertensive on atenolol 50 mg and he smokes 15 cigarettes daily. On examination he was febrile with a temperature of 38°C, pulse was 115 beats/min and blood pressure was 105/55 mmHg. He appeared pale with pitting oedema of both legs. He had no palpable lymph nodes.

Investigations are as shown:

Chest	X-ray Right lower lobe consolidation
Blood film	Prominent rouleaux formation
Hb	7.5 g/dl
WCC	19.4 × 10 ⁹ /l
Platelets	89 × 10 ⁹ /l
ESR	180 mm/h
Na ⁺	133 mmol/l
K ⁺	4.5 mmol/l
Urea	21 mmol/l
Creatinine	440 μmol/l
Total protein	90 g/l
Albumin	28 g/l
Corrected calcium	3.2 mmol/l
Urinalysis	protein +++++, blood +, MC&S (microscopy, culture and sensitivity test) – no growth

What underlying diagnosis best fits the clinical picture and investigation results?

- A

Bronchogenic carcinoma
- B

Hypercalcaemia
- C

Multiple myeloma
- D

Primary hyperparathyroidism
- E

Amylodosis

32484

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 124 of 203

Investigations are as shown:

A	Bronchogenic carcinoma
B	Hypercalcaemia
C	Multiple myeloma
D	Primary hyperparathyroidism
E	Amyloidosis

This man has an acute pneumonia, although the investigations we're presented with point towards underlying myeloma. The disease is characterised by abnormal proliferation of mature and immature plasma cells. The bone marrow is infiltrated by plasmacytomas (abnormal plasma cells) and there is overproduction of monoclonal immunoglobulin (Ig) – usually IgG in about 55%, IgA in about 20%; of these patients, 40% have Bence-Jones proteinuria. Light chain is found in 20% of patients, while IgD and IgE myeloma account for about 1% of cases.

Rate this question:

[Next Question](#)

Difficulty: Average

Difficulty: Average

Peer Responses %

Responses Correct:	0
Responses Incorrect:	24
Responses Total:	24
Responses - % Correct:	0%

Back to Filters

Question 125 of 203

A 35-year-old male was admitted unwell with severe dyspnoea. He reported two episodes of frank haemoptysis two weeks previously and another episode a few hours before admission. He had lost 2 kg in weight and he is constantly lethargic and nauseous. He is a non-smoker, drinks alcohol socially and he last travelled abroad 1 year ago to Thailand. On examination he is dyspnoeic at rest, and appeared pale with pitting oedema up to the knees. His BP is 125/72 mmHg.

Urgent investigations are as follows:

Hb	6.6 g/dl
WCC	12 × 10 ⁹ /l
Platelets	200 × 10 ⁹ /l
Na ⁺	132 mmol/l
K ⁺	6.0 mmol/l
Urea	22 mmol/l
Creatinine	480 μmol/l
Calcium	2.4 g/l
Albumin	26 g/l
Bilirubin	16 μmol/l
ALT	36 IU/l
AST	29 IU/l
Chest X-ray	Asymmetrical, bilateral air space shadowing
Thyroid function	Normal
Urinalysis	Protein +++, blood ++, culture no growth
Renal ultrasound scan	Normal

The patient was treated with intravenous fluid, cephalosporin and metronidazole. He was also transfused with 3 units of packed cells. Two days after admission, the patient’s condition continued to deteriorate with low urine output, increasing dyspnoea, and a further episode of frank haemoptysis.

Further investigations are shown below:

Chest X-ray	Increased bilateral air space shadowing
Hb	7.8 g/dl
WCC	12 × 10 ⁹ /l
Platelets	250 × 10 ⁹ /l
Na ⁺	129 mmol/l
K ⁺	6.5 mmol/l
Urea	28 mmol/l
Creatinine	650 μmol/l
Anti-GBM antibody	Positive
ANCA	Positive

What is the most likely underlying diagnosis?

- A

Infectious mononucleosis
- B

Primary syphilis
- C

Goodpasture’s syndrome
- D

Nephrotic syndrome
- E

Leptospirosis icterohaemorrhagia

32487

Submit

Previous Question

Skip Question

Calculator

✔

Normal Values

✔

Question 125 of 203

--	--

The patient was treated with intravenous fluid, cephalosporin and metronidazole. He was also transfused with 3 units of packed cells. Two days after admission, the patient's condition continued to deteriorate with low urine output, increasing dyspnoea, and a further episode of frank haemoptysis.

Further investigations are shown below:

What is the most likely underlying diagnosis?

A	Infectious mononucleosis
B	Primary syphilis
C	Goodpasture's syndrome
D	Nephrotic syndrome
E	Leptospirosis icterohaemorrhagica

Explanation:
Goodpasture's

Goodpasture's syndrome is characterised by circulating anti-GBM (glomerular basement membrane) antibodies in the blood with linear deposition in the glomerular basement membrane resulting in rapidly progressive glomerulonephritis (crescentic) and pulmonary haemorrhage.

The disease most often occurs in young men and is characterised

The disease most often occurs in young men and is characterised by severe haemoptysis, dyspnoea and rapidly progressive renal failure.

Anaemia, haematuria and proteinuria are common. Chest X-ray often shows bilateral asymmetrical shadowing of pulmonary haemorrhage. The combination of pulmonary haemorrhage and renal failure also occurs in vasculitis such as Wegener's granulomatosis, microscopic polyarteritis, and essential mixed cryoglobulinaemia. Vasculitis can usually be distinguished by the presence of anti-neutrophil cytoplasmic antibodies (ANCA) and findings from renal biopsy. Treatment is with pulse methylprednisolone, cyclophosphamide and plasma exchange. Given he has travelled to Thailand one year earlier, the foreign travel would be considered a "red herring".

Rate this question:

[Next Question](#)

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Responses Incorrect:	25
Responses Total:	25
Responses - % Correct:	0%

Back to Filters

Question 126 of 203

A 25-year old male is under investigation for long-standing hypokalaemia. He complains of persistent tiredness and muscle aches and finds it very difficult to do his job as a warehouse man. There is no other past medical history of note.

His most recent blood test results are shown below:

Sodium	140 mmol/l
Potassium	2.5 mmol/l
Urea	5.5 mmol/l
Creatinine	122 μ mol/l
Corrected calcium	1.90 mmol/l
Phosphate	0.7 mmol/l
Bicarbonate	14 mmol/l
Chloride	115 mmol/l (95-110 mmol/l)

What is the metabolic abnormality?

- A

Hypochloraemic metabolic alkalosis
- B

Hyperchloraemic metabolic acidosis
- C

Hyporeninaemic hypoaldosteronism
- D

High anion gap metabolic acidosis
- E

High anion gap metabolic alkalosis

32488

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 126 of 203

A 25-year old male is under investigation for long-standing hypokalaemia. He complains of persistent tiredness and muscle aches and finds it very difficult to do his job as a warehouse man. There is no other past medical history of note.

His most recent blood test results are shown below:

Sodium	140 mmol/l
Potassium	2.5 mmol/l
Urea	5.5 mmol/l
Creatinine	122 μmol/l
Corrected calcium	1.90 mmol/l
Phosphate	0.7 mmol/l
Bicarbonate	14 mmol/l
Chloride	115 mmol/l (95-110 mmol/l)

What is the metabolic abnormality?

- A

Hypochloraemic metabolic alkalosis
- B

Hyperchloraemic metabolic acidosis
- C

Hyporeninaemic hypoaldosteronism
- D

High anion gap metabolic acidosis
- E

High anion gap metabolic alkalosis

Explanation

The biochemical result shows a hyperchloraemic metabolic acidosis. Metabolic acidosis is due to the accumulation of any acid other than carbonic acid and a decrease in plasma bicarbonate. A series of disorders cause metabolic acidosis such as: ingestion of acid, generation of acid (lactic and ketoacidosis), impaired excretion by the kidneys and bicarbonate loss from the gastrointestinal tract. Calculation of anion gap is particularly useful to identify the cause of the acidosis. The main plasma anions are subtracted from the main cations to give the anion gap. The normal cations are - Na⁺, K⁺, Ca⁺⁺, Mg⁺⁺ and the anions are- Cl⁻ and HCO₃⁻. The normal anion gap is 10-18 mmol/l: (Na⁺ + K⁺) - (Cl⁻ + HCO₃⁻) In metabolic acidosis with normal anion gap, chloride is retained to balance the charge of the bicarbonate ions, which are lost. Causes of metabolic acidosis with normal anion gap are:

- loss of bicarbonate from the gastrointestinal tract: diarrhoea, ileostomy, ureterosigmoidostomy
- renal bicarbonate loss: renal tubular acidosis (proximal type 2), acetazolamide, tubular damage, hyperparathyroidism
- decreased hydrogen ion excretion from the kidneys: renal tubular acidosis (distal type1) and type 4 renal tubular acidosis.

32488

Rate this question:

⊖

★

★

★

★

★

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	26
Responses Total:	26
Responses - % Correct:	0%

Back to Filters

Question 127 of 203

A 65-year-old female presents for review. She is using acetazolamide and timolol for chronic glaucoma. Investigations reveal:

Sodium	145 mmol/l
Potassium	2.6 mmol/l
Urea	4.6 mmol/l
Creatinine	128 μmol/l
Corrected calcium	2.1 mmol/l
Chloride	118 mmol/l (95-110 mmol/l)
Bicarbonate	18 mmol/l (22-28 mmol/l)
Phosphate	0.65 mmol/l (0.8-1.5 mmol/l)

What is the cause of the metabolic abnormality?

- A

Bartter’s syndrome
- B

Nephrogenic diabetes insipidus
- C

Lactic acidosis
- D

Renal tubular acidosis type 1 (distal)
- E

Renal tubular acidosis type 2 (proximal)

32489

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 127 of 203

A 65-year-old female presents for review. She is using acetazolamide and timolol for chronic glaucoma. Investigations reveal:

Sodium	145 mmol/l
Potassium	2.6 mmol/l
Urea	4.6 mmol/l
Creatinine	128 μmol/l
Corrected calcium	2.1 mmol/l
Chloride	118 mmol/l (95–110 mmol/l)
Bicarbonate	18 mmol/l (22–28 mmol/l)
Phosphate	0.65 mmol/l (0.8–1.5 mmol/l)

What is the cause of the metabolic abnormality?



- A

Bartter’s syndrome
- B

Nephrogenic diabetes insipidus
- C

Lactic acidosis
- D

Renal tubular acidosis type 1 (distal)
- E

Renal tubular acidosis type 2 (proximal)

Explanation

Renal tubular acidosis is a disorder of the renal tubules’ ability to maintain acid-base balance leading to chronic hyperchloraemic metabolic acidosis.

The proximal tubule is the major site for reabsorption of filtered bicarbonate. In proximal RTA (type 2 RTA), bicarbonate reabsorption is defective, pRTA rarely occurs as an isolated defect of bicarbonate transport and is usually associated with multiple proximal tubule transport defects; therefore, urinary loss of glucose, amino acids, phosphate, uric acid, and other organic anions such as citrate can also occur (Fanconi syndrome). Plasma bicarbonate is <21 mmol/l, and urine pH may fall below 5.5 with acidification.

Features of RTA 2 are:

- acidosis
- hypokalaemia: due to inability of renal tubules to secrete hydrogen ions, potassium is preferentially excreted
- glycosuria
- aminoaciduria
- hyperchloraemia: chloride ions are mobilised to replace lost bicarbonate ions
- osteomalacia: excess hydrogen ions are buffered by calcuim, resulting in depletion of calcium from bone.

Causes and associations of RTA 2 are:

Fanconi’s syndrome, Wilson’s disease, cystinosis and hereditary fructose intolerance and acetazolamide, which leads to increased proximal tubule bicarbonate loss.

32489

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	27
Responses Total:	27
Responses - % Correct:	0%

Back to Filters

Question 128 of 203

A 38-year-old Nigerian woman presents with 4-week history of polyuria and polydipsia. She has a history of long-standing progressive dyspnoea, a dry cough and tiredness. A chest X-ray requested by the general practitioner was reported as abnormal with a bilateral hilar mass. On examination her BP is 142/72 mmHg, pulse is 75/min and regular. Her BMI is 25. There are scattered crackles on auscultation of the chest. You notice a rash on examination of her shins.

A water deprivation test was performed:

Water deprivation phase	
Plasma osmolality	298 mOsmol/kg (270-295 mOsmol/kg)
Urine osmolality	238 mOsmol/kg (350-1000 mOsmol/kg)
Desmopressin (DDAVP) phase	
Plasma osmolality	292 mOsmol/kg (270-295 mOsmol/kg)
Urine osmolality	680 mOsmol/kg (350-1000 mOsmol/kg)

What is the most likely underlying diagnosis?

- A

Diabetes mellitus
- B

Sarcoidosis
- C

Histiocytosis X
- D

Sjögren’s syndrome
- E

Wegener’s granulomatosis

32490

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 128 of 203

A 38-year-old Nigerian woman presents with 4-week history of polyuria and polydipsia. She has a history of long-standing progressive dyspnoea, a dry cough and tiredness. A chest X-ray requested by the general practitioner was reported as abnormal with a bilateral hilar mass. On examination her BP is 142/72 mmHg, pulse is 75/min and regular. Her BMI is 25. There are scattered crackles on auscultation of the chest. You notice a rash on examination of her shins.

A water deprivation test was performed:

Water deprivation phase	
Plasma osmolality	298 mOsmol/kg (270–295 mOsmol/kg)
Urine osmolality	238 mOsmol/kg (350–1000 mOsmol/kg)
Desmopressin (DDAVP) phase	
Plasma osmolality	292 mOsmol/kg (270–295 mOsmol/kg)
Urine osmolality	680 mOsmol/kg (350–1000 mOsmol/kg)

What is the most likely underlying diagnosis?



- A

Diabetes mellitus
- B

Sarcoidosis
- C

Histiocytosis X
- D

Sjögren’s syndrome
- E

Wegener’s granulomatosis

Explanation

The patient is deficient of antidiuretic hormone (ADH), therefore passing large volumes of diluted urine. This is due to the inability of the kidney to concentrate urine; as a result the urine osmolality is low. In the above case, during the water deprivation phase the urine osmolality is still low, but after DDAVP (desmopressin) the kidneys now concentrate the urine and the urine osmolality increases significantly from the initial value. Granulomatous diseases such as sarcoidosis, histiocytosis and tuberculosis could infiltrate the hypothalamus or pituitary gland causing diabetes insipidus. Intracranial sarcoidosis is the cause of her symptoms here. The rash on her shins is consistent with erythema nodosum.

32490

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	28
Responses Total:	28
Responses - % Correct:	0%

Back to Filters

Question 129 of 203

A 45-year-old mechanic was brought unconscious to the emergency department. He was found at the garage he owns by his wife having an epileptic fit, and had vomited twice. Apparently they have been having recent money worries. He is normally fit and well and has no past medical history, he is a non-smoker. He does not smell of alcohol.

Investigations reveal:

Computed tomography of the head	Normal
Chest X-ray	Normal
Sodium	132 mmol/l
Potassium	3.0 mmol/l
Urea	15 mmol/l
Creatinine	150 μmol/l
Bicarbonate	15 mmol/l
Calcium	2.0 mmol/l
Chloride	90 mmol/l
Glucose	5.0 mmol/l
Plasma osmolality	380 mOsmol/kg (275–295 mOsmol/kg)
Urine microscopy	Oxalate crystals



What is the most likely diagnosis?

- A

Lead poisoning
- B

Carbon monoxide intoxication
- C

Ethylene glycol poisoning
- D

Heroin overdose
- E

Rhabdomyolysis

32491

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 129 of 203

Investigations reveal:

Computed tomography of the head	Normal
Chest X-ray	Normal
Sodium	132 mmol/l
Potassium	3.0 mmol/l
Urea	15 mmol/l
Creatinine	150 μ mol/l
Bicarbonate	15 mmol/l
Calcium	2.0 mmol/l
Chloride	90 mmol/l
Glucose	5.0 mmol/l
Plasma osmolality	380 mOsmol/kg (275–295 mOsmol/kg)
Urine microscopy	Oxalate crystals

What is the most likely diagnosis?

A	Lead poisoning
B	Carbon monoxide intoxication
C	Ethylene glycol poisoning
D	Heroin overdose
E	Rhabdomyolysis

Ethylene glycol is a constituent of antifreeze used in cars. Metabolic products of the substance are directly responsible for the toxicity. Symptoms are: nausea, vomiting, convulsions, coma and respiratory distress with no alcohol odour on the breath. This differentiates ethylene glycol poisoning from other alcohols such as ethanol and methanol.

Severe metabolic acidosis, hypocalcaemia and renal failure are common. Renal failure is caused by oxalate crystalluria.

An osmolar gap exists when measured plasma osmolality is greater than calculated osmolality by >10 mOsmol/kg.

$$\text{Osmolality} = 2(\text{Na}^+ + \text{K}^+) + \text{urea} + \text{glucose}$$

Several unmeasured osmotically active substances in the plasma such as ethanol, methanol, ethylene glycol, mannitol and glycine are responsible for increased osmolar gap. Treatment is with inhibitors of alcohol dehydrogenase such as ethanol or fomepizole. Haemodialysis should also be employed to remove the substance and its metabolites, although this man's relatively late presentation is associated with a very poor prognosis.

Ethylene glycol is the only possible answer as it is the only option which leads to the formation of oxalate crystals. Carbon monoxide leads to unconsciousness and metabolic acidosis but carbon monoxide poisoning is usually intentional or as a result of combustion in an enclosed space, the mode of poisoning is therefore obvious.

Rate this question:

[Next Question](#)

Review Questions

Topic Questions

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	29
Responses Total:	29
Responses - % Correct:	0%

Back to Filters

Question 130 of 203

A 64-year-old man gives a two-week history of progressive lethargy and weakness. Eight weeks previously he was treated for hypertensive heart failure with 120 mg furosemide and commenced on 5 mg enalapril daily. He had not previously been treated with an ACE inhibitor. His haemoglobin at the time was 12.0, urea 14.2 mmol/l and creatinine 298 μmol/l. His blood pressure in clinic was 148/85 mmHg with no postural drop.

His blood results are shown below:

Hb	10.2 g/dl
MCV	89.2 fl
WCC	4.9 x 10 ⁹ /l
Platelets	175 x 10 ⁹ /l
Na ⁺	135 mmol/l
K ⁺	5.2 mmol/l
Urea	25.2 mmol/l
Creatinine	600 μmol/l

Which of the following is the most appropriate action?

- A

Reduce the dose of enalapril
- B

Reduce the dose of furosemide
- C

Stop the enalapril and furosemide
- D

Stop the enalapril
- E

Stop the furosemide

32492

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 130 of 203

His blood results are shown below:

Hb	10.2 g/dl
MCV	89.2 fl
WCC	$4.9 \times 10^9/l$
Platelets	$175 \times 10^9/l$
Na ⁺	135 mmol/l
K ⁺	5.2 mmol/l
Urea	25.2 mmol/l
Creatinine	600 mol/l

- | | |
|---|-----------------------------------|
| A | Reduce the dose of enalapril |
| B | Reduce the dose of furosemide |
| C | Stop the enalapril and furosemide |
| D | Stop the enalapril |
| E | Stop the furosemide |

Explanation

This man has pre-existing renal impairment that has worsened with the introduction of an ACE inhibitor, raising the possibility of renovascular disease. Given that his creatinine has doubled the most sensible option is to discontinue the ACE inhibitor. He does not appear volume depleted as there is no postural drop in blood pressure. Stopping his diuretics as well as the ACE inhibitor raises the chance of rebound pulmonary oedema; as such it is sensible to continue to current dose of furosemide for the moment.

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	30
Responses Total:	30
Responses - % Correct:	0%

Back to Filters

Question 131 of 203

A 28-year-old woman who is 14 weeks pregnant with her first pregnancy attends for a routine antenatal check. Her blood pressure is 142/92 mmHg. It is rechecked an hour later and found to be 138/90 mmHg. Her only past history of note includes two admissions to hospital with pyelonephritis as a child. The following investigations are performed:

Na+	137 mmol/l
K+	4.2 mmol/l
Urea	4.5 mmol/l
Creatinine	150 mol/l
Urinalysis	NAD
Ultrasound scan, kidneys:	
Right kidney BPD	10.8cm
Left kidney BPD	5.5cm
Irregular outline to left kidney	

What is the likely diagnosis?

- A

Chronic pyelonephritis
- B

Congenital renal atrophy
- C

Pre-eclampsia
- D

Glomerulonephritis
- E

Renal artery stenosis

32493

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Back to Filters

Question 131 of 203

A 28-year-old woman who is 14 weeks pregnant with her first pregnancy attends for a routine antenatal check. Her blood pressure is 142/92 mmHg. It is rechecked an hour later and found to be 138/90 mmHg. Her only past history of note includes two admissions to hospital with pyelonephritis as a child. The following investigations are performed:

Na+	137 mmol/l
K+	4.2 mmol/l
Urea	4.5 mmol/l
Creatinine	150 mol/l
Urinalysis	NAD
Ultrasound scan, kidneys:	
Right kidney BPD	10.8cm
Left kidney BPD	5.5cm
Irregular outline to left kidney	

What is the likely diagnosis?

- A

Chronic pyelonephritis
- B

Congenital renal atrophy
- C

Pre-eclampsia
- D

Glomerulonephritis
- E

Renal artery stenosis

Explanation

This woman has hypertension in pregnancy of which there are many causes. Her ultrasound scan showing a damaged left kidney is suggestive of chronic pyelonephritis. Chronic long-term renal damage related to pyelonephritis may predispose to the development of hypertension. Enalapril, bendrofluazide, hydralazine and atenolol are not recommended for the treatment of hypertension at this stage of pregnancy, but methyldopa would be an acceptable choice. Long-term management of the condition involves identification of any predisposing risk factors such as abnormal ureteric implantation or other structural problem that might be suitable for surgical repair.

32493

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	31
Responses Total:	31
Responses - % Correct:	0%

Question 132 of 203

—

—

32494

Submit

[Previous Question](#)
[Skip Question](#)

[Previous Question](#)
[Skip Question](#)

Back to Filters

Question 132 of 203

A 56-year-old man with past history of jejunocolic anastomosis for Crohn’s disease presents with symptoms of renal colic. He has suffered from Crohn’s for up to 25 years and is thought to have only the minimum amount of small bowel remaining to avoid significant malnutrition. On examination his BP is 105/70 mmHg, pulse is 80/min and regular and his BMI is 19. Investigations reveal presence of renal stones.

Which of the following is the most likely constituent of the stones?

A	Phosphate
B	Urate
C	Oxalate
D	Xanthine
E	Cysteine

Explanation

This man has had extensive small bowel resection for Crohn’s disease and this predisposes to the formation of calcium oxalate stones because of changes in the pattern of small bowel flora and electrolyte resorption. The best treatment to avoid further stone formation is to ensure adequate hydration, particularly in hot weather and dietary oxalate restriction. Sodium bicarbonate supplementation to cause alkalinisation of urine is the treatment for uric acid stones.

32494

Rate this question:      

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	32
Responses Total:	32
Responses - % Correct:	0%

Back to Filters

Question 133 of 203

A 17-year-old woman from West Africa presents to the clinic. She gives a history of several months of pitting oedema, which is now to mid-leg, and there is associated abdominal swelling. There is no shortness of breath, but she does complain of general malaise and intermittent fevers. There has been an episode of headache, upper abdominal pain and vomiting with associated diarrhoea 2 years ago but nil else of note. She was working on a fish farm during her time in Africa

Examination revealed evidence of ascites and marked lower limb oedema. Cardiovascular and respiratory examinations were unremarkable.

Investigations revealed:

Hb	10.1 g/dl
ALT	125 U/l
Creatinine	132 micromol/l
Other liver function tests	unremarkable
Albumin	26 g/l
Hepatic ultrasound	reveals evidence of hepatosplenomegaly
Urine dipstick	Blood +, protein +++

Which of the following investigations is the next most appropriate?

- A

Liver biopsy
- B

Urinary protein estimation
- C

Cardiac ultrasound scan
- D

Renal biopsy
- E

Hepatitis serology

32495

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 133 of 203

A 17-year-old woman from West Africa presents to the clinic. She gives a history of several months of pitting oedema, which is now to mid-leg, and there is associated abdominal swelling. There is no shortness of breath, but she does complain of general malaise and intermittent fevers. There has been an episode of headache, upper abdominal pain and vomiting with associated diarrhoea 2 years ago but nil else of note. She was working on a fish farm during her time in Africa

Examination revealed evidence of ascites and marked lower limb oedema. Cardiovascular and respiratory examinations were unremarkable.

Investigations revealed:

Hb	10.1 g/dl
ALT	125 U/l
Creatinine	132 micromol/l
Other liver function tests	unremarkable
Albumin	26 g/l
Hepatic ultrasound	reveals evidence of hepatosplenomegaly
Urine dipstick	Blood +, protein +++

Which of the following investigations is the next most appropriate?

- A

Liver biopsy
- B

Urinary protein estimation
- C

Cardiac ultrasound scan
- D

Renal biopsy
- E

Hepatitis serology

Explanation

We are aware that this patient has marked ascites and lower limb oedema. There are no signs of cardiac dysfunction on examination. Renal protein loss is one likely explanation for the oedema and hypoalbuminaemia, and it can be easily excluded. Significant exposure to mosquitoes is likely as this patient worked on a fish farm. She has a history suggestive of chronic malarial infection. It is well known that plasmodium malariae may be associated with membranous glomerulonephritis, the most likely cause of her significant proteinuria. Chloroquine would be appropriate treatment in this case.

32495

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	33
Responses Total:	33
Responses - % Correct:	0%

Back to Filters

Question 134 of 203

A 75-year-old male was reviewed in the emergency department with a 2-month history of dyspnoea, paraesthesia of both hands and feet and tiredness.

He is on atenolol 100 mg for long-standing hypertension and smokes 10 cigarettes daily. On examination he has oedema of his legs, jugular venous pressure is elevated and the cardiac apex beat is displaced downward to the 6th intercostal margin on the anterior axillary line. His BP is 132/82 mmHg, pulse is 85/min and regular. Abdominal examination reveals hepatosplenomegaly and bilateral painless ballotable loin masses.

Investigations reveal:

24 h urine collection	Protein 4.2 g
Hb	7.5 g/dl
WCC	10 × 10 ⁹ /l
Platelets	200 × 10 ⁹ /l
Na ⁺	134 mmol/l
K ⁺	5.3 mmol/l
Urea	20 mmol/l
Creatinine	390 μmol/l
Albumin	25 g/l
Total protein	93 g/l



What is the most likely cause of his renal dysfunction?

- A

Membranous nephropathy
- B

Cast nephropathy
- C

Congestive cardiac failure
- D

Amyloidosis
- E

Goodpasture’s syndrome

32496

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 134 of 203

He is on atenolol 100 mg for long-standing hypertension and smokes 10 cigarettes daily. On examination he has oedema of his legs, jugular venous pressure is elevated and the cardiac apex beat is displaced downward to the 6th intercostal margin on the anterior axillary line. His BP is 132/82 mmHg, pulse is 85/min and regular. Abdominal examination reveals hepatosplenomegaly and bilateral painless ballotable loin masses.

24 h urine collection	Protein 4.2 g
Hb	7.5 g/dl
WCC	$10 \times 10^9/\text{l}$
Platelets	$200 \times 10^9/\text{l}$
Na ⁺	134 mmol/l
K ⁺	5.3 mmol/l
Urea	20 mmol/l
Creatinine	390 $\mu\text{mol/l}$
Albumin	25 g/l
Total protein	93 g/l

- | | |
|---|----------------------------|
| A | Membranous nephropathy |
| B | Cast nephropathy |
| C | Congestive cardiac failure |
| D | Amyloidosis |
| E | Goodpasture's syndrome |

Amyloidosis is the accumulation in various tissues of deposited insoluble protein that impairs normal function. The fact that he has hepatosplenomegaly and bilateral ballotable masses (kidney enlargement) fits best with amyloid infiltration. Two major types of protein are deposited in tissues: protein light chain (AL) in primary amyloidosis and myeloma-associated amyloidosis, and non-immunoglobulin protein (AA) in secondary amyloidosis usually associated with chronic diseases such as: TB, bronchiectasis, osteomyelitis, rheumatoid arthritis, familial Mediterranean fever and Hodgkin's disease.

Diagnosis is based on clinical presentation, but the best screening tests are rectal and renal biopsy, subcutaneous fat aspiration and tests at other sites such as, skin, nerve and gingival biopsy. Tissue sampled is stained with Congo red and examined by polarised light to show 'apple green-birefringence' of amyloid protein.

Rate this question:

[Next Question](#)

[Previous Question](#)

[Tag Question](#)

[Feedback](#)

[End Session](#)

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	34
Responses Total:	34
Responses - % Correct:	0%

Back to Filters

Question 135 of 203

A 71-year-old man presents to the GP surgery for review. He has a past history of symptoms of prostatic hypertrophy with occasional wetting, but particularly difficulty starting urination, poor stream, and multiple visits to the toilet in the night. On examination his blood pressure is 165/90 mmHg; this is the third elevated measure in the past 3 visits to the surgery over a period of 6 months.

Investigations reveal;

Hb	13.9 g/dl
WCC	5.0 x10 ⁹ /l
PLT	305 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	5.0 mmol/l
Creatinine	150 μmol/l
PSA	3.1 μg/l

Which of the following would be the most appropriate blood pressure lowering agent given his prostatic problem?

- A

Doxazosin
- B

Bendroflumethiazide
- C

Furosemide
- D

Amlodipine
- E

Lisinopril

32497

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 135 of 203

Investigations reveal;

Hb	13.9 g/dl
WCC	5.0 x10 ⁹ /l
PLT	305 x10 ⁹ /l
Na ⁺	139 mmol/l
K ⁺	5.0 mmol/l
Creatinine	150 μ mol/l
PSA	3.1 μ g/l



A	Doxazosin
B	Bendroflumethiazide
C	Furosemide
D	Amlodipine
E	Lisinopril

Alpha blockade is a well recognised treatment for benign prostatic hypertrophy. It promotes smooth muscle relaxation improving urinary flow and reducing the problem of residual urine in the bladder post voiding. Many physicians also combine therapy with a 5-alpha reductase inhibitor, which acts over the longer term to reduce prostatic hypertrophy. By virtue of their mode of action, alpha-blockers may lead to hypotension and tachycardia. Psychomotor effects seen include insomnia, depression and anorexia. Decreased libido, ejaculation disorders and decreased volume of ejaculate were all said to occur with a frequency of more than 1% in clinical trials.

Rate this question:

[Next Question](#)

[Previous Question](#)

[Tag Question](#)

[Feedback](#)

[End Session](#)

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	35
Responses Total:	35
Responses - % Correct:	0%

Back to Filters

Question 136 of 203

A 49-year-old woman presents to the Emergency room complaining of haemoptysis. She has attended the GP on 3 previous occasions over the past 6 months complaining of recurrent nose bleeds and sinusitis. Examination of the chest reveals scattered crackles and signs of consolidation, particularly on the left hand side. On further questioning she admits to smoking 30 cigarettes per day.

Investigations;

Hb	11.0 g/dl
WCC	8.1 x 10 ⁹ /l
PLT	430 x 10 ⁹ /l
Na ⁺	140 mmol/l
K ⁺	5.7 mmol/l
Creatinine	190 μmol/l
c-ANCA	positive
CXR	Consolidation with evidence of granuloma formation

Which of the following is the most likely diagnosis?



- A

Tuberculosis
- B

Churg-Strauss vasculitis
- C

Wegener’s granulomatosis
- D

Sarcoidosis
- E

Anti-GBM disease

32498

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 136 of 203

Investigations;

Hb	11.0 g/dl
WCC	8.1 x 10 ⁹ /l
PLT	430 x 10 ⁹ /l
Na ⁺	140 mmol/l
K ⁺	5.7 mmol/l
Creatinine	190 μmol/l
c-ANCA	positive
CXR	Consolidation with evidence of granuloma formation

- | | |
|---|--------------------------|
| A | Tuberculosis |
| B | Churg-Strauss vasculitis |
| C | Wegener's granulomatosis |
| D | Sarcoidosis |
| E | Anti-GBM disease |

The picture seen here fits well with a diagnosis of Wegener's, with positive c-ANCA, anaemia with leukocytosis, thrombocytosis, granulomas on CXR and evidence of renal involvement. Diagnosis is confirmed by taking a histology sample from affected tissue. Easiest to obtain is usually a biopsy of nasal/ upper respiratory tract mucosa, but if this is impossible, lung or renal biopsy is the alternative. Most physicians begin treatment with a course of 3 days high dose IV methylprednisolone and cyclophosphamide with the aim of inducing remission from Wegener's. This is followed by conversion to lower doses of orals. IV immunoglobulin may be used in patients who are severely ill. More recently, IV rituximab (an anti CD20 antibody) has shown promising results in inducing a clinical response in patients resistant to medical treatment.

Rate this question:

[Next Question](#)

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	36
Responses Total:	36
Responses - % Correct:	0%

Question 137 of 203

Investigations;

Hb	13.2 g/dl
WCC	$7.2 \times 10^9/l$
PLT	$198 \times 10^9/l$
Na ⁺	137 mmol/l
K ⁺	3.6 mmol/l
Creatinine	135 micromol/l
Renal angiogram	"string of beads" appearance of left renal artery

—

- | | |
|---|--------------------------------------|
| A | Angioplasty |
| B | Add Atenolol 50mg |
| C | Add Doxazosin 4mg |
| D | Increase Ramipril to 10mg |
| E | Surgical renal artery reimplantation |

Submit

[Previous Question](#)
[Skip Question](#)

Skip Question

Question 137 of 203

Investigations;

Hb	13.2 g/dl
WCC	$7.2 \times 10^9/l$
PLT	$198 \times 10^9/l$
Na ⁺	137 mmol/l
K ⁺	3.6 mmol/l
Creatinine	135 micromol/l
Renal angiogram	"string of beads" appearance of left renal artery

A	Angioplasty
B	Add Atenolol 50mg
C	Add Doxazosin 4mg
D	Increase Ramipril to 10mg
E	Surgical renal artery reimplantation

Recent meta-analyses have suggested that atherosclerotic renal artery stenosis, medical management is as effective as angioplasty. This is in contrast to fibromuscular hyperplasia, which has the characteristic angiography appearance seen here. In this case angioplasty has superior outcomes with respect to control of blood pressure and long term sequelae. As such angioplasty is recommended as the most appropriate intervention. Surgical arterial re-implantation is not recommended, particularly in the light of modern angioplasty and stenting techniques.

Rate this question:

Next Question

[Previous Question](#)

[Tag Question](#)

[Feedback](#)

[End Session](#)

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	37
Responses Total:	37
Responses - % Correct:	0%

Question 138 of 203

Investigations;

Hb	13.1 g/dl
WCC	6.8 x10 ⁹ /l
PLT	212 x10 ⁹ /l
Na ⁺	138 mmol/l
K ⁺	3.1 mmol/l
Creatinine	110 micromol/l
Ultrasound renal tract	differential kidney size with small left kidney

11

- | | |
|---|----------------------------|
| A | Primary hyperaldosteronism |
| B | Renal artery stenosis |
| C | Essential hypertension |
| D | Reflux nephropathy |
| E | Phaeochromocytoma |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Question 138 of 203

Investigations;

Hb	13.1 g/dl
WCC	6.8 x10 ⁹ /l
PLT	212 x10 ⁹ /l
Na ⁺	138 mmol/l
K ⁺	3.1 mmol/l
Creatinine	110 micromol/l
Ultrasound renal tract	differential kidney size with small left kidney

Which of the following is the most likely diagnosis?

A	Primary hyperaldosteronism
B	Renal artery stenosis
C	Essential hypertension
D	Reflux nephropathy
E	Phaeochromocytoma

Explanation

There are multiple clues to the fact that this patient is a significant vasculopathy, with the previous inferior MI, poor peripheral circulation, and multiple bruits. This coupled with a differential kidney size on ultrasound is suggestive of renal artery stenosis, resultant secondary hyperaldosteronism leading to the hypokalaemia seen here. Angiography is the definitive investigation to confirm the diagnosis, although in practice, recent large studies have suggested that medical management of hypertension is as effective as vascular intervention.

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	38
Responses Total:	38
Responses - % Correct:	0%

Question 139 of 203

Investigations:

Hb	13.1 g/dl
WCC	$8.9 \times 10^9/l$
PLT	$183 \times 10^9/l$
Na ⁺	138 mmol/l
K ⁺	3.1 mmol/l
Bicarbonate	33 mmol/l
Creatinine	100 micromol/l

11

- | | |
|---|---------------------|
| A | Bartter's syndrome |
| B | Gitelman's syndrome |
| C | Conn's syndrome |
| D | Cushing's syndrome |
| E | Vasovagal syncope |

Submit

Skip Question

Back to Filters

Question 139 of 203

A 22-year-old woman is admitted with a third episode of syncope over the past 6 months. She says that she tries her best in hot weather to maintain her fluid intake, but often feels light headed. She does not take any regular medication apart from the combined oral contraceptive pill. On examination her BP is 100/60 mmHg, pulse is 75/min and regular. BMI is 23. There are no abnormal findings.

Investigations:

Hb	13.1 g/dl
WCC	8.9 x10 ⁹ /l
PLT	183 x10 ⁹ /l
Na ⁺	138 mmol/l
K ⁺	3.1 mmol/l
Bicarbonate	33 mmol/l
Creatinine	100 micromol/l

Which of the following is the most likely diagnosis?



- A

Bartter’s syndrome
- B

Gitelman’s syndrome
- C

Conn’s syndrome
- D

Cushing’s syndrome
- E

Vasovagal syncope

Explanation

Both Bartter’s syndrome and Gitelman’s present with hypokalaemic metabolic alkalosis in the absence of hypertension. This helps to differentiate them Conn’s syndrome which is of course associated with hypertension. Bartter’s normally presents in childhood with constipation, growth failure, muscle cramps and weakness. Gitelman’s on the other hand may be associated with milder disease and presents later. Gitelman’s syndrome is associated with reduced calcium excretion whereas Bartter’s is not, and this helps to differentiate between the two conditions. Renin and aldosterone are both elevated. Cushing’s syndrome is of course associated with obesity.

36476

Rate this question:

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	39
Responses Total:	39
Responses - % Correct:	0%

Back to Filters

Question 140 of 203

A 69-year-old woman is found on the floor having slipped in the bathroom at home. She has been lying on the floor for up to 8hrs before being found by her carer. There is a past history of ischaemic heart disease for which she is treated with Ramipril, Bisoprolol, Aspirin and Atorvastatin. Her BP is 122/72 mmHg, pulse is 80/min, sinus rhythm. She has signs of left lower lobe consolidation and there is extensive bruising over the left hip and pelvis. Urine is 3+ for blood.

You are concerned about the possibility of rhabdomyolysis.

Which of the following is the most appropriate investigation to confirm the diagnosis?

- A

Albumin
- B

Creatinine
- C

Creatine Kinase
- D

LDH
- E

Urine microscopy

37190

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 140 of 203

You are concerned about the possibility of rhabdomyolysis.



A	Albumin
B	Creatinine
C	Creatine Kinase
D	LDH
E	Urine microscopy

The options for diagnosing rhabdomyolysis include serum CK and urine microscopy. The presence of blood on urine dipstick and an absence of cells on microscopy would suggest that it is myoglobinuria rather than haematuria which is responsible for the positive test. However CK is the preferred answer, as the level is predictive of possible renal impairment. An LDH rise is seen in muscle necrosis but it occurs later in the process. Whilst elevated creatinine may be seen, it does not reveal the underlying cause of renal impairment.

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Session

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	40
Responses Total:	40
Responses - % Correct:	0%

Back to Filters

Question 141 of 203

A 62-year-old man presents to the Emergency Department with severe left loin pain and fevers which have increased over the past 48hrs. He has a history of chronic tophaceous gout for which he takes Allopurinol and Benzbromarone and hypertension for which he takes Amlodipine 5mg once daily. Examination reveals a BP of 100/60 mmHg, pulse is 95/min and regular. He is pyrexial 38.5°C. Abdominal examination reveals severe left loin pain.

Investigations;

Hb	13.1 g/dl	<div></div>
WCC	16.1 x10 ⁹ /l	
PLT	201 x10 ⁹ /l	
Na ⁺	138 mmol/l	
K ⁺	4.4 mmol/l	
Creatinine	187 micromol/l (up from 121 micromol/l at last clinic appointment)	
CRP	271 mg/l	
Urine	red cells++, white cells++, nitrites+	

Which of the following is the most appropriate next investigation?

- A

Cystoscopy
- B

IVP
- C

KUB
- D

Non-contrast enhanced CT
- E

Ultrasound abdomen

39028

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 141 of 203

A 62-year-old man presents to the Emergency Department with severe left loin pain and fevers which have increased over the past 48hrs. He has a history of chronic tophaceous gout for which he takes Allopurinol and Benzbromarone and hypertension for which he takes Amlodipine 5mg once daily. Examination reveals a BP of 100/60 mmHg, pulse is 95/min and regular. He is pyrexial 38.5°C. Abdominal examination reveals severe left loin pain.

Investigations;

Hb	13.1 g/dl	<div>≡</div>
WCC	16.1 x10 ⁹ /l	
PLT	201 x10 ⁹ /l	
Na ⁺	138 mmol/l	
K ⁺	4.4 mmol/l	
Creatinine	187 micromol/l (up from 121 micromol/l at last clinic appointment)	
CRP	271 mg/l	
Urine	red cells++, white cells++, nitrites+	

Which of the following is the most appropriate next investigation?

- A

Cystoscopy
- B

IVP
- C

KUB
- D

Non-contrast enhanced CT
- E

Ultrasound abdomen

Explanation

The potential diagnosis here is obstructed pyelonephritis due to a urate-containing stone within the left ureter. In this situation, with a significant rise in creatinine, contrast enhanced imaging carries risk. Out of the other remaining options, non-contrast enhanced CT is thought to be more sensitive in detecting obstruction versus ultrasound scanning. In the presence of significant infection, a conservative approach is not appropriate. IV antibiotics plus cystoscopy and potential JJ stent insertion is therefore the management of choice.

39028

Rate this question:

⊖

★

★

★

★

★

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	41
Responses Total:	41
Responses - % Correct:	0%

Back to Filters

Question 142 of 203

A 58-year-old man comes to the Oncology Clinic some 6 weeks after right nephrectomy for metastatic renal carcinoma that had resulted in a pathological fracture of the right humerus. He has recovered well from the surgery and is keen to discuss next steps. On examination his BP is 138/82 mmHg; pulse is 80/min and regular. His BMI is 23 and he has a nephrectomy scar on examination of the abdomen.

Investigations;

Hb	10.9 g/dl
WCC	9.1 x10 ⁹ /l
PLT	181 x10 ⁹ /l
ESR	64 mm/1 st hour
Na ⁺	137 mmol/l
K ⁺	4.5 mmol/l
Creatinine	162 micromol/l
Glucose	5.6 mmol/l
Bone scan	widespread hotspots consistent with metastatic disease

Treatment with recombinant IL-2 is planned.

Which of the following other therapies can be added?

- A

Bevacizumab
- B

Everolimus
- C

Pazopanib
- D

Sorafenib
- E

Sunitinib

40122

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 142 of 203

A 58-year-old man comes to the Oncology Clinic some 6 weeks after right nephrectomy for metastatic renal carcinoma that had resulted in a pathological fracture of the right humerus. He has recovered well from the surgery and is keen to discuss next steps. On examination his BP is 138/82 mmHg; pulse is 80/min and regular. His BMI is 23 and he has a nephrectomy scar on examination of the abdomen.

Investigations;

Hb	10.9 g/dl
WCC	9.1 x10 ⁹ /l
PLT	181 x10 ⁹ /l
ESR	64 mm/1 st hour
Na ⁺	137 mmol/l
K ⁺	4.5 mmol/l
Creatinine	162 micromol/l
Glucose	5.6 mmol/l
Bone scan	widespread hotspots consistent with metastatic disease

Treatment with recombinant IL-2 is planned.

Which of the following other therapies can be added?

- A

Bevacizumab
- B

Everolimus
- C

Pazopanib
- D

Sorafenib
- E

Sunitinib

Explanation

Sunitinib is a tyrosine kinase inhibitor recommended for patients with metastatic renal carcinoma with a good functional status who are suitable for cytokine therapy. In those who are not suitable for cytokine therapy, Pazopanib is an alternative option. The other choices are no longer recommended by NICE as therapies for metastatic renal cancer. Prognosis is closely related to the presence of metastases; in patients with distant metastases, 5-year survival is only 10%, versus 90% for those with early kidney cancer.
<http://www.medicines.org.uk/emc/medicine/18531>

40122

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	42
Responses Total:	42
Responses - % Correct:	0%

Back to Filters

Question 143 of 203

A 17-year-old woman who works at a local day nursery is admitted to the Emergency Department with a severe headache and weakness affecting the left hand side of her face. She has been unwell for the past few days with headaches, anorexia and diarrhoea. Apparently she has eaten the same food as a number of children at the nursery who have fallen ill with diarrhoea, two of whom are in the local hospital with renal failure. On examination her BP is 155/90 mmHg; pulse is 85/min and regular. There is evidence of a purpuric rash and petechial haemorrhages. Abdomen is soft and generally tender, and she has active bowel sounds. You confirm left upper motor neurone 7th nerve palsy.

Investigations;

Hb	9.9 g/dl (shistocytes on blood film)
WCC	12.1 x10 ⁹ /l
PLT	32 x10 ⁹ /l
Na ⁺	137 mmol/l
K ⁺	5.2 mmol/l
Creatinine	141 micromol/l
LDH	950 U/l
Urine	blood ++ protein ++



Which of the following is the most appropriate intervention?

- A

Ciprofloxacin
- B

IV immunoglobulin
- C

IV methylprednisolone
- D

Platelet transfusion
- E

Plasma exchange

40123

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 143 of 203

A 17-year-old woman who works at a local day nursery is admitted to the Emergency Department with a severe headache and weakness affecting the left hand side of her face. She has been unwell for the past few days with headaches, anorexia and diarrhoea. Apparently she has eaten the same food as a number of children at the nursery who have fallen ill with diarrhoea, two of whom are in the local hospital with renal failure. On examination her BP is 155/90 mmHg; pulse is 85/min and regular. There is evidence of a purpuric rash and petechial haemorrhages. Abdomen is soft and generally tender, and she has active bowel sounds. You confirm left upper motor neurone 7th nerve palsy.

Investigations;

Hb	9.9 g/dl (shistocytes on blood film)
WCC	12.1 x10 ⁹ /l
PLT	32 x10 ⁹ /l
Na ⁺	137 mmol/l
K ⁺	5.2 mmol/l
Creatinine	141 micromol/l
LDH	950 U/l
Urine	blood ++ protein ++



Which of the following is the most appropriate intervention?

- A

Ciprofloxacin
- B

IV immunoglobulin
- C

IV methylprednisolone
- D

Platelet transfusion
- E

Plasma exchange

Explanation

The history and blood picture seen here is consistent with thrombotic thrombocytopaenic purpura, (TTP). Plasma exchange is the initial therapy of choice, which has been associated with a reduction in mortality rate to less than 25% since its introduction. There is no evidence that corticosteroids or antibiotics have any benefit in the management of TTP. FFP is a potential interim solution until plasma exchange is available, but there is no role for IV immunoglobulin therapy. Platelet transfusion is contraindicated unless there is life-threatening haemorrhage.

40123

Rate this question:

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	43
Responses Total:	43
Responses - % Correct:	0%

Back to Filters

Question 144 of 203

A 21-year-old woman, who works as a receptionist in a doctor’s surgery, is admitted to the Emergency Department after fainting at work. She puts it down to it being a particularly hot day, but on further questioning admits to extreme fatigue and muscle cramps when she takes on relatively minor exercise, for as long as she can remember. Her mother, who has come to the Emergency Department to collect her, confirms she was always missing school sports because of lethargy and muscle pains. Examination reveals a BP of 100/70 mmHg; pulse is 75/min and regular. She is of normal height and her BMI is 22.

Investigations;

Hb	12.5 g/dl
WCC	6.1 x10 ⁹ /l
PLT	189 x10 ⁹ /l
Na ⁺	138 mmol/l
K ⁺	3.0 mmol/l
Bicarbonate	31 mmol/l
Creatinine	80 micromol/l

Which of the following is the most likely diagnosis?

- A

Bartter’s syndrome
- B

Conn’s syndrome
- C

Cushing’s syndrome
- D

Diuretic abuse
- E

Gitelman’s syndrome

40251

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 144 of 203

A 21-year-old woman, who works as a receptionist in a doctor’s surgery, is admitted to the Emergency Department after fainting at work. She puts it down to it being a particularly hot day, but on further questioning admits to extreme fatigue and muscle cramps when she takes on relatively minor exercise, for as long as she can remember. Her mother, who has come to the Emergency Department to collect her, confirms she was always missing school sports because of lethargy and muscle pains. Examination reveals a BP of 100/70 mmHg; pulse is 75/min and regular. She is of normal height and her BMI is 22.

Investigations;

Hb	12.5 g/dl
WCC	6.1 x10 ⁹ /l
PLT	189 x10 ⁹ /l
Na ⁺	138 mmol/l
K ⁺	3.0 mmol/l
Bicarbonate	31 mmol/l
Creatinine	80 micromol/l

Which of the following is the most likely diagnosis?

- A

Bartter’s syndrome
- B

Conn’s syndrome
- C

Cushing’s syndrome
- D

Diuretic abuse
- E

Gitelman’s syndrome

Explanation

This patient in all likelihood has suffered from hypokalaemia for many years, leading in conjunction with hypomagnesaemia to severe fatigue and muscle cramps. Given the long history, although she works in a doctor’s surgery, access to or abuse of diuretics seems unlikely. Both Conn’s and Cushing’s are associated with hypertension, Cushing’s also with weight gain, and Bartter’s usually presents in childhood because of failure to thrive. Workup to confirm the diagnosis includes a diuretic screen and urinary calcium excretion, which if elevated to levels >6.9mmol over a 24hr period suggests Bartter’s syndrome.

40251

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	44
Responses Total:	44
Responses - % Correct:	0%

Back to Filters

Question 145 of 203

A 38-year-old man with a 12-year history of HIV, with a good response to therapy, was admitted with a 12-day history of weakness, nausea, dysuria, myalgia and decreased urine output. His last CD4 lymphocyte count was 368 x10⁶/l (normal range 500-900). The current medications included Didanosine, Stavudine, Atorvastatin, Trimethoprim-sulfamethoxazole, and he had been commenced on Indinavir. On examination he was afebrile, pulse 100/min and regular, BP 176/99 mmHg, the JVP was not elevated; the chest and abdominal examinations were normal.

Investigations:

Hb	13.5 g/dl
WCC	6.7 x10 ⁹ /l
Platelets	200 x10 ⁹ /l
Na ⁺	135 mmol/l
K ⁺	5.2 mmol/l
Cl ⁻	102 mmol/l
Bicarbonate	16 mmol/l
Urea	29.2 mmol/l
Creatinine	765 micromol/l
Bone profile	normal
Creatine kinase	100 U/l
ANCA	negative
Anti-GBM	negative
Hep B/ Hep C	negative
ASOT	negative



Cultures of blood/urine/sputum were negative.

Which one of the following medications is the most likely cause of his acute renal failure?

- A

Didanosine
- B

Stavudine
- C

Indinavir
- D

Atorvastatin
- E

Trimethoprim-sulfamethoxazole

70057

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 145 of 203

A 38-year-old man with a 12-year history of HIV, with a good response to therapy, was admitted with a 12-day history of weakness, nausea, dysuria, myalgia and decreased urine output. His last CD4 lymphocyte count was 368 x10⁶/l (normal range 500-900). The current medications included Didanosine, Stavudine, Atorvastatin, Trimethoprim-sulfamethoxazole, and he had been commenced on Indinavir. On examination he was afebrile, pulse 100/min and regular, BP 176/99 mmHg, the JVP was not elevated; the chest and abdominal examinations were normal.

Investigations:

Hb	13.5 g/dl
WCC	6.7 x10 ⁹ /l
Platelets	200 x10 ⁹ /l
Na ⁺	135 mmol/l
K ⁺	5.2 mmol/l
Cl ⁻	102 mmol/l
Bicarbonate	16 mmol/l
Urea	29.2 mmol/l
Creatinine	765 micromol/l
Bone profile	normal
Creatine kinase	100 U/l
ANCA	negative
Anti-GBM	negative
Hep B/ Hep C	negative
ASOT	negative



Cultures of blood/urine/sputum were negative.

Which one of the following medications is the most likely cause of his acute renal failure?

- A

Didanosine
- B

Stavudine
- C

Indinavir
- D

Atorvastatin
- E

Trimethoprim-sulfamethoxazole

Explanation



C

Indinavir

Drugs used in patients with HIV may cause acute kidney injury (AKI). This occurs most commonly with protease inhibitors, in particular Indinavir, by causing intratubular crystal obstruction. It has also been reported with Ritonavir.

A

Didanosine

Didanosine is a nucleoside analogue (NRTI), the most common side effect being peripheral neuropathy. There is no lactate provided to suggest lactic acidosis as the cause of acidaemia.

B

Stavudine

Stavudine is a nucleoside analogue (NRTI), the most common side effect being peripheral neuropathy. There is no lactate provided to suggest lactic acidosis as the cause of acidaemia.

D

Atorvastatin

Atorvastatin may cause myositis but in this patient the serum creatine kinase is normal.

E

Trimethoprim-sulfamethoxazole

Trimethoprim-sulfamethoxazole may cause interstitial nephritis presenting with AKI but the patient has been on this drug for a while with no problems.

70057

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	45
Responses Total:	45
Responses - % Correct:	0%

Back to Filters

Question 146 of 203

A 38-year-old man with a 12-year history of HIV, with a good response to therapy, was admitted with a 12-day history of weakness, nausea, dysuria, myalgia and decreased urine output. His last CD4 lymphocyte count was 368 x10⁶/l (normal range 500-900). The current medications included Didanosine, Stavudine, Atorvastatin, Trimethoprim-sulfamethoxazole, and he had been commenced on Indinavir. On examination he was afebrile, pulse 100/min and regular, BP 176/99 mmHg, the JVP was not elevated; the chest and abdominal examinations were normal.

Investigations:

Hb	13.5 g/dl
WCC	6.7 x10 ⁹ /l
Platelets	200 x10 ⁹ /l
Na ⁺	135 mmol/l
K ⁺	5.2 mmol/l
Cl ⁻	102 mmol/l
Bicarbonate	16 mmol/l
Urea	29.2 mmol/l
Creatinine	765 micromol/l
Bone profile	normal
Creatine kinase	100 U/l
ANCA	negative
Anti-GBM	negative
Hep B/ Hep C	negative
ASOT	negative

Cultures of blood/urine/sputum were negative.

Which one of the following investigations will be most helpful in making the diagnosis?

- A

Native renal biopsy
- B

Renal ultrasound scan
- C

Urine for crystals
- D

X-ray studies of kidneys, ureters, bladder
- E

Serum immunoglobulins

70058

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 146 of 203

A 38-year-old man with a 12-year history of HIV, with a good response to therapy, was admitted with a 12-day history of weakness, nausea, dysuria, myalgia and decreased urine output. His last CD4 lymphocyte count was $368 \times 10^6/l$ (normal range 500-900). The current medications included Didanosine, Stavudine, Atorvastatin, Trimethoprim-sulfamethoxazole, and he had been commenced on Indinavir. On examination he was afebrile, pulse 100/min and regular, BP 176/99 mmHg, the JVP was not elevated; the chest and abdominal examinations were normal.

Investigations:

Hb	13.5 g/dl
WCC	$6.7 \times 10^9/l$
Platelets	$200 \times 10^9/l$
Na ⁺	135 mmol/l
K ⁺	5.2 mmol/l
Cl ⁻	102 mmol/l
Bicarbonate	16 mmol/l
Urea	29.2 mmol/l
Creatinine	765 micromol/l
Bone profile	normal
Creatine kinase	100 U/l
ANCA	negative
Anti-GBM	negative
Hep B/ Hep C	negative
ASOT	negative

Cultures of blood/urine/sputum were negative.

Which one of the following investigations will be most helpful in making the diagnosis?

- A

Native renal biopsy
- B

Renal ultrasound scan
- C

Urine for crystals
- D

X-ray studies of kidneys, ureters, bladder
- E

Serum immunoglobulins

Explanation

C Urine for crystals

This would confirm nephrolithiasis as the cause of AKI and specifically point to Inidinavir as the causative agent.

A Native renal biopsy

Nephrolithiasis may not be detected and the presence of acute tubular necrosis would not allow you to differentiate the offending agent.

B Renal ultrasound scan

Renal ultrasound scan would be useful if obstruction was suspected or to detect shrunken kidneys from chronic kidney disease. Although the blood pressure is high, there is no sign of fluid overload and the AKI is related to medication changes.

D X-ray studies of kidneys, ureters, bladder

The radiologic imaging procedures typically used in the diagnosis of ureteral stones appear to be unreliable in the diagnosis of non-opaque stones due to Indinavir.

E Serum immunoglobulins

There are no clinical or biochemical parameters to suggest myeloma or light chain deposition disease as the cause of AKI. There is no anemia and bone profile is normal.

70058

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	46
Responses Total:	46
Responses - % Correct:	0%

Back to Filters


Question 147 of 203

A 28-year-old Somali lady presented to the Emergency Department with a 3-week history of generalised weakness, anorexia, decreased appetite, and weight loss. She was noted to be increasingly short of breath and had intermittent headaches. She worked as a cleaner and has been in the UK for 3 years. She was unable to give a sexual history. There was no history of blood transfusions. On examination she appeared cachectic, afebrile, pulse 80/min and regular, BP 130/80 mmHg, JVP not seen. There was no peripheral oedema. Chest examination was normal. Fundoscopy revealed grade 2 hypertensive retinopathy. The remainder of the examination was normal.

Investigations:

Hb	9.2 g/dl
WCC	4.1 x10 ⁹ /l
Platelets	100 x10 ⁹ /l
Blood film	microcytosis
Na ⁺	138 mmol/l
K ⁺	3.2 mmol/l
Urea	18 mmol/l
Creatinine	205 micromol/l
Total protein	58 g/l
Albumin	18 g/l
LFTs	normal
Bone profile	normal
Urine microscopy	Intact red blood cells/granular casts, protein ++
Blood/urine culture	negative
CXR	normal
ECG	normal
Renal ultrasound scan	Highly echogenic kidneys
	Right kidney 13.5 cm, left kidney 14 cm

She underwent a native renal biopsy, the results of which are shown below;



What is the most likely diagnosis?

- A

Primary focal segmental glomerulosclerosis
- B

Schistosomiasis
- C

HIV-associated nephropathy
- D

Heroin-associated nephropathy
- E

Malignant hypertension

70059

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 148 of 203

A 70-year-old white man with a history of intermittent microscopic haematuria presented with generalised fatigue, cough, and vague abdominal pain and diarrhoea. He has a history of hypertension and was on Irbesartan. Baseline renal function was normal when checked by his GP 3 weeks previously.


Initial investigations;

Hb	7.8 g/dl
Haematocrit	0.4
Platelets	98 x109/l
Creatinine	250 micromol/l
Amylase	150 U/l
LFTs	normal

Urine microscopy was not done on admission.

Ultrasound scans of abdomen and pelvis were normal.

The CXR is shown below;



He was treated with Ciprofloxacin for a presumed chest infection. He then developed acute ischaemic bowel requiring resection, with non-specific pathological findings. The hospital course was complicated by hypotension and multi-organ failure. The Creatinine rose to 636 micromol/l with anuria; haemodialysis was initiated.

Urine microscopy	numerous red blood cells
Protein excretion	3g/24 hr
ANA, ANCA	negative
HBsAg	negative
HCV	negative
Cryoglobulins	negative
C3	107 mg/dl (65-190 mg/dl)
C4	38 mg/dl (15-50 mg/dl)
Protein electrophoresis	normal pattern

Renal biopsy was performed which showed crescentic glomerulonephritis with linear IgG staining on immunofluorescence.

What is the most likely diagnosis?

- A

Post-streptococcal glomerulonephritis
- B

Haemolytic uraemic syndrome
- C

Acute pyelonephritis
- D

Microscopic polyangiitis
- E

Anti-glomerular basement membrane disease

70060

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 148 of 203

A 70-year-old white man with a history of intermittent microscopic haematuria presented with generalised fatigue, cough, and vague abdominal pain and diarrhoea. He has a history of hypertension and was on Irbesartan. Baseline renal function was normal when checked by his GP 3 weeks previously.

Initial investigations;

Hb	7.8 g/dl
Haematocrit	0.4
Platelets	98 x109/l
Creatinine	250 micromol/l
Amylase	150 U/l
LFTs	normal

Urine microscopy was not done on admission.

Ultrasound scans of abdomen and pelvis were normal.

The CXR is shown below;



He was treated with Ciprofloxacin for a presumed chest infection. He then developed acute ischaemic bowel requiring resection, with non-specific pathological findings. The hospital course was complicated by hypotension and multi-organ failure. The Creatinine rose to 636 micromol/l with anuria; haemodialysis was initiated.

Urine microscopy	numerous red blood cells
Protein excretion	3g/24 hr
ANA, ANCA	negative
HBsAg	negative
HCV	negative
Cryoglobulins	negative
C3	107 mg/dl (65-190 mg/dl)
C4	38 mg/dl (15-50 mg/dl)
Protein electrophoresis	normal pattern

Renal biopsy was performed which showed crescentic glomerulonephritis with linear IgG staining on immunofluorescence.

What is the most likely diagnosis?

- A

Post-streptococcal glomerulonephritis
- B

Haemolytic uraemic syndrome
- C

Acute pyelonephritis
- D

Microscopic polyangiitis
- E

Anti-glomerular basement membrane disease

Explanation ⚙

- E

Anti-glomerular basement membrane disease

The initial presentation was of a patient with hematuria, subclinical lung haemorrhage and acute kidney injury. Often, lung haemorrhage may present without haemoptysis and in this case was incorrectly diagnosed as chest infection.

The clinical presentation and subsequent rapid deterioration was consistent with the syndrome of alveolar haemorrhage and renal failure caused by anti-glomerular basement membrane (GBM) antibodies (Goodpasture’s syndrome).

Diagnosis is made by the presence of circulating anti-GBM antibodies and on renal biopsy. Typical appearances are of diffuse proliferative glomerulonephritis with variable degrees of necrosis, crescent formation as in this case, glomerulosclerosis and tubular loss.

Treatment is with cytotoxic immunosuppression and plasma exchange. Poor prognostic factors are advanced renal failure on presentation and the degree of crescent formation.

Other causes of this syndrome are Granulomatosis with Polyangiitis (previously referred to as Wegener’s granulomatosis; ANCA-positive in 95% of patients), SLE, Churg-Strauss syndrome and mixed essential cryoglobulinaemia.

- A

Post-streptococcal glomerulonephritis

C3 was normal and PSGN rarely results in nephrotic range proteinuria. Renal biopsy is not performed in most patients to confirm the diagnosis of PSGN, since the resolution of PSGN typically begins within one week of presentation.

- B

Haemolytic uraemic syndrome

The renal biopsy findings point against this.

- C

Acute pyelonephritis

Does not explain the pulmonary findings.

- D

Microscopic polyangiitis

Microscopic polyangiitis may also cause this syndrome, but the negative ANCA makes it unlikely.

70060

Rate this question: ⊖ ★ ★ ★ ★ ★ ★

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	48
Responses Total:	48
Responses - % Correct:	0%

Back to Filters

Question 149 of 203

A 28-year-old man with Alport’s syndrome received a pre-emptive living related renal transplant from his mother. He has excellent primary graft function and the baseline Creatinine was 102 micromol/l. Three months later he presented with a 1-week history of decreased urine output with a raised creatinine of 188 micromol/l.

The biopsy did not show evidence of acute rejection. Urine microscopy showed dysmorphic red blood cells. He became progressively oligo-anuric and became dialysis dependent.

What is the most likely diagnosis?



- | | |
|---|--|
| A | Anti-glomerular basement membrane disease |
| B | Aspergillus pneumonia |
| C | Post-transplant lymphoproliferative disorder |
| D | Mycoplasma pneumonia |
| E | Granulomatosis with polyangiitis |

70061

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 149 of 203

A 28-year-old man with Alport’s syndrome received a pre-emptive living related renal transplant from his mother. He has excellent primary graft function and the baseline Creatinine was 102 micromol/l. Three months later he presented with a 1-week history of decreased urine output with a raised creatinine of 188 micromol/l.

The biopsy did not show evidence of acute rejection. Urine microscopy showed dysmorphic red blood cells. He became progressively oligo-anuric and became dialysis dependent.

What is the most likely diagnosis?



- A

Anti-glomerular basement membrane disease
- B

Aspergillus pneumonia
- C

Post-transplant lymphoproliferative disorder
- D

Mycoplasma pneumonia
- E

Granulomatosis with polyangiitis

Explanation



- A

Anti-glomerular basement membrane disease

Alport’s is an inherited disorder resulting in microscopic haematuria, progressive nephritis with renal impairment, sensorineural deafness and ocular abnormalities. It is due to mutations of tissue-specific type IV collagen chains, leading to formation of autoantibodies similar to those on Goodpasture’s syndrome. Therefore, a minority of Alport’s patients with a renal transplant develop a rapidly progressive glomerulonephritis indistinguishable from Goodpasture’s syndrome but without pulmonary haemorrhage. Treatment is similar to that of de novo anti-GBM disease but treatment efficacy is limited.

- B

Aspergillus pneumonia

Opportunistic infections are common in transplant recipients, especially with over-immunosuppression, but would not explain the graft loss. Other common infections in transplant patients include cytomegalovirus, varicella zoster virus and fungal diseases.

- C

Post-transplant lymphoproliferative disorder

Post-transplant lymphoproliferative disorder is an uncommon but serious manifestation of immunosuppression. The early onset of disease in this patient with likely minimal immunosuppression from a live donor transplant makes this answer incorrect. The majority of PTLD are B-cell clonality associated with Epstein-Barr virus. Treatment is often by a reduction of the immunosuppression and chemotherapy. Graft loss is common.

- D

Mycoplasma pneumonia

Opportunistic infections are common in transplant recipients, especially with over-immunosuppression, but would not explain the graft loss.

- E

Granulomatosis with polyangiitis

The short duration of illness and known association of anti-GBM in patients with Alport’s syndrome.

70061

Rate this question:

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	49
Responses Total:	49
Responses - % Correct:	0%

Back to Filters

Question 150 of 203

A 33-year-old white man was admitted with a 1-week history of rash, pyrexia, generalised weakness and decreased urine output. On examination he was obese, with a temperature of 38.5°C, BP 156/70 mmHg. He had nail-fold vasculitic skin lesions. On auscultation he had a soft systolic murmur. Abdominal examination was limited given his size, but unremarkable.

Initial investigations reveal;

Hb	9 g/dl
WCC	18.5 x10 ⁹ /l
Neutrophils	16.4 x10 ⁹ /l
Platelets	80 x10 ⁹ /l
Creatinine	336 micromol/l
Urine microscopy	Red cell casts
Hepatitis B	negative
Hepatitis C	negative
ANCA/GBM	negative
Cryoglobulins	positive
ASOT	negative
Protein electrophoresis	negative

Ultrasound renal scan of the abdomen showed an 11-cm spleen. Renal biopsy was not performed, as it was not technically possible.

What is the most likely diagnosis?

- A

Microscopic polyangiitis
- B

Infective endocarditis
- C

Acute pyelonephritis
- D

HIV-associated nephropathy
- E

Renal tuberculosis

70062

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 150 of 203

A 33-year-old white man was admitted with a 1-week history of rash, pyrexia, generalised weakness and decreased urine output. On examination he was obese, with a temperature of 38.5°C, BP 156/70 mmHg. He had nail-fold vasculitic skin lesions. On auscultation he had a soft systolic murmur. Abdominal examination was limited given his size, but unremarkable.

Initial investigations reveal;

Hb	9 g/dl
WCC	18.5 x109/l
Neutrophils	16.4 x109/l
Platelets	80 x109/l
Creatinine	336 micromol/l
Urine microscopy	Red cell casts
Hepatitis B	negative
Hepatitis C	negative
ANCA/GBM	negative
Cryoglobulins	positive
ASOT	negative
Protein electrophoresis	negative

Ultrasound renal scan of the abdomen showed an 11-cm spleen. Renal biopsy was not performed, as it was not technically possible.

What is the most likely diagnosis?

- A

Microscopic polyangiitis
- B

Infective endocarditis
- C

Acute pyelonephritis
- D

HIV-associated nephropathy
- E

Renal tuberculosis

Explanation



- B

Infective endocarditis

The history and investigations are consistent with infective endocarditis. One third of patients with bacterial endocarditis develop AKI. The glomerulonephritis is associated with deposition of immune complexes containing bacterial antigens in glomeruli. Cryoglobulins (polyclonal or type III) are present in 50% of subjects.

- A

Microscopic polyangiitis

Microscopic polyangiitis may present with nail-fold vasculitic lesions but these patients are commonly ANCA-positive.

- C

Acute pyelonephritis

The heart murmur, negative urinary sediment and positive cryoglobulins make this unlikely.

- D

HIV-associated nephropathy

There is no weight loss or lymphopenia. HIVAN occurs almost exclusively in black patients, and they tend to be normotensive.

- E

Renal tuberculosis

Renal TB is associated with an insidious onset of renal impairment with asymptomatic urinary abnormalities such as persistent sterile pyuria. It may or may not be associated with extrarenal systemic manifestations.

70062

Rate this question: ⓪★★★★★

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	50
Responses Total:	50
Responses - % Correct:	0%

Question 151 of 203

The following is the MRI scan of his abdomen and brain;




A	Autosomal dominant polycystic kidney disease
B	Autosomal recessive polycystic kidney disease
C	Tuberose sclerosis
D	von-Hippel Lindau disease
E	Histiocytosis X

Submit

Skip Question

Question 151 of 203

The following is the MRI scan of his abdomen and brain;



What is the diagnosis?

- | | |
|---|---|
| A | Autosomal dominant polycystic kidney disease |
| B | Autosomal recessive polycystic kidney disease |
| C | Tuberose sclerosis |
| D | von-Hippel Lindau disease |
| E | Histiocytosis X |

Explanation



- D von-Hippel Lindau disease

Von Hippel-Lindau disease is an autosomal dominant condition that manifests in CNS haemangioblastomas, renal and pancreatic cysts, renal carcinoma and pheochromocytomas. Renal cysts are usually multiple and bilateral and are often associated with solid tumours. Other causes of renal cystic disease include ADPKD, ARPKD, tuberose sclerosis complex, medullary cystic disease and renal cystic dysplasia.

- A Autosomal dominant polycystic kidney disease

This would not explain the MRI brain cyst. ADPKD commonly presents in adulthood and may be associated with intracranial berry aneurysms.

- B Autosomal recessive polycystic kidney disease

This commonly presents in either the neonatal period or up to adolescence with renal impairment.

- C Tuberose sclerosis

In TSC 80% of affected individuals have seizures and manifest with angiomyolipomas in skin and visceral organs at a younger age.

- E Histiocytosis X

Langerhans cell histiocytosis, which is also called Histiocytosis X is incorrect. It is a multisystem disease with predominant bone manifestations.

Rate this question:

[Next Question](#)

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	51
Responses Total:	51
Responses - % Correct:	0%

Back to Filters

Question 152 of 203

A 50-year-old white woman presented to her GP with a 2-week history of nausea, vomiting, and anorexia. She had mild epistaxis and scant haemoptysis for the last 6 to 8 months, with three episodes of sinusitis, treated with antibiotics and inhaled steroids. Her serum creatinine on presentation was 144 micromol/l with protein ++ and blood +++. Urine microscopy revealed granular casts and dysmorphic red blood cells. She was referred to the hospital medical team the next day when her creatinine was 175 micromol/l. Her BP was measured at 160/98 mmHg, with no rashes or other findings on physical examination.

Investigations:

Creatinine	256 micromol/l
Haematocrit	0.36
Platelets	300 x109/l
WCC	6.1 x109/l
ANA	negative
C3	71 mg/dl (65-190 mg/dl)
C4	37 mg/dl (15-50 mg/dl)

What is the most likely diagnosis?

- A

pANCA-positive vasculitis
- B

cANCA-positive vasculitis
- C

Lupus nephritis
- D

Post-streptococcal glomerulonephritis
- E

Henoch-Schönlein purpura

70064

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 152 of 203

Investigations:

Creatinine	256 micromol/l
Haematocrit	0.36
Platelets	300 x10 ⁹ /l
WCC	6.1 x10 ⁹ /l
ANA	negative
C3	71 mg/dl (65-190 mg/dl)
C4	37 mg/dl (15-50 mg/dl)

What is the most likely diagnosis?

- | | |
|---|---------------------------------------|
| A | pANCA-positive vasculitis |
| B | cANCA-positive vasculitis |
| C | Lupus nephritis |
| D | Post-streptococcal glomerulonephritis |
| E | Henoch-Schönlein purpura |

Explanation



- B cANCA-positive vasculitis

This is a classic presentation of Granulomatosis with Polyangiitis (GPA), (previously referred to as Wegener's granulomatosis). It is a necrotising granulomatous vasculitis affecting the small vessels, associated with circulating anti-neutrophil cytoplasmic antibodies (ANCA). It may manifest in a combination of upper and/or lower respiratory tract and/or renal disease. On immunofluorescence, ANCA staining is cytoplasmic with specificity to proteinase-3.

- | | |
|---|---------------------------|
| A | pANCA-positive vasculitis |
|---|---------------------------|

pANCA-positive vasculitis is not classically associated with upper respiratory tract symptoms.

- | | |
|---|-----------------|
| C | Lupus nephritis |
|---|-----------------|

Lupus nephritis is associated with a positive ANA and hypocomplementaemia.

- D Post-streptococcal glomerulonephritis

This does not explain the insidious upper respiratory tract symptoms. In addition, C3 would have likely been low.

- | | |
|---|--------------------------|
| E | Henoch-Schönlein purpura |
|---|--------------------------|

HSP is not classically associated with upper respiratory tract

© 2006 The Authors
Journal compilation © 2006 Blackwell Publishing Ltd

Feedback

End Session

Difficulty: Average

Peer Responses %


Session Progress	
Responses Correct:	0
Responses Incorrect:	52
Responses Total:	52
Responses % Correct:	0%

Question 153 of 203

Investigations:

Creatinine	68 micromol/l
Bilirubin	5 micromol/l
ALT	30 U/l
ALP	98 U/l
Gamma-GT	45 U/l
Total protein	32 g/l
Albumin	18 g/l
Cholesterol	9.8 mmol/l
Urine microscopy	granular casts
Urine protein	14.2 g/day
Hepatitis serology:	
B e antigen	negative
B core IgG antibody	positive
B surface antigen	positive
C antibody	negative

Renal biopsy was performed and is shown below;



What is the most likely diagnosis?

- | | |
|---|--------------------------------------|
| A | Renal TB |
| B | Mesangiocapillary glomerulonephritis |
| C | Minimal change disease |
| D | Diabetic glomerulosclerosis |
| E | Membranous nephropathy |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Question 153 of 203

Investigations,

Renal biopsy was performed and is shown below;



A	Renal TB
B	Mesangiocapillary glomerulonephritis
C	Minimal change disease
D	Diabetic glomerulosclerosis
E	Membranous nephropathy

The answer is Membranous nephropathy -

A	Renal TB
---	----------

B	Mesangiocapillary glomerulonephritis
---	--------------------------------------

C Minimal change disease

D Diabetic glomerulosclerosis

Rate this question: 

Rate this question:

[Next Question](#)

End Session

Difficulty: Average

Responses Correct:	0
Responses Incorrect:	53

[Blog](#) [About Pastest](#) [Contact Us](#) [Help](#)

Question 154 of 203

A	Focal segmental glomerulosclerosis
B	Membranous nephropathy
C	Light-chain disease
D	Diabetic glomerulosclerosis
E	Minimal change disease


Submit

Skip Question

Back to Filters

Question 154 of 203

A 68-year-old Ghanaian man with a 10-year history of type 2 diabetes presented with a 3-week history of generalised oedema. In spite of insulin treatment, his diabetes has been poorly controlled. He has background retinopathy. Investigations revealed normal renal function, albumin 12 g/l, negative ANA, negative ANCA, negative GBM, negative HBsAg, negative hepatitis C antibody, negative cryoglobulins, negative serum electrophoresis. 24 hour urine protein excretion was 15.6. g.He underwent a renal biopsy shown below;



What is the most likely diagnosis?

- A Focal segmental glomerulosclerosis
- B Membranous nephropathy
- C Light-chain disease
- D Diabetic glomerulosclerosis**
- E Minimal change disease

Explanation



- D Diabetic glomerulosclerosis**

The duration of poorly controlled diabetes and evidence of microvascular disease make diabetic nephropathy the most likely diagnosis. The biopsy shows mesangial expansion with a Kimmelstiel-Wilson nodule at the top of the glomerulus, typical of diabetic glomerulosclerosis.

- A Focal segmental glomerulosclerosis

The renal biopsy findings are not consistent with this, along with the long duration of diabetes.

- B Membranous nephropathy

The incidence of membranous nephropathy is increased in diabetics compared with non-diabetic patients but on light microscopy the glomeruli will show capillary loop thickening and normal architecture respectively.

- C Light-chain disease

Inconsistent with the history of poorly controlled diabetes and normal electrophoresis.

- E Minimal change disease

The incidence of minimal-change disease is increased in diabetics compared with non-diabetic patients but on light microscopy the glomeruli will show no change at all or at most mild mesangial proliferation.

70066

Rate this question: ⚙️ ⭐ ⭐ ⭐ ⭐ ⭐

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	54
Responses Total:	54
Responses - % Correct:	0%

Question 155 of 203

Examining his legs you note the following;



Creatinine	162 micromol/l
LFTs	normal
FBC	normal
Urine microscopy	red cell casts
ANCA	negative
ASOT	normal
Hepatitis B/C	negative

A	IgA nephropathy
B	Infective endocarditis
C	Henoch-Schönlein purpura
D	Erythema nodosum
E	Polyarteritis nodosa

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 155 of 203

A 19-year-old man presented with a 2-day history of a painful rash, constipation, lethargy and generalised body pain. He has no significant past medical history apart from hospitalisation as a child for peritonitis secondary to a ruptured appendix.

Examining his legs you note the following;



Investigations;

Creatinine	162 micromol/l
LFTs	normal
FBC	normal
Urine microscopy	red cell casts
ANCA	negative
ASOT	normal
Hepatitis B/C	negative



What is the diagnosis?

- A

IgA nephropathy
- B

Infective endocarditis
- C

Henoch-Schönlein purpura
- D

Erythema nodosum
- E

Polyarteritis nodosa

Explanation ⚙

- C

Henoch-Schönlein purpura

In a young patient with microscopic haematuria and renal impairment, the differential diagnosis includes HSP, IgA nephropathy and thin-membrane disease. HSP is uncommon after the second decade of life and the renal involvement is often transient. The rash is a purpuric vasculitis, usually spreading on the extensor surfaces. Histology shows a leucocytoclastic vasculitis with IgA deposits in blood vessel walls.

- A

IgA nephropathy

Both IgA nephropathy (IgAN) and Henoch-Schönlein nephritis are characterised by mesangial IgA deposition. The latter is differentiated from IgAN by extrarenal manifestations, such as purpura (as in this gentleman), polyarthralgia, and abdominal pain caused by gut vasculitis with IgA deposition.

- B

Infective endocarditis

It does not explain all the extra-renal manifestations and normal FBC.

- D

Erythema nodosum

Erythema nodosum is characterised by red or violet subcutaneous nodules that usually develop in a pretibial location.

- E

Polyarteritis nodosa

PAN typically presents with systemic symptoms (fatigue, weight loss, weakness, fever, arthralgias) and signs (skin lesions, hypertension, renal insufficiency, neurologic dysfunction, abdominal pain) of multisystem involvement.

70067

Rate this question: ⊖ ★ ★ ★ ★ ★

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	55
Responses Total:	55
Responses - % Correct:	0%

Back to Filters

Question 156 of 203

A 24-year-old South Asian woman presented with acute onset of left-sided flank pain with microscopic haematuria. She had a KUB X-ray performed. She has a history of recurrent urinary tract infections since childhood and in the past was investigated and treated by a Urology Specialist. Her maternal first cousin is on peritoneal dialysis for end-stage renal disease.

Her KUB X-ray is shown below;



Her investigations;

Creatinine	96 micromol/l
Ca ²⁺	2.48 mmol/l
PO ₄ ⁻	1.02 mmol/l
Albumin	40 g/l
Urine protein	0.5g/24h
Creatinine clearance	68 ml/minute
24-h urine calcium	10 mmol (<7.5 mmol)
24-h urine oxalate	1.4 mmol (<0.36 mmol)
24-h urine urate	3.2 mmol (<4.5 mmol)

What is the most likely diagnosis?

- A

Primary hyperoxaluria
- B

Primary hyperparathyroidism
- C

Secondary hyperparathyroidism
- D

Nephrogenic diabetes insipidus
- E

Surreptitious antacid ingestion

70068

Submit

Previous Question

Skip Question

Calculator

✔

Normal Values

✔

Back to Filters

Question 156 of 203

A 24-year-old South Asian woman presented with acute onset of left-sided flank pain with microscopic haematuria. She had a KUB X-ray performed. She has a history of recurrent urinary tract infections since childhood and in the past was investigated and treated by a Urology Specialist. Her maternal first cousin is on peritoneal dialysis for end-stage renal disease.

Her KUB X-ray is shown below;



Her investigations;

Creatinine	96 micromol/l
Ca ²⁺	2.48 mmol/l
PO ₄ ⁻	1.02 mmol/l
Albumin	40 g/l
Urine protein	0.5g/24h
Creatinine clearance	68 ml/minute
24-h urine calcium	10 mmol (<7.5 mmol)
24-h urine oxalate	1.4 mmol (<0.36 mmol)
24-h urine urate	3.2 mmol (<4.5 mmol)



What is the most likely diagnosis?

- A

Primary hyperoxaluria
- B

Primary hyperparathyroidism
- C

Secondary hyperparathyroidism
- D

Nephrogenic diabetes insipidus
- E

Surreptitious antacid ingestion

Explanation



- A

Primary hyperoxaluria

Elevated urinary oxalate levels may be due to increased dietary intake, malabsorption or an inherited enzyme deficiency that leads to excessive metabolism of oxalate (primary hyperoxaluria). There are three types: types I and III are due to an enzyme defect in the liver glyoxalate pathway and in type II there is failure of reduction of glyoxalate to glycolate. Type I is the commonest and results in widespread calcium oxalate deposition throughout the body.

Treatment of primary hyperoxaluria is aimed at increasing urinary pH to make calcium oxalate more soluble. This is by administering supplemental citrate and magnesium. Renal insufficiency is common and patients require a combined liver and kidney transplant in type I disease.

- B

Primary hyperparathyroidism

Her normal serum calcium and phosphate levels are not consistent with hyperparathyroidism.

- C

Secondary hyperparathyroidism

Her normal serum calcium and phosphate levels are not consistent with hyperparathyroidism.

- D

Nephrogenic diabetes insipidus

This is characterised by polyuria and polydipsia. The acquired form may be due to electrolyte abnormalities (hypokalaemia, hypercalcaemia), chronic renal failure or lithium therapy. Nephrocalcinosis is uncommon.

- E

Surreptitious antacid ingestion

The recurrent childhood urinary tract infections and family history of renal failure make this unlikely.

70068

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	56
Responses Total:	56
Responses - % Correct:	0%

Question 157 of 203

Investigations;

Hb	6.2 g/dl
Haematocrit	0.24
Platelets	56 x10 ⁹ /l
Blood film	Fragmented red cells
Urea	14 mmol/l
Creatinine	236 micromol/l



- | | |
|---|---|
| A | Transfusion with packed red cells to aim for a haemoglobin of 10 g/dl |
| B | Intravenous vitamin K |
| C | Intravenous Tranexamic acid |
| D | Intravenous fresh frozen plasma |
| E | Haemodialysis |

Submit

Skip Question

Question 157 of 203

Investigations;

Hb	6.2 g/dl
Haematocrit	0.24
Platelets	56 x10 ⁹ /l
Blood film	Fragmented red cells
Urea	14 mmol/l
Creatinine	236 micromol/l

3

- | | |
|---|---|
| A | Transfusion with packed red cells to aim for a haemoglobin of 10 g/dl |
| B | Intravenous vitamin K |
| C | Intravenous Tranexamic acid |
| D | Intravenous fresh frozen plasma |
| E | Haemodialysis |

- | | |
|---|---------------------------------|
| D | Intravenous fresh frozen plasma |
|---|---------------------------------|

A Transfusion with packed red cells to aim for a haemoglobin of 10 g/dl

B	Intravenous vitamin K
---	-----------------------

C	Intravenous Tranexamic acid
---	-----------------------------

E	Haemodialysis
---	---------------

[Next Question](#)

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	57
Responses Total:	57
Responses - % Correct:	0%

Back to Filters

Question 158 of 203

A 44-year-old man has been referred to the nephrologists with a serum creatinine of 210 micromol/l. He has a 15-year history of osteoarthritis. One year ago he was diagnosed with diabetes mellitus and essential hypertension. His current medications include Ramipril 10 mg, Aspirin 75 mg, Diclofenac 150 mg and Atorvastatin 10 mg daily. On examination his BP was 110/70 mmHg. Systemic examination was normal.

Investigations;

Hb	10.9 g/dl
WCC	7.2 x10 ⁹ /l
Platelets	210 x10 ⁹ /l
Na ⁺	138 mmol/l
K ⁺	4.2 mmol/l
Urea	14.2 mmol/l
Creatinine	224 micromol/l
Ca ²⁺	2.28 mmol/l
PO ₄ ⁻	1.05 mmol/l
Albumin	38 g/l
HbA1c	37 mmol/mol (5.5%)
ECG	normal
Urine protein	0.9 g/24h

Renal ultrasound scan reveals unobstructed echogenic kidneys, right kidney 9.4 cm, left kidney 8.8 cm

What is the most likely cause of his renal impairment?

- A

Renovascular disease
- B

Hypertensive nephrosclerosis
- C

Analgesic nephropathy
- D

Herbal nephropathy
- E

Diabetic nephropathy

70070

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 158 of 203

A 44-year-old man has been referred to the nephrologists with a serum creatinine of 210 micromol/l. He has a 15-year history of osteoarthritis. One year ago he was diagnosed with diabetes mellitus and essential hypertension. His current medications include Ramipril 10 mg, Aspirin 75 mg, Diclofenac 150 mg and Atorvastatin 10 mg daily. On examination his BP was 110/70 mmHg. Systemic examination was normal.

Investigations;

Hb	10.9 g/dl
WCC	7.2 x10 ⁹ /l
Platelets	210 x10 ⁹ /l
Na ⁺	138 mmol/l
K ⁺	4.2 mmol/l
Urea	14.2 mmol/l
Creatinine	224 micromol/l
Ca ²⁺	2.28 mmol/l
PO ₄ ⁻	1.05 mmol/l
Albumin	38 g/l
HbA1c	37 mmol/mol (5.5%)
ECG	normal
Urine protein	0.9 g/24h

Renal ultrasound scan reveals unobstructed echogenic kidneys, right kidney 9.4 cm, left kidney 8.8 cm

What is the most likely cause of his renal impairment?

- A

Renovascular disease
- B

Hypertensive nephrosclerosis
- C

Analgesic nephropathy
- D

Herbal nephropathy
- E

Diabetic nephropathy

Explanation Settings icon

- C

Analgesic nephropathy

The patient in question has chronic renal failure due to the long-term use of non-steroidal anti-inflammatory drugs (NSAIDs). Chronic NSAID abuse leads to renal injury due to renal papillary necrosis and chronic interstitial nephritis. The degree of renal injury correlates with the duration of use and analgesic load. There is no specific treatment apart from avoidance of NSAIDs. In this situation the proteinuria raises the possibility of a membranous nephritis pattern of disease.

- A

Renovascular disease

As he is on ACE inhibitors, the absence of bruits and symmetrical kidneys make renovascular disease unlikely.

- B

Hypertensive nephrosclerosis

The duration of diabetes mellitus and the normal blood pressure with no evidence of other end-organ damage (normal fundi, no LVH on voltage criteria in the ECG) excludes hypertensive nephrosclerosis.

- D

Herbal nephropathy

There is no clue in the history that he has been taking over-the-counter herbal medications. Some Chinese herbs have been implicated in the development of rapidly progressive renal failure due to interstitial nephritis and progressive tubulo-interstitial fibrosis.

- E

Diabetic nephropathy

The duration of diabetes mellitus and the normal blood pressure with no evidence of other end-organ damage (normal fundi, no LVH on voltage criteria in the ECG) excludes diabetic nephropathy.

70070

Rate this question: Feedback icon Star 1 Star 2 Star 3 Star 4 Star 5 Star 6

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	58
Responses Total:	58
Responses - % Correct:	0%

Back to Filters

Question 159 of 203

A 66-year-old man on peritoneal dialysis for end-stage renal failure presents with a 3-month history of tiredness secondary to anaemia, in spite of increased subcutaneous erythropoietin therapy.

Investigations;

Hb	6.2 g/dl
WCC	6.2 x10 ⁹ /l
Differential WCC	normal
Platelets	200 x10 ⁹ /l
Haematocrit	0.2
MCV	80 fl
Reticulocyte count	0%
Faecal occult bloods	negative
Colonoscopy	normal
Upper GI endoscopy	normal
Haptoglobins	normal

What is the most likely cause of his anaemia?

- A

Hypothyroidism
- B

Myelodysplasia
- C

Myelofibrosis
- D

Pure red cell aplasia
- E

Chronic lymphocytic leukaemia

70071

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 159 of 203

A 66-year-old man on peritoneal dialysis for end-stage renal failure presents with a 3-month history of tiredness secondary to anaemia, in spite of increased subcutaneous erythropoietin therapy.

Investigations;

Hb	6.2 g/dl
WCC	6.2 x10 ⁹ /l
Differential WCC	normal
Platelets	200 x10 ⁹ /l
Haematocrit	0.2
MCV	80 fl
Reticulocyte count	0%
Faecal occult bloods	negative
Colonoscopy	normal
Upper GI endoscopy	normal
Haptoglobins	normal

What is the most likely cause of his anaemia?

- A

Hypothyroidism
- B

Myelodysplasia
- C

Myelofibrosis
- D

Pure red cell aplasia
- E

Chronic lymphocytic leukaemia

Explanation



- D

Pure red cell aplasia

Pure red cell aplasia (PRCA) is a rare condition defined by the absence of erythroblasts in the bone marrow, leading to profound anaemia with normal leucocyte and platelet counts, and characterised by low or absent circulating reticulocytes. Serum iron and ferritin rise sharply as iron cannot be incorporated into the erythrocytes.

PRCA occurs due to neutralising antibodies to erythropoietin. There has been a recent increase in PRCA in patients with chronic kidney disease on subcutaneous erythropoietin alpha. Treatment includes discontinuation of the EPO preparation and repeated transfusions.

- A

Hypothyroidism

The normal MCV and WCC make hypothyroidism unlikely.

- B

Myelodysplasia

The normal MCV and WCC make myelodysplasia unlikely. The myelodysplastic syndromes comprise a heterogeneous group of malignant hematopoietic stem cell disorders characterised by dysplastic and ineffective blood cell production and a variable risk of transformation to acute leukaemia.

- C

Myelofibrosis

In myelofibrosis the blood film shows a leucoerythroblastic picture with an increased reticulocyte count.

- E

Chronic lymphocytic leukaemia

The normal MCV and WCC make chronic lymphocytic leukaemia unlikely.

70071

Rate this question:

Next Question

- Previous Question

Tag Question
- Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	59
Responses Total:	59
Responses - % Correct:	0%

Question 160 of 203

Investigations:

Hb	9 g/dl
Haematocrit	0.3
MCV	80 fl
Urea	21mmol/l
Creatinine	423 micromol/l
PTH	8.5 pmol/l
Ferritin	14 microg/l



- | | |
|---|-----------------------------|
| A | Oral ferrous sulphate |
| B | Intravenous iron |
| C | Intravenous erythropoietin |
| D | Red cell transfusion |
| E | Subcutaneous erythropoietin |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 160 of 203

A 34-year-old man with chronic kidney disease is seen in the Nephrology Clinic. He is asymptomatic. His current medications include atenolol 100 mg once daily, Ramipril 10 mg once daily, alfacalcidol 0.25 micrograms once daily, Calcichew[®] 1 tablet with meals and a multivitamin preparation

Investigations:

Hb	9 g/dl
Haematocrit	0.3
MCV	80 fl
Urea	21mmol/l
Creatinine	423 micromol/l
PTH	8.5 pmol/l
Ferritin	14 microg/l

What is the definitive intervention for his anaemia?



- A

Oral ferrous sulphate
- B

Intravenous iron
- C

Intravenous erythropoietin
- D

Red cell transfusion
- E

Subcutaneous erythropoietin

Explanation



- B

Intravenous iron

Anaemia is almost universal in patients with chronic kidney disease; it is characteristically normochromic and normocytic and is associated with erythropoietin (Epo) deficiency and shortening of red cell survival.

Intravenous iron is the most appropriate intervention taking into account the very low ferritin of 14 and given that oral iron absorption is often defective in chronic renal failure.

- A

Oral ferrous sulphate

Due to defective oral iron absorption, and the fact that the ferritin level is only 14, oral iron replacement is highly unlikely to be effective.

- C

Intravenous erythropoietin

This is only used in dialysis patients and given via the dialysis circuit.

- D

Red cell transfusion

Red cell transfusions should be avoided unless in an emergency, as this patient will develop circulating antibodies which would make future transplantation more difficult.

- E

Subcutaneous erythropoietin

Eventually, patients are commenced on erythropoietin. Iron stores should be optimised before beginning erythropoietin.

70314

Rate this question:

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %


Session Progress

Responses Correct:	0
Responses Incorrect:	60
Responses Total:	60
Responses - % Correct:	0%

Back to Filters

Question 161 of 203

The pedigree of a family affected by a genetic disease linked to progressive renal failure, hearing loss and haematuria come to the clinic for review.



Given the suspected diagnosis, which of the following correctly reflects the inheritance pattern for this condition?

- A Autosomal dominant
- B Autosomal recessive
- C X-linked dominant
- D X-linked recessive
- E X-linked recessive plus spontaneous mutation

70496

Submit

Previous Question Skip Question

Calculator

Normal Values

Question 161 of 203

A	Autosomal dominant
B	Autosomal recessive
C	X-linked dominant
D	X-linked recessive
E	X-linked recessive plus spontaneous mutation

The answer is C), X-linked dominant -

Autosomal dominant, A), is incorrect. The fact that this is a renal condition suggestive of Alport's drives us towards X-linked dominant as the likely diagnosis.

Autosomal recessive, B), is incorrect. The high rates of heritability from generation to generation, (up to 50% of offspring), are much more suggestive of either X-lined dominant or autosomal dominant inheritance.

X-linked recessive, D), is incorrect. The fact that females are affected by the condition effectively rules out x-linked recessive inheritance.

X-linked recessive plus spontaneous mutation, E), is incorrect. This would not be consistent with the number of females affected by the condition.

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %


Session Progress

Responses Correct:	0
Responses Incorrect:	61
Responses Total:	61
Responses - % Correct:	0%

Back to Filters

Question 162 of 203

A 45-year-old man presents with a history of poor urine output over the past 2 days. He has recently returned from a holiday in Thailand, after which he had suffered diarrhoea about a week ago; otherwise he has no other significant history. On examination he had a blood pressure of 98/50 mmHg and a pulse rate of 110 bpm. A diagnosis is reached of haemolytic-uraemic syndrome associated with diarrhoea. Gram stain of the organism found in the stool is shown below.



What would be the most useful treatment for this patient?

- | | |
|---|---|
| A | Antibiotics |
| B | Diuretics |
| C | Heparin |
| D | Intravenous immunoglobulins |
| E | Supportive (dialysis, fluids, blood transfusions) |

70755

Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Back to Filters

Question 162 of 203

A 45-year-old man presents with a history of poor urine output over the past 2 days. He has recently returned from a holiday in Thailand, after which he had suffered diarrhoea about a week ago; otherwise he has no other significant history. On examination he had a blood pressure of 98/50 mmHg and a pulse rate of 110 bpm. A diagnosis is reached of haemolytic-uraemic syndrome associated with diarrhoea. Gram stain of the organism found in the stool is shown below.



What would be the most useful treatment for this patient?

- A Antibiotics
- B Diuretics
- C Heparin
- D Intravenous immunoglobulins
- E Supportive (dialysis, fluids, blood transfusions)

Explanation



- E Supportive (dialysis, fluids, blood transfusions)

Supportive treatment is the mainstay of treatment for haemolytic-uraemic syndrome (HUS). Although in some cases, plasma exchange will remove the inhibitor of the von Willebrand factor protease enzyme. About 70% of affected patients will recover with no long-term problems with supportive treatment alone.

- A Antibiotics

There is no evidence that antibiotics will shorten the duration of illness in HUS. For infective diarrhoea, in general, giving antibiotics may prolong infection.

- B Diuretics

Giving diuretics may potentially worsen the acute renal failure by causing further hypotension. This man needs fluid resuscitation as evidenced by the low blood pressure and tachycardia.

- C Heparin

Anticoagulation is contraindicated in HUS, particularly in children, because the condition is frequently complicated by both bleeding and hypertension. In adults, heparin increases mortality because of increased bleeding.

- D Intravenous immunoglobulins

There is no role of immunoglobulin replacement in HUS. There are some immune disorders whereby replacement of immunoglobulins may be useful such as multiple sclerosis, but not in this case.

70755

Rate this question: ⚙️ ⭐ ⭐ ⭐ ⭐ ⭐

Next Question

- Previous Question
- Tag Question
- Feedback
- End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	62
Responses Total:	62
Responses - % Correct:	0%

Question 163 of 203

Blood results at this time show:

Na ⁺	139 mmol/l
K ⁺	4.6 mmol/l
Urea	15.0 mmol/l
Creatinine	179 μmol/l
CRP	150 mg/l
ESR	87 mm/h
ANA	Positive
ANCA	Negative
Hb	10 g/dl
WCC	11.4 × 10 ⁹ /l
Neutrophils	8.0 × 10 ⁹ /l
Eosinophils	0.09 × 10 ⁹ /l
PLT	560 × 10 ⁹ /l
HBsAg	Positive
HBeAg	Negative
Hep B anti-core AB	Positive
Hep C AB	Negative

21

- | | |
|---|----------------------------------|
| A | Microscopic polyangiitis |
| B | Polyarteritis nodosa |
| C | Streptokinase-induced vasculitis |
| D | Systemic lupus erythematosus |
| E | Wegener's granulomatosis |

Submit

Skip Question

Back to Filters

Question 163 of 203

A 42-year-old banker was admitted with a 2-hour history of central crushing chest pain radiating down his left arm. He admits to abusing intravenous drugs in the past and is a life-long smoker. His past medical history includes hypertension and irritable bowel syndrome. On admission, a fast-track troponin level was raised. ECG demonstrated anterolateral ST segments elevated >2 mm. He was treated for a myocardial infarction (MI) with intravenous thrombolysis followed by intravenous heparin infusion. ST segments returned to normal. Three days later in hospital this man developed a purpuric rash over his legs. On examination, there were normal heart sounds with no added murmurs. Chest examination revealed clinically a small left-sided pleural effusion. The abdominal examination was unremarkable. Neurological examination revealed weakness of his left arm: the patient felt this has been present for a while now since sustaining an injury playing football. ECG demonstrated sinus rhythm and anterolateral Q waves.

Blood results at this time show:

Na ⁺	139 mmol/l
K ⁺	4.6 mmol/l
Urea	15.0 mmol/l
Creatinine	179 μmol/l
CRP	150 mg/l
ESR	87 mm/h
ANA	Positive
ANCA	Negative
Hb	10 g/dl
WCC	11.4 × 10 ⁹ /l
Neutrophils	8.0 × 10 ⁹ /l
Eosinophils	0.09 × 10 ⁹ /l
PLT	560 × 10 ⁹ /l
HBsAg	Positive
HBeAg	Negative
Hep B anti-core AB	Positive
Hep C AB	Negative

What is the most likely diagnosis?

- A

Microscopic polyangiitis
- B

Polyarteritis nodosa
- C

Streptokinase-induced vasculitis
- D

Systemic lupus erythematosus
- E

Wegener’s granulomatosis

Explanation

- B

Polyarteritis nodosa

The American College of Rheumatology states that patients must have three of the following to have a diagnosis of polyarteritis nodosa (PAN): weight loss >5 kg, livedo reticularis, testicular pain/tenderness, myalgia, mono/polyneuropathy, diastolic BP >90 mmHg, elevated urea/creatinine, hepatitis B virus. This man clearly has a systemic vasculitis given a history of cardiac, gastrointestinal and renal involvement. The recent neurological involvement clinches the diagnosis of PAN.

- A

Microscopic polyangiitis

In this scenario, the test for antineutrophil cytoplasmic antibodies (ANCA) is negative which rules out microscopic polyangiitis.

- C

Streptokinase-induced vasculitis

A vasculitis secondary to streptokinase occurs mainly in those who have received previous streptokinase therapy and does not explain the other manifestations. It is an immune-complex phenomenon and antistreptokinase antibodies can be measured. Renal impairment can occur.

- D

Systemic lupus erythematosus

Patients with SLE tend to present with other skin manifestations and low platelets.

- E

Wegener’s granulomatosis

Wegener’s granulomatosis affects the respiratory tract more commonly. There is no mention of ear, nose and throat symptoms in this case.

70756

Rate this question: ⚙️☆☆☆☆☆

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	63
Responses Total:	63
Responses - % Correct:	0%

Back to Filters

Question 164 of 203

A 40-year-old man who is undergoing regular haemodialysis is receiving erythropoietin, but his Hb is still low. There is no evidence of active or past infection. Other than chronic renal failure of unknown cause, he has no other past medical history. He is on oral iron, which was started after a confirmed low ferritin. Upon examination, his BMI is 28 kg/m², and chest and abdominal examination was unremarkable.

Investigations showed:

Na ⁺	138 mmol/l
K ⁺	4.8 mmol/l
Urea	15.0 mmol/l
Creatinine	530 μmol/l
CRP	20 mg/l
Hb	7 g/dl
WCC	11.4 × 10 ⁹ /l
PLT	560 × 10 ⁹ /l

What is the next step in his management?

- A

Check for antibodies against erythropoietin
- B

Give higher doses of iron tablets
- C

Give folic acid
- D

Give IV iron
- E

Transfuse

70757

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 164 of 203

Investigations showed:

Na ⁺	138 mmol/l
K ⁺	4.8 mmol/l
Urea	15.0 mmol/l
Creatinine	530 μmol/l
CRP	20 mg/l
Hb	7 g/dl
WCC	$11.4 \times 10^9/l$
PLT	$560 \times 10^9/l$

1.

2.

3.

- | | |
|---|---|
| A | Check for antibodies against erythropoietin |
| B | Give higher doses of iron tablets |
| C | Give folic acid |
| D | Give IV iron |
| E | Transfuse |

- | | |
|---|--------------|
| D | Give IV iron |
|---|--------------|

A Check for antibodies against erythropoietin

B Give higher doses of iron tablets

C	Give folic acid
---	-----------------

E	Transfuse
---	-----------

[Next Question](#)

Th O Li

End Session

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	64
Responses Total:	64
Responses - % Correct:	0%

Back to Filters

Question 165 of 203

A 40-year-old woman attended Accident and Emergency Department. She is on established peritoneal dialysis (PD) and has presented with generalised abdominal pain, vomiting and pyrexia. She has a past medical history of hypertension. On questioning, she says her menstruation is erratic and she has not had a menstrual period for several months now. Furthermore, she says she has not passed any urine for a few days. On physical examination, she has a temperature of 38°C, heart rate 100 bpm and a blood pressure of 140/80 mmHg. Her abdomen is distended and there is generalised tenderness with no guarding or rigidity.

Investigations showed:

Na ⁺	136 mmol/l
K ⁺	4.9 mmol/l
Urea	15.0 mmol/l
Creatinine	530 μmol/l
CRP	150 mg/l
Hb	9 g/dl
WCC	15.4 × 10 ⁹ /l
PLT	560 × 10 ⁹ /l

What is the likely cause of her abdominal pain?

- A

Appendicitis
- B

Ectopic pregnancy
- C

Menstruation
- D

Peritonitis
- E

UTI

70758

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 165 of 203

Investigations showed:

Na ⁺	136 mmol/l
K ⁺	4.9 mmol/l
Urea	15.0 mmol/l
Creatinine	530 μmol/l
CRP	150 mg/l
Hb	9 g/dl
WCC	15.4 × 10 ⁹ /l
PLT	560 × 10 ⁹ /l

A	Appendicitis
B	Ectopic pregnancy
C	Menstruation
D	Peritonitis
E	UTI

D	Peritonitis
---	-------------

A	Appendicitis
---	--------------

B	Ectopic pregnancy
---	-------------------

C	Menstruation
---	--------------

Rate this question:

[Previous Question](#)

[Tag Question](#)

[Feedback](#)

[End Session](#)

Peer Responses %

Session Progress	
Responses Correct:	0
Responses Incorrect:	65
Responses Total:	65
Percentage of 64 Questions:	0%

Question 166 of 203

Investigations showed:

Na ⁺	138 mmol/l
K ⁺	4.8 mmol/l
Urea	15.0 mmol/l
Creatinine	320 μmol/l
CRP	20 mg/l
Hb	15 g/dl
WCC	11.4 × 10 ⁹ /l
PLTg	560 × 10 ⁹ /l
Erythropoietin level	Raised

11

- | | |
|---|--|
| A | Arterial blood gases (ABGs) |
| B | Bone marrow examination to exclude myeloproliferative disorder |
| C | Chest X-ray |
| D | Lactate dehydrogenase (LDH) levels |
| E | Renal USS |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 166 of 203

A 25-year-old woman presents to medical clinic complaining of back pain. Past medical history of note includes hypertension which is currently monitored by the GP. Apparently one of her family died suddenly of a stroke at the age of 41. On physical examination, her blood pressure is 175/80 mmHg, BMI was 27 kg/m², her chest was clear and abdomen examination was unremarkable.

Investigations showed:

Na ⁺	138 mmol/l
K ⁺	4.8 mmol/l
Urea	15.0 mmol/l
Creatinine	320 μmol/l
CRP	20 mg/l
Hb	15 g/dl
WCC	11.4 × 10 ⁹ /l
PLTg	560 × 10 ⁹ /l
Erythropoietin level	Raised

What investigation should be performed next?



- A

Arterial blood gases (ABGs)
- B

Bone marrow examination to exclude myeloproliferative disorder
- C

Chest X-ray
- D

Lactate dehydrogenase (LDH) levels
- E

Renal USS

Explanation



- E

Renal USS

The combination of back pain and raised Hb should raise the suspicion of renal cystic disease. One cause may be autosomal-dominant polycystic kidney disease (ADPKD) accounting for the death of a relative at an early age. Because this lady is 25 years old, the kidneys would not have been so enlarged to have been detectable on physical examination, therefore the abdominal examination is unremarkable. APKD is a common autosomal dominant disease with a gene defect on the short arm of chromosome 16 and usually presents in the fourth and fifth decades. APKD may present at any age from the second decade with acute loin pain or haematuria due to cyst haemorrhage or infection, loin pain due to increased renal size, subarachnoid haemorrhage due to berry aneurysm rupture, complications of hypertension, complications of liver cyst formation or symptoms of chronic renal failure.

- A

Arterial blood gases (ABGs)

The ABG result will be normal in ADKPD and therefore would not help in discerning any differentials.

- B

Bone marrow examination to exclude myeloproliferative disorder

The bone marrow in ADPKD may be normal providing there is no associated renal bone disease associated with renal failure. There would be an increased production of erythropoietin from the kidneys though.

- C

Chest X-ray

In ADPKD, the chest X-ray is typically normal.

- D

Lactate dehydrogenase (LDH) levels

LDH may be abnormal in myeloproliferative and lymphoproliferative disorders. Haemolytic anaemia would be one example where LDH would be elevated. In ADPKD, the LDH level would be normal.

70759

Rate this question:

Next Question

- Previous Question

Tag Question
- Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	66
Responses Total:	66
Responses - % Correct:	0%

Back to Filters

Question 167 of 203

A 19-year-old man with malaria has been admitted to the Zambia University Teaching Hospital. He has no past medical history of note. On admission he was started on oral chloroquine. After being on the ward for 3 days, his condition continues to worsen and a nurse notices that he has not passed urine. He is already managed with IV fluids.

His blood results are as follows:

Na ⁺	138 mmol/l
K ⁺	6.2 mmol/l
Urea	54 mmol/l
Creatinine	785 mol/l
Hb	8.7 g/dl
MCV	80 fl
WCC	6 × 10 ⁹ /l
Platelets	213 × 10 ⁹ /l
Blood film	+++ for ring forms

The patient is catheterised and only 50 ml of dark urine is drained. He is given IV calcium for CV protection because of his potassium.

What is the appropriate step in the immediate management of this patient’s infection?

- A

Arrange a blood transfusion
- B

Peritoneal dialysis
- C

Repeat the blood film
- D

Start intravenous quinine
- E

Urine microscopy and culture

70760

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 167 of 203

His blood results are as follows.

Na	138 mmol/l
K ⁺	6.2 mmol/l
Urea	54 mmol/l
Creatinine	785 μmol/l
Hb	8.7 g/dl
MCV	80 fl
WCC	$6 \times 10^9/l$
Platelets	$213 \times 10^9 /l$
Blood film	+++ for ring forms

What is the appropriate step in the immediate management of this patient's infection?

- | | |
|---|------------------------------|
| A | Arrange a blood transfusion |
| B | Peritoneal dialysis |
| C | Repeat the blood film |
| D | Start intravenous quinine |
| E | Urine microscopy and culture |

- D Start intravenous quinine

A Arrange a blood transfusion

There is no role for blood transfusion in malaria treatment unless there is co-existent symptomatic anaemia.

- B Peritoneal dialysis

This patient has acute renal failure which may be reversible with malaria treatment and supportive care which includes intravenous fluids. There is no indication for acute dialysis in this scenario. Acute dialysis may be required if there is pulmonary oedema resistant to medical treatment, life-threatening hyperkalaemia, flash pulmonary oedema, metabolic acidosis resistant to medical treatment or uraemic pericarditis or encephalopathy.

- C Repeat the blood film

A repeat blood film will confirm the diagnosis, but will not be helpful in this scenario as the patient needs treatment.

- E Urine microscopy and culture

In this scenario, he has not passed urine because he has acute renal failure; therefore waiting to obtain a urine sample would not be an appropriate option. Furthermore, this would not help with the diagnosis of malaria

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	67
Responses Total:	67
Responses - % Correct:	0%

External Links

cks.nice.org.uk/malaria#!scenario



Back to Filters

Question 168 of 203

A 61-year-old woman is being reviewed in the medical follow-up clinic. She received a renal transplant 6 months ago because of end-stage renal failure. She has been fine but complains that she feels embarrassed whenever she laughs because her gums have become very prominent. Her past medical history, apart from the renal transplant, includes long-standing hypertension. Ciclosporin was started after her renal transplant, and this is the only medication she is taking. On examination, she has a blood pressure of 140/80 mmHg, BMI of 27 kg/m², gums are swollen, chest is clear, and non-tender renal transplant.

Investigations show:

Na ⁺	138 mmol/l
K ⁺	4.2 mmol/l
Urea	12 mmol/l
Creatinine	250 mol/l
Hb	8.7 g/dl
WCC	6 × 10 ⁹ /l
PLT	213 × 10 ⁹ /l

What other complications of ciclosporin treatment should this patient be aware of?

- A

Alopecia
- B

Hypertrichosis
- C

Pigmentation
- D

Purpura
- E

Staining of the teeth

70761

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 168 of 203

Investigations show:

Na ⁺	138 mmol/l
K ⁺	4.2 mmol/l
Urea	12 mmol/l
Creatinine	250 μmol/l
Hb	8.7 g/dl
WCC	$6 \times 10^9 / l$
PLT	$213 \times 10^9 / l$

- | | |
|---|-----------------------|
| A | Alopecia |
| B | Hypertrichosis |
| C | Pigmentation |
| D | Purpura |
| E | Staining of the teeth |

- | | |
|---|----------------|
| B | Hypertrichosis |
|---|----------------|

A	Alopecia
---	----------

Alopecia is a known side-effect of another renal transplant immunosuppressant – tacrolimus.

- C Pigmentation

Ciclosporin does not affect pigmentation of the skin. Any change in pigmentation of moles in renal transplant should be taken seriously because there is an increased chance of malignancy whilst taking immunosuppression. Renal transplant patients are often advised to wear sun block when exposed to the sun.

- | | |
|---|---------|
| D | Purpura |
|---|---------|

Warfarin may have the side effect of purpura.

- E Staining of the teeth

Minocycline and doxycycline may have the side effects of dental staining

[Next Question](#)

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	68
Responses Total:	68
Responses - % Correct:	0%

Back to Filters

Question 169 of 203

A 42-year-old man with severe acute left-sided flank pain is referred by his GP to A&E. This is the first time he has had such severe sudden pain. He has no past medical history of cardiovascular or renal disease, his brother died of a subarachnoid haemorrhage. On examination his blood pressure is 170/100 mmHg, pulse 78 bpm and regular. His jugular venous pressure (JVP) is not raised, heart sounds are normal and his chest is clear. His liver is not palpable but he has a mass in the left flank. The mass is bimanually palpable and you can get above it.

Blood results are as follows:

Na ⁺	143 mmol/l
K ⁺	4.2 mmol/l
Urea	10.2 mmol/l
Creatinine	205 μmol/l
Hb	15.0 g/dl
WCC	5.1 × 10 ⁹ /l
MCV	81 fl
Platelets	243 × 10 ⁹ /l
ESR	8 mm/h
Urine dipstick	Blood +++, protein +

Which first line investigation would you request?

- A

CT scan of the abdomen
- B

Intravenous urogram (IVU)
- C

MRI scan of the abdomen
- D

Ultrasound of the abdomen
- E

Urine microscopy

70762

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 169 of 203

A 42-year-old man with severe acute left-sided flank pain is referred by his GP to A&E. This is the first time he has had such severe sudden pain. He has no past medical history of cardiovascular or renal disease, his brother died of a subarachnoid haemorrhage. On examination his blood pressure is 170/100 mmHg, pulse 78 bpm and regular. His jugular venous pressure (JVP) is not raised, heart sounds are normal and his chest is clear. His liver is not palpable but he has a mass in the left flank. The mass is bimanually palpable and you can get above it.

Blood results are as follows:

Na ⁺	143 mmol/l
K ⁺	4.2 mmol/l
Urea	10.2 mmol/l
Creatinine	205 μmol/l
Hb	15.0 g/dl
WCC	5.1 × 10 ⁹ /l
MCV	81 fl
Platelets	243 × 10 ⁹ /l
ESR	8 mm/h
Urine dipstick	Blood +++, protein +

Which first line investigation would you request?

- A

CT scan of the abdomen
- B

Intravenous urogram (IVU)
- C

MRI scan of the abdomen
- D

Ultrasound of the abdomen
- E

Urine microscopy

Explanation



- D

Ultrasound of the abdomen

Ultrasound of the abdomen is easily performed, non-invasive and plays a critical diagnostic role. The suspicion here would be one of polycystic kidney disease. There is an association with Berry aneurysm which may well have resulted in the death of his brother. Aggressive management of hypertension is the most appropriate way to prevent or at least slow the accelerated deterioration in renal function associated with this condition.

- A

CT scan of the abdomen

CT scan of the abdomen is incorrect. Although a CT scan of the abdomen would also show polycysts in the kidneys, it would not be as easily performed compared with ultrasound. It would also expose the patient to radiation.

- B

Intravenous urogram (IVU)

Intravenous urogram (IVU) is incorrect. IVU may be normal in polycystic kidney disease. There would be associated unnecessary contrast exposure.

- C

MRI scan of the abdomen

MRI scan of the abdomen is incorrect. MRI compared with ultrasound abdomen is more difficult to perform. There is nephrogenic systemic fibrosis with the use of contrast agent in MRI scans in patients with renal failure.

- E

Urine microscopy

Urine microscopy is incorrect. The urine microscopy would typically be normal in polycystic kidney disease unless there are associated complications such as bleeding in the cysts where there may be haematuria.

70762

Rate this question:

Next Question

- Previous Question

Tag Question
- Feedback

End Session

Difficulty: Easy

Peer Responses %

Session Progress


Responses Correct:	0
Responses Incorrect:	69
Responses Total:	69
Responses - % Correct:	0%

Back to Filters

Question 170 of 203

A 44-year-old woman presents to her GP for review. She has attended the GP on a number of occasions during the past year for sinusitis and is now concerned that the bridge of her nose has collapsed and that she may require cosmetic surgery. Upon examination, blood pressure is 160/70 mmHg, chest is clear and abdominal examination is unremarkable. Chest X-ray is abnormal and reveals multiple nodules. Routine blood testing reveals a creatinine of 205 μ mol/l.

Renal biopsy reveals:



Which of the following is the best initial treatment for this condition?

- | | |
|---|--------------------------------------|
| A | Azathioprine |
| B | Corticosteroids |
| C | Cyclophosphamide |
| D | Corticosteroids and cyclophosphamide |
| E | Methotrexate |

70763

Submit

Previous QuestionSkip Question

Calculator✔


Normal Values✔

Back to Filters

Question 170 of 203

A 44-year-old woman presents to her GP for review. She has attended the GP on a number of occasions during the past year for sinusitis and is now concerned that the bridge of her nose has collapsed and that she may require cosmetic surgery. Upon examination, blood pressure is 160/70 mmHg, chest is clear and abdominal examination is unremarkable. Chest X-ray is abnormal and reveals multiple nodules. Routine blood testing reveals a creatinine of 205 μ mol/l.

Renal biopsy reveals:



Which of the following is the best initial treatment for this condition?

- A Azathioprine
- B Corticosteroids
- C Cyclophosphamide
- D Corticosteroids and cyclophosphamide**
- E Methotrexate

Explanation



- D Corticosteroids and cyclophosphamide**

The clinical history is highly suggestive of Wegener's granulomatosis. Often patients present with severe rhinorrhoea, complicated by nasal ulceration and later cough with haemoptysis. Chest X-ray reveals nodular changes \pm pneumonic features. Renal involvement is characterised by microvascular glomerulonephritis, which is what is illustrated in the picture. Treatment is with cyclophosphamide in combination with corticosteroids in patients with severe disease.

- A Azathioprine

Azathioprine may be used in the maintenance therapy for Wegener's granulomatosis when the patient is in remission.

- B Corticosteroids

Corticosteroids would treat the inflammation, but would not be used alone in therapy. Before the introduction of steroids, over 75% of patients who had vital organ involvement and vasculitis would die. However, after the introduction of steroids, 5-year mortality fell to 50% and further decreased to 12% with the addition of cytotoxic drugs for polyarteritis nodosa. Few data exist for the use of steroids alone for Wegener's granulomatosis, but we would expect a similar reduction in death; therefore the addition of cytotoxic medications in addition to steroids is important.

- C Cyclophosphamide

Cyclophosphamide is a well-established immunosuppressant therapy for vasculitis, but it would not be used alone in the case of severe disease.

- E Methotrexate

Methotrexate would be used in the treatment of rheumatoid vasculitis but not in Wegener's granulomatosis.

70763

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	70
Responses Total:	70
Responses - % Correct:	0%

Question 171 of 203

His twin brother's investigations:

Sodium	140 mmol/l
Potassium	4.2 mmol/l
Urea	7.2 mmol/l
Creatinine	80 μmol/l
Hb	12.0 g/dl
WCC	$5.1 \times 10^9/l$
Urine dipstick	Trace protein

-
-
-

- | | |
|---|-----------------------|
| A | Heterotopic isograft |
| B | Heterotopic xenograft |
| C | Orthotopic allograft |
| D | Orthotopic autograft |
| E | Orthotopic isograft |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Question 171 of 203

His twin brother's investigations:

Sodium	140 mmol/l
Potassium	4.2 mmol/l
Urea	7.2 mmol/l
Creatinine	80 μmol/l
Hb	12.0 g/dl
WCC	$5.1 \times 10^9/l$
Urine dipstick	Trace protein

This type of transplantation is described as:

- | | |
|---|-----------------------|
| A | Heterotopic isograft |
| B | Heterotopic xenograft |
| C | Orthotopic allograft |
| D | Orthotopic autograft |
| E | Orthotopic isograft |

Explanation

- | | |
|---|----------------------|
| A | Heterotopic isograft |
|---|----------------------|

Transplants are classified as:

- autografts: in which the same individual acts as both donor and recipient;
- isografts: in which the donor and recipient are genetically identical;
- allografts: where the donor and recipient are genetically dissimilar but belong to the same species;
- xenografts: in which the donor and recipient belong to different species.

In orthotopic transplants the transplanted part is placed in its normal anatomical location, while in heterotopic transplants it is placed in a different anatomical location.

- | | |
|---|-----------------------|
| B | Heterotopic xenograft |
|---|-----------------------|

Xenografts are from different species such as pig and human.

- | | |
|---|----------------------|
| C | Orthotopic allograft |
|---|----------------------|

Donor and recipient are genetically dissimilar but belong to the same species.

- | | |
|---|----------------------|
| D | Orthotopic autograft |
|---|----------------------|

The same individual acts as both donor and recipient.

- | | |
|---|---------------------|
| E | Orthotopic isograft |
|---|---------------------|

The transplanted part is placed in the normal anatomical location.

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	71
Responses Total:	71
Responses - % Correct:	0%

Back to Filters

Question 172 of 203

A 61-year-old man who came to your clinic complains of weakness and loss of weight. His voice is hoarse. He has no past medical history of note. On examination he has pedal oedema up to the tibial tuberosity, enlarged tongue and liver edge palpable up to three finger breadths below the right costal margin.

Investigations

Na ⁺	140 mmol/l
K ⁺	4.2 mmol/l
Urea	7.2 mmol/l
Creatinine	80 mol/l
Hb	12.0 g/dl
WCC	5.1 × 10 ⁹ /l
Urine dipstick	3+++ protein

What is the most appropriate test that will lead to the correct diagnosis?

- A

Liver spleen scan
- B

24 h urinary protein
- C

Subcutaneous fat biopsy
- D

Urine protein electrophoresis
- E

Urine protein immunophoresis

70765

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 172 of 203

A 61-year-old man who came to your clinic complains of weakness and loss of weight. His voice is hoarse. He has no past medical history of note. On examination he has pedal oedema up to the tibial tuberosity, enlarged tongue and liver edge palpable up to three finger breadths below the right costal margin.

Investigations

Na ⁺	140 mmol/l
K ⁺	4.2 mmol/l
Urea	7.2 mmol/l
Creatinine	80 mol/l
Hb	12.0 g/dl
WCC	5.1 × 10 ⁹ /l
Urine dipstick	3+++ protein

What is the most appropriate test that will lead to the correct diagnosis?

- ALiver spleen scan
- B24 h urinary protein
- CSubcutaneous fat biopsy
- DUrine protein electrophoresis
- EUrine protein immunophoresis

Explanation

- CSubcutaneous fat biopsy

The clinical features of the patient are suggestive of amyloidosis. It is characterised by extracellular deposition of fibrous protein in various tissues and organs. It may be primary or associated with other chronic diseases like myeloma or rheumatoid arthritis. Diagnosis is made by typical findings and demonstration of amyloid fibrils by Congo red staining under polarised light. Abdominal subcutaneous fat pad aspirate or rectal submucosal biopsy are often performed to reach a final diagnosis.

- ALiver spleen scan

In amyloid there may be an enlarged spleen and liver. This is not specific for amyloidosis. Lymphoproliferative diseases may also cause hepatosplenomegaly. Therefore, this would not be the investigation of choice for diagnosis.

- B24 h urinary protein

Heavy proteinuria is a feature of amyloidosis. However, proteinuria may be found in other glomerular diseases such as minimal-change glomerulopathy and membranous glomerulopathy. Therefore, quantification of protein would not give the definitive diagnosis.

- DUrine protein electrophoresis

Urine protein electrophoresis would be abnormal in myeloma.

- EUrine protein immunophoresis

Similarly, in myeloma the urine immunophoresis would be abnormal.

70765

Rate this question: ⚙️ ⭐ ⭐ ⭐ ⭐ ⭐

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	72
Responses Total:	72
Responses - % Correct:	0%

Question 173 of 203

Investigations:

Na ⁺	140 mmol/l
K ⁺	5.5 mmol/l
Urea	15 mmol/l
Creatinine	230 μmol/l
Hb	12.0 g/dl
WCC	$5.1 \times 10^9/l$
Urine dipstick	Red blood cells (RBCs) +++ White blood cells (WBCs) +

11

- | | |
|---|---------------------------------|
| A | Multiple myeloma |
| B | Nephrolithiasis with haematuria |
| C | Profound dehydration |
| D | Rhabdomyolysis |
| E | Viral syndrome |

Submit

Skip Question

Question 173 of 203

Investigations:

Na ⁺	140 mmol/l
K ⁺	5.5 mmol/l
Urea	15 mmol/l
Creatinine	230 μmol/l
Hb	12.0 g/dl
WCC	5.1 × 10 ⁹ /l
Urine dipstick	Red blood cells (RBCs) +++ White blood cells (WBCs) +

- | | |
|---|---------------------------------|
| A | Multiple myeloma |
| B | Nephrolithiasis with haematuria |
| C | Profound dehydration |
| D | Rhabdomyolysis |
| E | Viral syndrome |

- | | |
|---|----------------|
| D | Rhabdomyolysis |
|---|----------------|

A Multiple myeloma

B Nephrolithiasis with haematuria

C	Profound dehydration
---	----------------------

E	Viral syndrome
---	----------------

Rate this question:

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	73
Responses Total:	73
Responses - % Correct:	0%

Back to Filters

Question 174 of 203

A 23-year-old male presents with a 4-day history of sore throat. He also mentions a change in colour of urine and it looks blood-stained. There are no other associated urinary complaints. His past medical history is unremarkable. Upon examination his temperature is 36°C, blood pressure 130/80 mmHg, chest and abdominal examination unremarkable. Chest X-ray is normal. Blood results are normal. Renal biopsy shows mesangial proliferation with positive immunofluorescence for immunoglobulin A (IgA) and C3.

What is the most likely diagnosis?

- A

Adult polycystic kidney disease
- B

Henoch-Schönlein purpura
- C

IgA nephropathy
- D

Kawasaki disease
- E

Post-streptococcal glomerulonephritis

70767

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 174 of 203

A 23-year-old male presents with a 4-day history of sore throat. He also mentions a change in colour of urine and it looks blood-stained. There are no other associated urinary complaints. His past medical history is unremarkable. Upon examination his temperature is 36°C, blood pressure 130/80 mmHg, chest and abdominal examination unremarkable. Chest X-ray is normal. Blood results are normal. Renal biopsy shows mesangial proliferation with positive immunofluorescence for immunoglobulin A (IgA) and C3.

What is the most likely diagnosis?



- A

Adult polycystic kidney disease
- B

Henoch-Schönlein purpura
- C

IgA nephropathy
- D

Kawasaki disease
- E

Post-streptococcal glomerulonephritis

Explanation



- C

IgA nephropathy

Immunoglobulin A (IgA) nephropathy, or Berger’s disease, is the commonest cause of glomerulonephritis worldwide. A typical patient is a young male with asymptomatic microscopic haematuria or episodic macroscopic haematuria, precipitated by infection (e.g. pharyngitis). Recovery is usually rapid between attacks. The pathology consists of focal proliferative glomerulonephritis with mesangial deposits of IgA. Prognosis is usually good in patients with normal blood pressure, normal kidney function and absence of proteinuria at presentation. Approximately 20% of patients develop renal failure 20 years from the time of diagnosis.

- A

Adult polycystic kidney disease

Polycystic kidney disease is not associated with sore throat or any abnormality in immunofluorescence. Diagnosis is with genetic studies or ultrasound identification of cysts.

- B

Henoch-Schönlein purpura

There would be a history of skin rash with this condition.

- D

Kawasaki disease

Kawasaki disease is associated with sore throat. However, it generally occurs in children and there may be coronary artery aneurysms and peeling of the skin of the hands and feet.

- E

Post-streptococcal glomerulonephritis

Post-streptococcal glomerulonephritis is characterised by sudden onset of haematoproteinuria, oedema and hypertension. It is not associated with predominantly IgA in the renal biopsy.

70767

Rate this question:

Next Question

- Previous Question

Feedback
- Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	74
Responses Total:	74
Responses - % Correct:	0%

Back to Filters

Question 175 of 203

A 39-year-old man comes to the clinic for follow-up after a short hospital admission for an episode of renal colic. At the time of admission the patient had an intravenous pyelogram performed that showed mid-ureteric calculi on the left side. There was delayed uptake and excretion of the contrast in the left kidney, consistent with obstruction. There was also a filling defect at the level of L5 consistent with the calcification seen on pre-contrast films. While in the hospital, he underwent ureteroscopic stone extraction of the left mid-ureteral calculi. The calculi were sent to the laboratory for chemical analysis, and the image is shown below. He denies any history of calculi in the past or history of calculi in the family.

Image of calculi:



What will be the likely composition of the stones?

A	Calcium oxalate
B	Cysteine
C	Oxalate
D	Magnesium-ammonium-phosphate
E	Uric acid

70768

Submit

Previous Question

Skip Question

Calculator

✔

Normal Values

✔

Back to Filters

Question 175 of 203

A 39-year-old man comes to the clinic for follow-up after a short hospital admission for an episode of renal colic. At the time of admission the patient had an intravenous pyelogram performed that showed mid-ureteric calculi on the left side. There was delayed uptake and excretion of the contrast in the left kidney, consistent with obstruction. There was also a filling defect at the level of L5 consistent with the calcification seen on pre-contrast films. While in the hospital, he underwent ureteroscopic stone extraction of the left mid-ureteral calculi. The calculi were sent to the laboratory for chemical analysis, and the image is shown below. He denies any history of calculi in the past or history of calculi in the family.

Image of calculi:



What will be the likely composition of the stones?

- A Calcium oxalate
- B Cysteine
- C Oxalate
- D Magnesium-ammonium-phosphate
- E Uric acid

Explanation ⚙

- A Calcium oxalate

This patient has a radio-opaque calculus. Calcium oxalate-containing stones account for nearly 60% of all kidney stones. They are radio-opaque. Cysteine stones result from cysteinuria, which is due to an inherited autosomal recessive disorder of renal tubular reabsorption of four amino acids. Uric acids stones are not radio-opaque. Magnesium-ammonium-phosphate stones are infectious in aetiology. The most common organisms are *Proteus*, *Kleibsiella* and *Pseudomonas*.

- B Cysteine

Cysteine stones are suspected in childhood stone formers. Mutation in the gene encoding an amino acid transporter leads to wasting of the cationic amino acids cysteine, ornithine, lysine and arginine.

- C Oxalate

Hyperoxaluria can be associated with increased dietary oxalate or enhanced oxalate uptake due to ileal inflammation or resection, such as in inflammatory disease, and may lead to the formation of oxalate stones. Normally stones are in combination with calcium rather than oxalate alone.

- D Magnesium-ammonium-phosphate

These stones are formed as a result of infection by *Proteus*, *Klebsiella* or *Serratia*, which are capable of splitting urea to ammonium and hydroxyl ions (thus increasing urinary pH), predisposing towards struvite stones. Such stones act as a reservoir for infection, and often expand to fill as much of the collecting system. Typically seen on KUB.

- E Uric acid

Pure uric acid stones are uncommon. Increase in urinary urate with reduction in urinary pH to less than 5.5 leads to the formation of insoluble uric acid stones. Increase in urinary urate occurs when there is hyperuricaemia (10-20% will have frank gout) of any cause. Uric acid stones are radiolucent on KUB.

70768

Rate this question: ⊖ ★ ★ ★ ★ ★

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	75
Responses Total:	75
Responses - % Correct:	0%

Back to Filters

Question 176 of 203

A 35-year-old man attends the renal clinic for the first time. He has been referred by his GP because of a progressive decline in renal function over several months. His past medical history includes type 2 diabetes mellitus, repeated urinary tract infections as a child and recent admission to hospital with renal colic. On examination, he is apyrexial, his blood pressure is 180/70 mmHg, and chest and abdominal examination are unremarkable.

Investigations:

Sodium	140 mmol/l
Potassium	5.5 mmol/l
Urea	15 mmol/l
Creatinine	230 mol/l
HbA1C	6% (126 mg/dl)
Hb	10.0 g/dl
WCC	5.1 × 10 ⁹ /l
Urine dipstick	No red blood cells White blood cells (WBCs) ++
Urine microscopy	WBC casts

What is the most likely diagnosis?

- A

Diabetic nephropathy
- B

Hypertensive nephropathy
- C

Reflux nephropathy
- D

Tuberculosis
- E

Urinary tract infection

70769

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 176 of 203

Sodium	140 mmol/l
Potassium	5.5 mmol/l
Urea	15 mmol/l
Creatinine	230 μmol/l
HbA1C	6% (126 mg/dl)
Hb	10.0 g/dl
WCC	$5.1 \times 10^9/l$
Urine dipstick	No red blood cells White blood cells (WBCs) ++
Urine microscopy	WBC casts

- | | |
|---|--------------------------|
| A | Diabetic nephropathy |
| B | Hypertensive nephropathy |
| C | Reflux nephropathy |
| D | Tuberculosis |
| E | Urinary tract infection |

- | | |
|---|--------------------|
| C | Reflux nephropathy |
|---|--------------------|

A	Diabetic nephropathy
---	----------------------

B Hypertensive nephropathy

D	Tuberculosis
---	--------------

E	Urinary tract infection
---	-------------------------

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

100

Responses incorrect:	76
Responses Total:	76
Responses - % Correct:	0%

Question 177 of 203

Investigations:

Na ⁺	140 mmol/l
K ⁺	5.5 mmol/l
Urea	15 mmol/l
Creatinine	230 μmol/l
Hb	10.8 g/dl
WCC	12.0 × 10 ⁹ /l
ESR	84 mm/h
Urine dipstick	Blood ++ Protein ++
Chest X-ray	Cavitating lesion in the right upper lobe

What is the most likely diagnosis?

- | | |
|---|---------------------------------------|
| A | Churg-Strauss syndrome |
| B | Wegener's granulomatosis |
| C | Polyarteritis nodosa |
| D | Post-streptococcal glomerulonephritis |
| E | Subacute bacterial endocarditis |

Submit

[Previous Question](#)

Skip Question

Calculator

Normal Values

Back to Filters

Question 177 of 203

A 31-year-old, previously healthy man presents with a 6-month history of nasal congestion and sinusitis. Over a 6-month period he was treated with antibiotic therapy, but his symptoms persisted. His current complaints include frontal headache and a nose bleed. On examination, he is pyrexial and dyspnoeic. He has a rash on his legs (image below) and a right foot drop.

Investigations:

Na ⁺	140 mmol/l
K ⁺	5.5 mmol/l
Urea	15 mmol/l
Creatinine	230 μmol/l
Hb	10.8 g/dl
WCC	12.0 × 10 ⁹ /l
ESR	84 mm/h
Urine dipstick	Blood ++ Protein ++
Chest X-ray	Cavitating lesion in the right upper lobe

Rash:



What is the most likely diagnosis?

- A Churg–Strauss syndrome
- B Wegener’s granulomatosis**
- C Polyarteritis nodosa
- D Post-streptococcal glomerulonephritis
- E Subacute bacterial endocarditis

Explanation

- B Wegener’s granulomatosis**

Wegener’s granulomatosis (WG) is a necrotising vasculitis involving the respiratory tract and kidneys. It causes epistaxis, sinusitis, destruction of the nasal cartilage and glomerulonephritis and renal failure. The serum contains antibodies reacting with the cytoplasm of neutrophils (cANCA).

WG is an idiopathic, systemic inflammatory disease characterised by the presence of granulomas, necrosis and/or vasculitis. Although WG typically affects the upper and lower airways and the kidneys, it may involve any organ system. WG is classified, along with microscopic polyangiitis (MPA) and Churg–Strauss syndrome (CSS), as a primary systemic small-medium-sized vessel vasculitis associated with anti-neutrophil cytoplasmic antibodies (ANCAs). However, WG can also affect medium and even large arteries, and may lack an association with ANCAs. It is a renal/pulmonary disease which often presents with haemoptysis, sinusitis, oral lesions and otitis.

The diagnosis of WG is based on a combination of clinical, laboratory and, if necessary, pathological features. If a typical clinical picture is associated with a positive ANCA finding with specificity for proteinase 3 (PR3), the diagnosis of WG can be presumed. It remains incumbent for the clinician to be certain that WG ‘mimics’, especially granulomatous infection, are not present. In the setting of only a moderate suspicion of WG and a negative ANCA result, it would be judicious to pursue histological support for the diagnosis. The outcome of WG has been dramatically altered by treatment with glucocorticoids (GCs) and cyclophosphamide (CP).

- A Churg–Strauss syndrome

There would be respiratory symptoms such as asthmatic wheeze. Investigations in Churg–Strauss syndrome will show microscopic evidence of red cells and casts in the urine, with protein on dipstick. In the full blood count there would be either prominent eosinophilia (>10% total white cell count or absolute count of >1.5 × 10⁹/l) or a predominance of eosinophils in the biopsy of affected tissues. This eosinophilia is highly steroid-responsive and resolves rapidly on treatment.

- C Polyarteritis nodosa

Polyarteritis nodosa is a systemic vasculitis. There would be palpable purpura and livedo reticularis. Nerve involvement may include paraesthesia. There is no sinusitis. Investigations in polyarteritis nodosa may include a raised creatinine kinase due to muscle injury, raised ESR and ANCA negative. Hepatitis B would be positive in the minority. The diagnosis is made on angiography, which would show arterial aneurysms in the renal or mesenteric tree.

- D Post-streptococcal glomerulonephritis

This generally presents with sore throat. The streptococcal titre would be positive. There would be no cavitating lung lesions on chest X-ray, as present in this case.

- E Subacute bacterial endocarditis

You would expect mention of a murmur with endocarditis. Duke’s criteria are used for diagnosis of endocarditis.

70770

Rate this question:

Next Question

Previous Question Tag Question

Feedback End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	77
Responses Total:	77
Responses - % Correct:	0%

Question 178 of 203

Investigations

Serum calcium	2.05 mmol/l
Serum phosphate	1.8 mmol/l
Serum ALP	170 U/l
Na ⁺	143 mmol/l
K ⁺	5.5 mmol/l
Cr	220 μmol/l
Urea	9.1 mmol/l

3

- | | |
|---|--------------------------------|
| A | Aluminium toxicity |
| B | Decreased vitamin D metabolism |
| C | Osteomalacia |
| D | Primary hyperparathyroidism |
| E | Secondary hyperparathyroidism |

Submit

Skip Question

Question 178 of 203

Investigations

Which of the following pathological processes is an unlikely contributing factor to the above findings?

- ### Explanation

- A Aluminium toxicity

B Decreased vitamin D metabolism

C	Osteomalacia
---	--------------

E Secondary hyperparathyroidism

Rate this question:

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Easy

Peer Responses %

Back to Filters

Question 179 of 203

A 45-year-old male from Ghana underwent a live-related renal transplant 3 weeks ago for end-stage renal failure due to hypertensive nephropathy. His current medications are: ciclosporin, mycophenolate mofetil, prednisolone, cotrimoxazole, ranitidine, amlodipine and atenolol. There were no complications after the procedure and his creatinine normalised a few days after surgery. He has no symptoms. Two days ago his serum creatinine was 155 $\mu\text{mol/l}$.

Investigations

Na ⁺	143 mmol/l
K ⁺	4.5 mmol/l
Cr	165 $\mu\text{mol/l}$
Urea	10.5 mmol/l

What definitive investigation will confirm that acute rejection is the problem?

- A

24 h urine with creatinine estimation
- B

Ciclosporin level
- C

Renal biopsy
- D

Renal ultrasound scan
- E

Serum immunoglobulin

70772

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 179 of 203

A 45-year-old male from Ghana underwent a live-related renal transplant 3 weeks ago for end-stage renal failure due to hypertensive nephropathy. His current medications are: ciclosporin, mycophenolate mofetil, prednisolone, cotrimoxazole, ranitidine, amlodipine and atenolol. There were no complications after the procedure and his creatinine normalised a few days after surgery. He has no symptoms. Two days ago his serum creatinine was 155 mol/l.

Investigations

Na ⁺	143 mmol/l
K ⁺	4.5 mmol/l
Cr	165 mol/l
Urea	10.5 mmol/l

What definitive investigation will confirm that acute rejection is the problem?

- A

24 h urine with creatinine estimation
- B

Ciclosporin level
- C

Renal biopsy
- D

Renal ultrasound scan
- E

Serum immunoglobulin

Explanation



- C

Renal biopsy

Acute rejection develops in 30-50% of renal transplant recipients. The risk is greater in the first 3 weeks after transplant. In patients on ciclosporin there are often no symptoms. Patients with deteriorating renal function with normal ciclosporin level should be investigated by percutaneous renal biopsy.

- A

24 h urine with creatinine estimation

Creatinine estimation would not help in this situation because a rise in serum creatinine is a good enough indication that there may be abnormality in renal function. Urinary creatinine would only be helpful in conditions such as alcoholism or extremes in weight.

- B

Ciclosporin level

A high ciclosporin level may affect creatinine levels by causing acute toxicity. However, this will not determine whether there is any acute rejection.

- D

Renal ultrasound scan

Renal ultrasound scan will identify any obstruction or possible urinary leak post-transplant. It cannot diagnose acute rejection.

- E

Serum immunoglobulin

There are no characteristic changes in serum immunoglobulins that are associated with rejection.

70772

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	79
Responses Total:	79
Responses - % Correct:	0%

Back to Filters

Question 180 of 203

A 17-year-old patient attends a follow-up clinic appointment. She has a family history of autosomal dominant polycystic kidney disease. Her older brother was recently diagnosed with the condition and she was referred by her general practitioner for screening. She has remained asymptomatic and was seen in clinic previously by a colleague. On physical examination she has a blood pressure of 135/80 mmHg, with chest and abdominal examination unremarkable.

Investigations:

Na ⁺	143 mmol/l
K ⁺	3.9 mmol/l
Urea	7 mmol/l
Creatinine	70 μmol/l
Renal ultrasound	No renal cysts detected

What is the next best management step in this case?

- A

Cerebral magnetic resonance angiogram
- B

Computerised tomography of the abdomen
- C

Discharge from clinic
- D

Repeat renal ultrasound after 1 year
- E

Repeat renal ultrasound after 3 years

70773

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 180 of 203

A 17-year-old patient attends a follow-up clinic appointment. She has a family history of autosomal dominant polycystic kidney disease. Her older brother was recently diagnosed with the condition and she was referred by her general practitioner for screening. She has remained asymptomatic and was seen in clinic previously by a colleague. On physical examination she has a blood pressure of 135/80 mmHg, with chest and abdominal examination unremarkable.

Investigations:

Na ⁺	143 mmol/l
K ⁺	3.9 mmol/l
Urea	7 mmol/l
Creatinine	70 μmol/l
Renal ultrasound	No renal cysts detected

What is the next best management step in this case?

- A

Cerebral magnetic resonance angiogram
- B

Computerised tomography of the abdomen
- C

Discharge from clinic
- D

Repeat renal ultrasound after 1 year
- E

Repeat renal ultrasound after 3 years

Explanation



- E

Repeat renal ultrasound after 3 years

Autosomal polycystic kidney disease (ADPKD) genes are located on chromosomes 16 (majority) and 4 (minority). The best screening tool is ultrasound. In patients below the age of 20 years you can get false-negative test results, and it is recommended to screen patients after the age of 20 years.

- A

Cerebral magnetic resonance angiogram

There is an association with intra-cranial aneurysms and ADPKD, but screening for these is not indicated in this patient. Routine screening for intracranial aneurysms is recommended only for high-risk patients, such as those with a previous rupture, a positive family history of an intra-cerebral bleed, warning symptoms, and prior surgery that is likely to be associated with hypertension.

- B

Computerised tomography of the abdomen

A computerised tomography scan, although more sensitive than ultrasound, involves a high radiation dose to this young girl.

- C

Discharge from clinic

This girl is may have ADPKD, which would have future implications. Therefore, it is essential to follow up and not simply discharge from the clinic.

- D

Repeat renal ultrasound after 1 year

1 year would be too early - this girl would not be 20 years old at this point. As explained above, there would be false-negative test results when screening too early.

70773

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	80
Responses Total:	80
Responses - % Correct:	0%

Back to Filters

Question 181 of 203

A previously fit and well 23-year-old woman fell down a flight of stairs. A magnetic resonance scan of the spine has revealed cord compression and she is awaiting the neurosurgeons. She is catheterised and a urine sample is sent to the lab. Her past medical history has no illness of note. The house officer is contacted with the urine microscopy result and phones you for advice. Urine microscopy reveals the presence of epithelial squames, a few red cells of normal morphology are detected and hyaline and fine granular casts are present. There are no white cells and no bacteria on Gram stain.

What is the most likely cause of this result?

- A

Acute tubular necrosis
- B

Atypical urinary tract infection
- C

Glomerulonephritis
- D

Normal urine
- E

Renal haematoma

70774

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 181 of 203

A previously fit and well 23-year-old woman fell down a flight of stairs. A magnetic resonance scan of the spine has revealed cord compression and she is awaiting the neurosurgeons. She is catheterised and a urine sample is sent to the lab. Her past medical history has no illness of note. The house officer is contacted with the urine microscopy result and phones you for advice. Urine microscopy reveals the presence of epithelial squames, a few red cells of normal morphology are detected and hyaline and fine granular casts are present. There are no white cells and no bacteria on Gram stain.

What is the most likely cause of this result?

- A

Acute tubular necrosis
- B

Atypical urinary tract infection
- C

Glomerulonephritis
- D

Normal urine
- E

Renal haematoma

Explanation

- D

Normal urine

The above result is compatible with a normal urine specimen. The squames and normal morphology red cells are probably the result of the catheterisation procedure. Fine granular casts and hyaline casts are precipitated protein and occur in normal urine.

- A

Acute tubular necrosis

In acute tubular necrosis, the presence of ‘muddy brown casts’ is pathognomonic for the condition.

- B

Atypical urinary tract infection

White blood cells and white cell casts suggest conditions such as urinary tract infection and pyelonephritis.

- C

Glomerulonephritis

Dysmorphic red blood cells can suggest glomerular disease. Red cell casts suggest glomerulonephritis. Urine microscopy can reveal bacteria, yeasts, and crystals.

- E

Renal haematoma

Renal haematoma may show red cells or red cell casts in the urine.

70774

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	81
Responses Total:	81
Responses - % Correct:	0%

Back to Filters

Question 182 of 203

A 56-year-old lady is admitted to the ward. She developed a sore throat a week ago and the general practitioner commenced her on penicillin. She now feels more unwell and has developed generalised joint pains and shivering attacks with a rash. For bipolar disorder she has also been taking lithium for the past 2 years. On physical examination her blood pressure is 140/80 mmHg, and chest and abdominal examination is unremarkable.

Investigations:

Haemoglobin	12.2 g/dl
White cell count	11.2 × 10 ⁹ /l
Platelets	Platelets
Na ⁺	135 mmol/l
K ⁺	4.3 mmol/l
Urea	15.2 mmol/l
Creatinine	160 μmol/l
Urine microscopy	Eosinophils, red blood cells

What is the most likely diagnosis?

- A

Acute tubulo-interstitial nephritis
- B

Immunoglobulin A (IgA) nephropathy
- C

Lithium toxicity
- D

Post-streptococcal glomerulonephritis
- E

Systemic lupus erythematosus

70775

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 182 of 203

A 56-year-old lady is admitted to the ward. She developed a sore throat a week ago and the general practitioner commenced her on penicillin. She now feels more unwell and has developed generalised joint pains and shivering attacks with a rash. For bipolar disorder she has also been taking lithium for the past 2 years. On physical examination her blood pressure is 140/80 mmHg, and chest and abdominal examination is unremarkable.

Investigations:

Haemoglobin	12.2 g/dl
White cell count	11.2 × 10 ⁹ /l
Platelets	Platelets
Na ⁺	135 mmol/l
K ⁺	4.3 mmol/l
Urea	15.2 mmol/l
Creatinine	160 μmol/l
Urine microscopy	Eosinophils, red blood cells

What is the most likely diagnosis?



- A

Acute tubulo-interstitial nephritis
- B

Immunoglobulin A (IgA) nephropathy
- C

Lithium toxicity
- D

Post-streptococcal glomerulonephritis
- E

Systemic lupus erythematosus

Explanation



- A

Acute tubulo-interstitial nephritis

Acute tubule-interstitial nephritis presents with arthralgia, skin rashes, fever and renal failure. It is mainly caused by drug reactions, commonly non-steroidal anti-inflammatory agents, antibiotics (penicillin, sulphonamides, cephalosporins, rifampicin, quinolones), allopurinol and phenytoin. Investigations can reveal eosinophilia, urinary eosinophils and raised immunoglobulin E (IgE) levels.

- B

Immunoglobulin A (IgA) nephropathy

There would not be eosinophils in the urine. There would be haematuria. IgA nephropathy is the most common primary glomerulonephritis in the world.

- C

Lithium toxicity

Lithium toxicity presents with neurological symptoms of tremor, weakness, and dysarthria. It can also cause nephrogenic diabetes insipidus.

- D

Post-streptococcal glomerulonephritis

Although blood and protein in the urine after an episode of sore throat may suggest post-streptococcal glomerulonephritis, there would not be any eosinophils in the urine.

- E

Systemic lupus erythematosus

Systemic lupus erythematosus can cause a chronic tubule-interstitial nephritis. Apart from joint pains there are no other symptoms of SLE, such as malar rash, alopecia or weakness.

70775

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	82
Responses Total:	82
Responses - % Correct:	0%

Back to Filters

Question 183 of 203

A 52-year-old gentleman with diabetes presents with weight loss, fevers and dull, persistent left loin pain. The general practitioner has been treating him for relapsing urinary tract infections with oral antibiotics. On examination, his temperature is 37.9°C, blood pressure is 130/80 mmHg and chest and abdominal examination is unremarkable.

Investigations:

Haemoglobin	14.3 g/dl
White cell count	18 × 10 ⁹ /l
Platelets	510 × 10 ⁹ /l
Na ⁺	145 mmol/l
K ⁺	4.2 mmol/l
Urea	12 mmol/l
Creatinine	120 μmol/l
ESR	67 mm/h (0-20)

Computerised tomography scan shows a heterogeneous non-enhancing mass on the left kidney, which is hydronephrotic. The right kidney is normal.

Renal biopsy shows lipid-laden macrophages with lymphocytes and polymorphonuclear leucocytes.

What is the definitive treatment in this patient?

- A

Antituberculous treatment
- B

Intravenous antibiotics
- C

Left nephrectomy
- D

Lithotripsy
- E

Radiotherapy

70776

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 183 of 203

Investigations:

Haemoglobin	14.3 g/dl
White cell count	$18 \times 10^9/\text{l}$
Platelets	$510 \times 10^9/\text{l}$
Na ⁺	145 mmol/l
K ⁺	4.2 mmol/l
Urea	12 mmol/l
Creatinine	120 $\mu\text{mol/l}$
ESR	67 mm/h (0-20)

Renal biopsy shows lipid-laden macrophages with lymphocytes and polymorphonuclear leucocytes.

What is the definitive treatment in this patient?

- | | |
|---|---------------------------|
| A | Antituberculous treatment |
| B | Intravenous antibiotics |
| C | Left nephrectomy |
| D | Lithotripsy |
| E | Radiotherapy |

Explanation

- | | |
|---|------------------|
| C | Left nephrectomy |
|---|------------------|

Xanthogranulomatous pyelonephritis can be difficult to distinguish from a renal cell carcinoma. It typically presents with symptoms of fever, weight loss and loin pain. It is more common in diabetics, the immunocompromised and patients with obstructive uropathy. The most common organism is *Proteus mirabilis*. Histology combined with clinical and radiological evidence confirms the diagnosis. The definitive treatment is nephrectomy, as medical therapy is inefficient.

- | | |
|---|---------------------------|
| A | Antituberculous treatment |
|---|---------------------------|

This man does not have renal tuberculosis. Tuberculosis of the kidneys would present with sterile pyuria. Diagnosis may be obtained with three early morning urine cultures.

- B Intravenous antibiotics

It is difficult for intravenous antibiotics alone to treat xanthogranulomatous pyelonephritis completely.

- | | |
|---|-------------|
| D | Lithotripsy |
|---|-------------|

Lithotripsy is an ultrasound treatment for renal stones. It is not utilised in this case.

- | | |
|---|--------------|
| E | Radiotherapy |
|---|--------------|

Radiotherapy will not help in the treatment for this mass. As it is an infection, it is important to include antibiotics and nephrectomy for treatment

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	83
Responses Total:	83
Responses - % Correct:	0%

Back to Filters

Question 184 of 203

A 29-year-old woman with renal disease secondary to systemic lupus erythematosus is seen in the rheumatology clinic. She has had three urinary tract infections in the past year and was recently admitted to the emergency department with acute severe unilateral abdominal pain which settled spontaneously after several hours. On physical examination, her blood pressure is 130/80 mmHg and chest and abdominal examination is unremarkable.

On her last outpatient visit a number of investigations were requested, the results of which are now available:

Na ⁺	141 mmol/l
Serum potassium	3.3 mmol/l
Urea	9.0 mmol/
Creatinine	188 μmol/l
Serum bicarbonate	8 mmol/l
Urine	pH 7.4

What is the most likely underlying cause of these results and possibly for some of her recent presentations?

- A

Bartter syndrome
- B

Type 1 (distal) renal tubular acidosis
- C

Type 2 (proximal) renal tubular acidosis
- D

Type 4 renal tubular acidosis
- E

Staghorn calculus leading to recurrent urinary sepsis

70778

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 184 of 203

A 29-year-old woman with renal disease secondary to systemic lupus erythematosus is seen in the rheumatology clinic. She has had three urinary tract infections in the past year and was recently admitted to the emergency department with acute severe unilateral abdominal pain which settled spontaneously after several hours. On physical examination, her blood pressure is 130/80 mmHg and chest and abdominal examination is unremarkable.

On her last outpatient visit a number of investigations were requested, the results of which are now available:

Na ⁺	141 mmol/l
Serum potassium	3.3 mmol/l
Urea	9.0 mmol/
Creatinine	188 μmol/l
Serum bicarbonate	8 mmol/l
Urine	pH 7.4

What is the most likely underlying cause of these results and possibly for some of her recent presentations?



- A

Bartter syndrome
- B

Type 1 (distal) renal tubular acidosis
- C

Type 2 (proximal) renal tubular acidosis
- D

Type 4 renal tubular acidosis
- E

Staghorn calculus leading to recurrent urinary sepsis

Explanation



- B

Type 1 (distal) renal tubular acidosis

This lady has a compensated metabolic acidosis with relatively high urinary pH and low potassium. These are features of type 1 (distal) renal tubular acidosis (RTA), and are probably secondary to her lupus-associated renal impairment. The suggestion of urinary calculi and recurrent urinary infections support this as these are recognised complications of the inability to acidify the urine. A staghorn calculus with recurrent sepsis would not in itself explain her biochemical picture, although nephrocalcinosis can also be associated with RTA. Nephrocalcinosis and recurrent urinary sepsis are also associated with type 1 (distal) RTA, and in the context of her rheumatological disease it is more likely that this is the underlying cause. It would also be highly unusual to find this condition coexisting with lupus renal disease.

- A

Bartter syndrome

Bartter syndrome is an inherited or sometimes sporadic disorder leading to hypokalaemia; however, this would not cause the urinary abnormalities seen here and, if inherited, would have presented earlier.

- C

Type 2 (proximal) renal tubular acidosis

Proximal renal tubular acidosis is due to impaired retention of bicarbonate in the proximal tubule leading to bicarbonate wasting and a systemic acidosis. It presents as a hyperchloraemic metabolic acidosis, usually with other features of proximal tubular dysfunction. Common causes include myeloma, amyloidosis, cystinosis and Wilson’s disease.

- D

Type 4 renal tubular acidosis

This is more common than type 1 and 2 renal tubular acidosis. It is due to hypoaldosteronism, usually hyporeninaemic hypoaldosteronism.

- E

Staghorn calculus leading to recurrent urinary sepsis

Staghorn calculus is caused by infection with *Proteus*, *Klebsiella* or *Serratia*. Typically, these stones are seen on KUB. It is not associated with SLE.

70778

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Difficult

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	84
Responses Total:	84
Responses - % Correct:	0%

Question 185 of 203

Investigations:

Urinalysis:

Two recent urine cultures have shown mixed growth on one and no growth on a second.

11

Submit

Skip Question

Back to Filters

Question 185 of 203

A 64-year-old man presents with a 2-month history of dysuria, urinary urgency and pelvic discomfort. He was investigated last year for proteinuria and haematuria picked up on urinalysis at the general practitioner's surgery, although no specific cause was found. Repeated urine cultures sent at this time showed no growth. He also reports weight loss over the last year of two stone.

Investigations:

Haemoglobin (Hb)	11.9 g/dl
White cell count (WCC)	9.2 × 10 ⁹ /l
Platelets	479 × 10 ⁹ /l
Mean corpuscular volume (MCV)	92 fl
Mean corpuscular haemoglobin content (MCHC)	32 g/dl
Na ⁺	139 mmol/l
K ⁺	3.8 mmol/l
Urea	6.9 mmol/l
Creatinine	101 μmol/l
Erythrocyte sedimentation rate (ESR)	73 mm/h
C-reactive protein	56 mg/l

Urinalysis:

Protein	1+
Blood	1+
Leucoctyes	3+
Glucose	Negative

Two recent urine cultures have shown mixed growth on one and no growth on a second.

What is the most likely diagnosis?



- A

Chronic/recurrent *E. coli* infection
- B

Chronic prostatitis
- C

Renal cell carcinoma
- D

Renal tract tuberculosis
- E

Squamous cell carcinoma of the bladder

Explanation



- D

Renal tract tuberculosis

This man has very long-standing urinalysis abnormalities with weight loss and symptoms of bladder irritation. In addition, he has evidence of an inflammatory response with a mild anaemia. These raise the suspicion of either malignant disease or severe chronic infective disease such as tuberculosis. The renal tract is a common site for tuberculous infection after the lungs, and is commonly involved in association with primary TB infection (albeit often without being detected clinically). It can be difficult to diagnose and requires repeated early morning urine cultures and occasionally more invasive steps such as urinary tract biopsy on cystoscopy.

- A

Chronic/recurrent *E. coli* infection

Occult bacterial infection is unlikely to be the cause in view of the repeated negative urinary cultures.

- B

Chronic prostatitis

Although prostatic symptoms are present, such as dysuria and urgency. Penile discharge may be present and we would expect the urinary growth to be positive for bacteria.

- C

Renal cell carcinoma

The symptoms of bladder irritation are unlikely to occur with renal cell carcinoma.

- E

Squamous cell carcinoma of the bladder

Although squamous cell carcinoma of the bladder is a possibility, this would be more likely to be a transitional cell lesion.

70779

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Difficult

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	85
Responses Total:	85
Responses - % Correct:	0%

Back to Filters

Question 186 of 203

A 22-year-old student presents with dark urine with the appearance of ‘black tea’, followed by swelling of the eyelids associated with malaise, anorexia and weakness. She has no relevant medical history, although she reports having had an infection at the site of a skin piercing approximately 4 weeks previously, for which she was prescribed some antibiotics. She took these for two days but stopped because the infection cleared and she developed some abdominal pain while taking them. On physical examination, her blood pressure is 125/70 mmHg, with chest and abdominal examination unremarkable.

Urinalysis:

Protein	3+
Blood	3+
Nitrites	Negative
Leucocytes	Negative

The likely diagnosis is?



- A

Acute intermittent porphyria
- B

Acute variegate porphyria
- C

Alkaptonuria
- D

Mesangioproliferative glomerulonephritis (immunoglobulin A [IgA] nephropathy)
- E

Post-streptococcal glomerulonephritis

70780

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 186 of 203

A 22-year-old student presents with dark urine with the appearance of ‘black tea’, followed by swelling of the eyelids associated with malaise, anorexia and weakness. She has no relevant medical history, although she reports having had an infection at the site of a skin piercing approximately 4 weeks previously, for which she was prescribed some antibiotics. She took these for two days but stopped because the infection cleared and she developed some abdominal pain while taking them. On physical examination, her blood pressure is 125/70 mmHg, with chest and abdominal examination unremarkable.

Urinalysis:

Protein	3+
Blood	3+
Nitrites	Negative
Leucocytes	Negative

The likely diagnosis is?



- A

Acute intermittent porphyria
- B

Acute variegate porphyria
- C

Alkaptonuria
- D

Mesangioproliferative glomerulonephritis (immunoglobulin A [IgA] nephropathy)
- E

Post-streptococcal glomerulonephritis

Explanation



- E

Post-streptococcal glomerulonephritis

This young woman has presented with a nephritic syndrome with frank (probably haemolysed) blood in her urine. The length of time after her skin infection makes the likely diagnosis a post-streptococcal glomerulonephritis (Option E).

- A

Acute intermittent porphyria

Acute intermittent porphyria can cause urinary discoloration when the urine is left to stand in sunlight, but this lady has dark urine when passed.

- B

Acute variegate porphyria

The clinical features in any case are not those of this form of porphyria, which typically presents with neuropathy, severe episodes of abdominal pain and neuropsychiatric disturbance. When comparing acute intermittent porphyria (AIP) and variegate porphyria, the latter rarely presents with acute episodes (unlike the former), and often the only reported symptom is excessive skin sensitivity to sunlight.

- C

Alkaptonuria

Alkaptonuria is an inborn error of the metabolism of tyrosine, resulting again in urine which turns black on standing due to excretion of homogentisic acid, which is oxidised on contact with air.

- D

Mesangioproliferative glomerulonephritis (immunoglobulin A [IgA] nephropathy)

Immunoglobulin A (IgA) nephropathy usually occurs with upper respiratory tract infections and usually either at the same time or within 2–3 days of these.

70780

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Difficult

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	86
Responses Total:	86
Responses - % Correct:	0%

Question 187 of 203

Urinalysis:

The likely diagnosis is?

11

Skip Question

Back to Filters

Question 187 of 203

A 28-year-old lawyer, originally from Singapore and who has recently returned from the Far East, presents complaining of passing bright red urine, facial puffiness and malaise. He also has a severe sore throat, which began two or three days before he noticed the red urine. His past medical history includes hay fever. On physical examination his blood pressure is 130/80 mmHg, and chest and abdominal examination is unremarkable.

Urinalysis:

Protein	2+
Blood	3+
Nitrites	Negative
Leucocytes	Negative

The likely diagnosis is?



- A

Focal segmental glomerulonephritis
- B

Mesangioproliferative glomerulonephritis (immunoglobulin A [IgA] nephropathy)
- C

Membranous glomerulonephritis
- D

Post-streptococcal glomerulonephritis
- E

Schistosomiasis

Explanation



- B

Mesangioproliferative glomerulonephritis (immunoglobulin A [IgA] nephropathy)

Mesangioproliferative glomerulonephritis (immunoglobulin A [IgA] nephropathy) is primarily a disease of young adults who develop ‘synpharyngitic’ episodes of glomerulonephritis, i.e. episodes which either coincide or occur within 2–3 days of an upper respiratory tract infection. It is the commonest glomerulonephritis (GN) in adults in the West. It is particularly common in the Far East, which is thought to be associated with a high incidence of the human leucocyte antigen (HLA) DQW7 haplotype in the local population.

- A

Focal segmental glomerulonephritis

We would expect significant proteinuria with focal segmental glomerulonephritis. The history of sore throat is not typical.

- C

Membranous glomerulonephritis

Membranous nephropathy is the most common cause of nephrotic syndrome (oedema, hypoalbuminaemia, nephrotic-range proteinuria) in adult patients. There is no haematuria, unlike in this scenario.

- D

Post-streptococcal glomerulonephritis

Post-streptococcal GN usually occurs several weeks after a streptococcal infection of either the upper respiratory tract or the skin.

- E

Schistosomiasis

Schistosomiasis, also known as bilharzia, is a disease caused by parasitic flatworms called schisotosomes. The urinary tract or the intestines may be infected. Signs and symptoms may include abdominal pain, diarrhoea, bloody stool or blood in the urine. Those who have been infected for a long time may experience liver damage, kidney failure, infertility or bladder cancer. In children, it may cause poor growth and learning difficulties.

70781

Rate this question:

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Difficult

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	87
Responses Total:	87
Responses - % Correct:	0%

Back to Filters

Question 188 of 203

A 40-year-old man with life-long episodes of fever and acute severe abdominal and chest pain presents to outpatients. He has lived most of his life in Turkey and several members of his family have suffered with similar symptoms, often being admitted as emergencies to local hospitals. He has been taking high doses of painkillers and anti-inflammatory medications for many years to help with his symptoms. On physical examination, his temperature is 38°C, blood pressure is 130/80 mmHg and chest and abdominal examination is unremarkable.

He was last seen over a year ago and a comparison of his blood results from that attendance and this shows:

Results now:

Na ⁺	138 mmol/l
K ⁺	4.6 mmol/l
Urea	12.4 mmol/l
Creatinine	180 μmol/l

Results 14 months ago:

Na ⁺	142 mmol/l
K ⁺	3.8 mmol/l
Urea	8.3 mmol/l
Creatinine	109 μmol/l

The single most useful test to confirm the pathological basis of his renal dysfunction would be?

- A

Intravenous urogram (IVU)
- B

MEFV gene analysis for familial Mediterranean fever (FMF)
- C

Rectal biopsy
- D

Renal biopsy
- E

Serum amyloid P component (SAP scan)

70782

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 188 of 203

A 40-year-old man with life-long episodes of fever and acute severe abdominal and chest pain presents to outpatients. He has lived most of his life in Turkey and several members of his family have suffered with similar symptoms, often being admitted as emergencies to local hospitals. He has been taking high doses of painkillers and anti-inflammatory medications for many years to help with his symptoms. On physical examination, his temperature is 38°C, blood pressure is 130/80 mmHg and chest and abdominal examination is unremarkable.

He was last seen over a year ago and a comparison of his blood results from that attendance and this shows:

Results now:

Na ⁺	138 mmol/l
K ⁺	4.6 mmol/l
Urea	12.4 mmol/l
Creatinine	180 μmol/l

Results 14 months ago:

Na ⁺	142 mmol/l
K ⁺	3.8 mmol/l
Urea	8.3 mmol/l
Creatinine	109 μmol/l

The single most useful test to confirm the pathological basis of his renal dysfunction would be?

- A

Intravenous urogram (IVU)
- B

MEFV gene analysis for familial Mediterranean fever (FMF)
- C

Rectal biopsy
- D

Renal biopsy
- E

Serum amyloid P component (SAP scan)

Explanation



- D

Renal biopsy

This man is likely to have familial Mediterranean FEVER (FMF), an inherited condition resulting in repeated episodes of polyserositis mimicking acute abdominal surgical presentations such as acute appendicitis, and often leading to unnecessary surgical exploration until the diagnosis is made. It is relatively common in the Middle East, especially in Turkey, Syria, Lebanon, Israel and Egypt. It relates to abnormal triggering of the inflammatory response and can lead to AA-type amyloid deposition and therefore renal failure. However, patients with this condition frequently take high doses of analgesics and anti-inflammatory drugs, which puts them at risk of analgesic nephropathy.

- A

Intravenous urogram (IVU)

IVU might show characteristic ‘cup and spill’ calices suggesting possible analgesic nephropathy, but this would not be confirmatory.

- B

MEFV gene analysis for familial Mediterranean fever (FMF)

Although gene testing might confirm the suspected diagnosis, it would not conclusively indicate the cause of the renal impairment.

- C

Rectal biopsy

Rectal biopsy followed by examination with polarised light leading to ‘apple-green’ birefringence might confirm amyloidosis, but this may not indicate the cause for the renal impairment if not identified.

- E

Serum amyloid P component (SAP scan)

Serum amyloid P component (SAP) scan may likewise confirm the presence of amyloid, but a renal biopsy would be diagnostic for either analgesic- or amyloid-associated nephropathy.

70782

Rate this question:

Next Question

Previous Question

Feedback

Tag Question

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	88
Responses Total:	88
Responses - % Correct:	0%

Back to Filters

Question 189 of 203

A 63-year-old man with known chronic renal failure (CRF) was admitted to another hospital 6 months ago with hypertension and congestive cardiac failure (CCF) where he was treated with furosemide (frusemide) 120 mg and enalapril 5 mg. His haemoglobin (Hb) concentration at that time was 11.0 g/dl and his creatinine level was 300 μmol/l. He has now presented to his regular hospital with lethargy. During the intervening period his GP has uptitrated his enalapril to 20 mg per day. His blood pressure is 115/70 mmHg, his jugular venous pressure (JVP) is visible and not raised. Chest is clear and abdomen is soft and non-tender. There is no peripheral oedema.

Investigations show:

Hb	8.0 g/dl
WCC	5.1 × 10 ⁹ /l
Na ⁺	140 mmol/l
K ⁺	5.3 mmol/l
Urea	15.0 mmol/l
Creatinine	380 μmol/l
Urine dipstick	Protein 3+, blood none

What would be the most appropriate management of this patient?

- A

Reduce enalapril
- B

Reduce furosemide (frusemide)
- C

Stop all drugs
- D

Stop enalapril
- E

Stop furosemide (frusemide)

70784

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 189 of 203

Investigations show:

Hb	8.0 g/dl
WCC	$5.1 \times 10^9/l$
Na ⁺	140 mmol/l
K ⁺	5.3 mmol/l
Urea	15.0 mmol/l
Creatinine	380 μmol/l
Urine dipstick	Protein 3+, blood none

What would be the most appropriate management of this patient?

- | | |
|---|-------------------------------|
| A | Reduce enalapril |
| B | Reduce furosemide (frusemide) |
| C | Stop all drugs |
| D | Stop enalapril |
| E | Stop furosemide (frusemide) |

Explanation



- | | |
|---|------------------|
| A | Reduce enalapril |
|---|------------------|

B	Reduce furosemide (frusemide)
---	-------------------------------

Reduction of the loop diuretic dose would contribute to the hyperkalaemia.

- | | |
|---|----------------|
| C | Stop all drugs |
|---|----------------|

D	Stop enalapril
---	----------------

E	Stop furosemide (frusemide)
---	-----------------------------

Temporary withdrawal of the loop diuretic may cause rebound pulmonary oedema.

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	89
Responses Total:	89
Responses - % Correct:	0%

Back to Filters

Question 190 of 203

A 24-year-old white man is referred to the Renal Clinic for further investigation of an abnormal urinalysis result, detected at a routine private health insurance medical some 3 months earlier. At this medical he was noted to have a BP of 159/89 mmHg. Urinalysis revealed 2+ blood and 1+ protein. The patient feels well and denies any significant past medical history. He is on no regular medication and denies any illicit drug use. As he was adopted, he is unable to say whether there is a family history of renal disease. On examination, he has a blood pressure of 154/96 mmHg. The rest of the clinical examination is unremarkable.

Laboratory and radiology results are as follows:

Urine microscopy	Red blood cells visualised
Na ⁺	140 mmol/l
K ⁺	3.7 mmol/l
Urea	5.3 mmol/l
Creatinine	95 μmol/l
Complement C3	108 mg/dl (65–190 mg/dl)
Complement C4	37 mg/dl (15–50 mg/dl)

Renal ultrasound scan:

Right kidney	10 cm, with two simple cysts
Left kidney	10 cm, normal morphology

What is the next most appropriate course of action?

-
- A

Check antiglomerular basement membrane (anti-GBM) antibodies
- B

Carry out genetic testing for adult polycystic kidney disease
- C

Check immunoglobulin levels
- D

Prescribe an antihypertensive drug
- E

Renal biopsy

70785

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 190 of 203

Laboratory and radiology results are as follows:

Renal ultrasound scan:

What is the next most appropriate course of action?

- ### Explanation

- The differential diagnosis includes IgA nephropathy and thin basement membrane disease. Although the gold standard diagnostic test is a renal biopsy it would not help management at present. A renal biopsy confers risk, and therefore there is no point in carrying out a potentially kidney-threatening procedure for little gain. The patient's hypertension does need treatment, however, so he should be commenced on an antihypertensive drug. He should be monitored periodically in the renal clinic for any progression of proteinuria and/or a fall in renal function.

- The patient is too well to have Goodpasture's syndrome, so it would not be helpful to check anti-GBM antibodies.

- The two simple cysts in the normal-sized right kidney are incidental and are of no clinical significance. Therefore, this patient does not need to be screened for adult polycystic kidney disease.

- The differential diagnosis includes IgA nephropathy and thin basement membrane disease. If the patient has IgA disease, his serum IgA levels may well be abnormal, but checking the levels would not constitute a diagnostic test.

- The gold standard way to distinguish between the two main differential diagnoses in this case, which is IgA nephropathy and thin membrane disease, is to carry out a renal biopsy, but at this stage it would not impact on management (although it is something that should be considered at a later date).

Rate this question:

Next Question

[Previous Question](#)

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

[Blog](#) [About Pastest](#) [Contact Us](#) [Help](#)

Back to Filters

Question 191 of 203

A 78-year old white man presents to the Emergency Department with a 1-day history of haemoptysis and a 2-week history of general malaise, shortness of breath, cough and intermittent fevers. He was prescribed clarithromycin for a presumed chest infection 10 days ago, but he has failed to respond to treatment. On examination, he has a temperature of 37.8°C; he is tachypnoeic and has evidence of bilateral lung consolidation.

Investigations show:

Na ⁺	143 mmol/l
K ⁺	5.5 mmol/l
Serum bicarbonate	19 mmol/l
Urea	18 mmol/l
Creatinine	220 μmol/l
CRP	120 mg/l
ESR	90 mm/first hour
Hb	10.5 g/dl
WCC	16 × 10 ⁹ /l
Complement C3	73 mg/dl (65-190 mg/dl)
Antibodies to nuclear antigens (ANA)	Positive
Antineutrophil cytoplasmic antigens (ANCA)	Positive
Antiproteinase 3	Positive
Antiglomerular basement membrane antibodies	Negative
Urine dipstick	3+ blood and 2+ protein
Urine microscopy	Red cell casts
Chest X-ray	Bilateral infiltrates

What is the most appropriate treatment for this patient?

- A

High-dose prednisolone alone
- B

High-dose prednisolone and cyclophosphamide
- C

Immunoglobulin IV
- D

Plasmapheresis
- E

Supportive care and IV antibiotics

70786

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 191 of 203

A 78-year old white man presents to the Emergency Department with a 1-day history of haemoptysis and a 2-week history of general malaise, shortness of breath, cough and intermittent fevers. He was prescribed clarithromycin for a presumed chest infection 10 days ago, but he has failed to respond to treatment. On examination, he has a temperature of 37.8°C; he is tachypnoeic and has evidence of bilateral lung consolidation.

Investigations show:

Na ⁺	143 mmol/l
K ⁺	5.5 mmol/l
Serum bicarbonate	19 mmol/l
Urea	18 mmol/l
Creatinine	220 μmol/l
CRP	120 mg/l
ESR	90 mm/first hour
Hb	10.5 g/dl
WCC	16 × 10 ⁹ /l
Complement C3	73 mg/dl (65-190 mg/dl)
Antibodies to nuclear antigens (ANA)	Positive
Antineutrophil cytoplasmic antigens (ANCA)	Positive
Antiproteinase 3	Positive
Antiglomerular basement membrane antibodies	Negative
Urine dipstick	3+ blood and 2+ protein
Urine microscopy	Red cell casts
Chest X-ray	Bilateral infiltrates

What is the most appropriate treatment for this patient?

- A

High-dose prednisolone alone
- B

High-dose prednisolone and cyclophosphamide
- C

Immunoglobulin IV
- D

Plasmapheresis
- E

Supportive care and IV antibiotics

Explanation



- B

High-dose prednisolone and cyclophosphamide

The patient has evidence of a rapidly progressive glomerulonephritis. The differential diagnosis includes pauci-immune small vessel vasculitides (Wegener’s granulomatosis, polyarteritis nodosa (PAN) and microscopic polyangiitis), Goodpasture’s syndrome, systemic lupus erythematosus (SLE) and IgA nephropathy. The presence of a pulmonary-renal syndrome coupled with the pattern of autoantibodies points to a diagnosis of Wegener’s granulomatosis. The most appropriate treatment for this is high-dose steroids with cyclophosphamide, as this can induce remission in 80-90% of patients.

- A

High-dose prednisolone alone

High-dose prednisolone on its own would not be enough immunosuppression for rapidly progressive glomerulonephritis.

- C

Immunoglobulin IV

There is no evidence that there is any benefit for IV immunoglobulins in this scenario. In fact, it may be important to *remove* the antigenic immunoglobulins for treatment, which is the concept for plasmapheresis.

- D

Plasmapheresis

When there is pulmonary haemorrhage associated with Wegener’s granulomatosis then plasmapheresis would be indicated.

- E

Supportive care and IV antibiotics

It is important to stop progression of the inflammation and damage; therefore supportive care will not be the appropriate treatment. Appropriate immunosuppression in the form of prednisolone and cyclophosphamide is crucial.

70786

Rate this question: ⚙️☆☆☆☆☆

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	91
Responses Total:	91
Responses - % Correct:	0%

Question 192 of 203

Her blood results are as follows:

She has had a test of her renin and aldosterone activity both of which were high.

3

Skip Question

Back to Filters

Question 192 of 203

A 45-year-old woman presents with refractory hypertension first picked up on a routine health check. She is a long-term smoker. She has mild asthma and eczema but no other significant medical history. She has recently complained to her general practitioner of pains in her calves which have occurred while she is walking her dog. Her blood pressure remains 150/95 mmHg in spite of multiple medical therapy trials.

Her blood results are as follows:

Na ⁺	144 mmol/l
K ⁺	3.2 mmol/l
Urea	4.8 mmol/l
Creatinine	76 μmol/l

She has had a test of her renin and aldosterone activity both of which were high.

The most likely diagnosis is:

- A

Bilateral adrenal hyperplasia
- B

Cushing’s disease
- C

Primary hyperaldosteronism (Conn’s syndrome)
- D

Renovascular hypertension secondary to fibromuscular dysplasia
- E

Renovascular hypertension secondary to renal arterial sclerosis

Explanation



- E

Renovascular hypertension secondary to renal arterial sclerosis

This woman has evidence of secondary hyperaldosteronism with high normal sodium and low potassium levels in the presence of high renin and aldosterone activity. This is likely to be a result of renal artery stenosis from atherosclerosis. She has a risk factor for arteriosclerosis and has symptoms that could reflect peripheral vascular disease which is recognised to occur with atherosclerotic renal artery sclerosis. Limb arteries can be affected by fibromuscular dysplasia, but this occurs only in about 5% of cases.

- A

Bilateral adrenal hyperplasia

Adrenal hyperplasia may lead to increased secretion of aldosterone. This is primary hyperaldosteronism, and therefore the renin would be suppressed.

- B

Cushing’s disease

Potassium is normal in Cushing’s disease because the underlying disease is due to an excess of cortisol production.

- C

Primary hyperaldosteronism (Conn’s syndrome)

In Conn’s syndrome the renin activity would be suppressed.

- D

Renovascular hypertension secondary to fibromuscular dysplasia

Her age makes fibromuscular dysplasia less likely as it usually presents between the ages of 30 and 40 years. In addition, arteriosclerosis is a more common cause of renal artery stenosis than fibromuscular dysplasia.

70787

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	92
Responses Total:	92
Responses - % Correct:	0%

Back to Filters

Question 193 of 203

A 45-year-old man who is a known alcohol abuser is under investigation for hyponatraemia. He drinks several lagers daily, he is otherwise well and not on any medication. He smokes 15 cigarettes/day. On physical examination his blood pressure is 130/80 mmHg, chest and abdominal examination is unremarkable.

Investigations reveal:

Chest X-ray	Normal
Computed tomography (CT) of the head	Normal
Thyroid function	Normal
Na ⁺	125 mmol/l
K ⁺	4.5 mmol/l
Urea	3.5 mmol/l
Creatinine	110 μmol/l
Glucose	4.5 mmol/l
Plasma osmolality	285 mosmol/kg
Calculated osmolality	269 mosmol/kg

What is the cause of the hyponatraemia?

- A

Addison’s disease
- B

Beer potomania
- C

Hypothyroidism
- D

Salt-losing nephropathy
- E

Syndrome of inappropriate antidiuretic hormone secretion (SIADH)

70788

Submit

Previous QuestionSkip Question

Calculator

Normal Values

Back to Filters

Question 193 of 203

A 45-year-old man who is a known alcohol abuser is under investigation for hyponatraemia. He drinks several lagers daily, he is otherwise well and not on any medication. He smokes 15 cigarettes/day. On physical examination his blood pressure is 130/80 mmHg, chest and abdominal examination is unremarkable.

Investigations reveal:

Chest X-ray	Normal
Computed tomography (CT) of the head	Normal
Thyroid function	Normal
Na ⁺	125 mmol/l
K ⁺	4.5 mmol/l
Urea	3.5 mmol/l
Creatinine	110 μmol/l
Glucose	4.5 mmol/l
Plasma osmolality	285 mosmol/kg
Calculated osmolality	269 mosmol/kg

What is the cause of the hyponatraemia?



- A

Addison's disease
- B

Beer potomania
- C

Hypothyroidism
- D

Salt-losing nephropathy
- E

Syndrome of inappropriate antidiuretic hormone secretion (SIADH)

Explanation



- B

Beer potomania

Beer potomania is a recognised cause of hyponatraemia in alcohol misusers. The electrolyte imbalance normally corrects itself with cessation of alcohol. The osmolar gap is due to the presence of osmotically active ethanol in the blood.

- A

Addison's disease

Addison's disease is due to deficiency in corticosteroids. Therefore, we would expect a high potassium and a postural drop in blood pressure or low blood pressure.

- C

Hypothyroidism

The osmolar gap would be normal. There may be associated symptoms such as weight gain, low mood, reduced appetite and intolerance to the cold.

- D

Salt-losing nephropathy

There may be variety of causes of salt-losing nephropathy including urea-induced osmotic diuresis, glomerulotubular imbalance and impaired tubular salt resorption. The osmolar gap would be normal.

- E

Syndrome of inappropriate antidiuretic hormone secretion (SIADH)

SIADH is a diagnosis of exclusion. It is defined by hyponatraemia and hypo-osmolality resulting from inappropriate and continued secretion of ADH. Patients would be euvolaemic and the osmolar gap would be normal.

70788

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Difficult

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	93
Responses Total:	93
Responses - % Correct:	0%

Back to Filters

Question 194 of 203

A 52-year-old man presents with several months’ history of generalised swelling, fatigue, cough, dyspnoea and several episodes of haemoptysis. There is no significant past medical history and he did not take any regular medication. He smokes 20 cigarettes per day and drinks 14 units of alcohol per week. On examination, he is grossly oedematous and has ascites. Cardiorespiratory examination is unremarkable and there are no neurological signs or rashes.

Investigation results are below:

Hb	10.2 g/dl
WCC	6.0 × 10 ⁹ /l
PLT	380 × 10 ⁹ /l
Mean corpuscular volume (MCV)	90fl
Na ⁺	145 mmol/l
K ⁺	3.7 mmol/l
Urea	8.2 mmol/l
Bilirubin	16 μmol/l
Creatinine	180 μmol/l
Albumin	22 g/l
Aspartate transaminase (AST)	32 U/l
Alkaline phosphatase	120 U/l
Urinalysis	Protein +++
24-hour urinary protein excretion	5g
Chest radiograph	Enlarged right hilum
Echocardiogram	Mild left ventricular impairment, no valve lesion
Abdominal ultrasound scan	Normal-sized kidneys, no abnormality seen

A renal biopsy was performed.

What is it most likely to show?

-
- A

Crescent formation
- B

Kimmelstiel–Wilson lesions
- C

Fusion of podocyte foot processes on electron microscopy
- D

Necrotising granulomata
- E

Thickened glomerular basement membrane with deposits of IgG and C3

70792

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 194 of 203

Hb	10
----	----

A renal biopsy was performed.

what is it most likely to show?

- 「**Thinking**」

- membranous glomerulonephritis which is associated with an underlying bronchial carcinoma. Membranous glomerulonephritis accounts for 20–30% of adult nephrotic syndromes, and there will be underlying malignancy in 10% of cases. Other causes of membranous nephropathy are drugs (eg gold and penicillamine), autoimmune diseases (eg systemic lupus erythematosus (SLE)) and infections (eg hepatitis). Approximately one-third of patients will progress to end-stage renal failure within 10–20 years of diagnosis. Typical renal biopsy features of membranous nephropathy is thickened glomerular basement membrane with deposits of IgG and C3. Treatment may include chlorambucil and cyclophosphamide.

- presentation would be a quickly deteriorating renal function with blood and protein in the urine. Crescentic glomerulonephritis may be found in anti-glomerular basement membrane disease and lupus nephritis.

- scenario there is no mention of history of diabetes and there is no sugar in the urine. The first sign of diabetic nephropathy is microalbuminuria.

- This feature is found in minimal change disease. Although minimal change disease can present with nephrotic syndrome, it is more common in young children. The prognosis is generally good and steroid responsive.

- Granulomas are an organised collection of macrophages. There are a variety of conditions which can cause necrotising granulomata, of which the most classic is tuberculosis. In this scenario there is no history that is indicative of tuberculosis such as fever, weight loss or any contact history.

100% 99% 98% 97% 96% 95% 94% 93% 92% 91% 90% 89% 88% 87% 86% 85% 84% 83% 82% 81% 80% 79% 78% 77% 76% 75% 74% 73% 72% 71% 70% 69% 68% 67% 66% 65% 64% 63% 62% 61% 60% 59% 58% 57% 56% 55% 54% 53% 52% 51% 50% 49% 48% 47% 46% 45% 44% 43% 42% 41% 40% 39% 38% 37% 36% 35% 34% 33% 32% 31% 30% 29% 28% 27% 26% 25% 24% 23% 22% 21% 20% 19% 18% 17% 16% 15% 14% 13% 12% 11% 10% 9% 8% 7% 6% 5% 4% 3% 2% 1% 0%

End Session

Peer Responses %

Responses - % Correct.

Question 195 of 203

Investigation results are below:

What is the correct way to treat her condition?

- A Admission and cautious oral angiotensin-converting enzyme (ACE) inhibitor introduction
- B IV furosemide
- C IV labetalol
- D IV sodium nitroprusside
- E Immediate haemodialysis with ultrafiltration

Submit

Skip Question

Back to Filters

Question 195 of 203

A 34-year-old woman is admitted as an emergency with a 24-hour history of severe headaches, blurred vision and shortness of breath. She had been previously fit and well with no significant past medical history and apart from some recent fatigue, had no other symptoms. She smoked 10 cigarettes per day and drank very little alcohol. She was not sure about family history as she had been adopted. On examination she was fully alert and oriented with warm peripheries and a blood pressure of 185/115 mmHg. She had bi-basal chest crepitations, bilateral irregular masses palpable in both flanks and hepatomegaly. Fundoscopy revealed flame haemorrhages and exudates with indistinct disc margins. There were no other focal neurological signs.

Investigation results are below:

Hb	10.2 g/dl
WCC	4.4 x 10 ⁹ /l
PLT	160 x 10 ⁹ /l
Mean corpuscular volume (MCV)	92 fl
Na ⁺	143 mmol/l
K ⁺	4.6 mmol/l
Urea	17 mmol/l
Creatinine	250 µmol/l
Bilirubin	15 µmol/l
Alkaline phosphatase	92 U/l
Aspartate transaminase (AST)	32 U/l
Gamma-glutamyl transferase (GGT)	42 U/l
Urinalysis	Protein ++ blood++
Electrocardiogram (ECG)	Sinus rhythm, left ventricular hypertrophy with strain pattern
Chest X-ray	Cardiomegaly, upper lobe diversion, alveolar shadowing

What is the correct way to treat her condition?



- A

Admission and cautious oral angiotensin-converting enzyme (ACE) inhibitor introduction
- B

IV furosemide
- C

IV labetalol
- D

IV sodium nitroprusside
- E

Immediate haemodialysis with ultrafiltration

Explanation



- A

Admission and cautious oral angiotensin-converting enzyme (ACE) inhibitor introduction

This patient has malignant hypertension secondary to previously undiagnosed polycystic kidney disease. This causes vascular fibrinoid necrosis and is usually the first manifestation of hypertension, but it can sometimes occur in known hypertensive individuals. Death mainly occurs from heart failure, renal failure or stroke, with an untreated mortality of approximately 80%. Management with bed rest and a gradual reduction of the blood pressure over 24–48 h with oral antihypertensives is usually sufficient (aim for a diastolic pressure of 100–110 mmHg). If the blood pressure is reduced too rapidly it may cause cerebral, renal, retinal or myocardial infarction. Parenteral therapy should be reserved for cases when very rapid blood pressure reduction is required, such as hypertensive encephalopathy or aortic dissection. The agents of choice are usually labetalol or sodium nitroprusside. Patients with malignant hypertension and a serum creatinine of >300 µmol/l often progress to end-stage renal failure.

- B

IV furosemide

Furosemide will help the pulmonary oedema in this case, but it is important to first control the blood pressure.

- C

IV labetalol

Labetalol is an appropriate antihypertensive to use; however, it should not to be given intravenously in this situation. This is because it is important not to drop the blood pressure too rapidly as it could lead to a stroke.

- D

IV sodium nitroprusside

Cautious control of the blood pressure is needed in this scenario and it should be administered in the simplest method. Therefore IV sodium nitroprusside is not needed here. In addition, there are some potential side-effects of sodium nitroprusside, including skin rashes, nausea, warmth and redness under the skin or darkening of the veins under the skin.

- E

Immediate haemodialysis with ultrafiltration

Although this woman has pulmonary oedema and signs of fluid overload, her renal impairment is not severe and there are no immediate indications for acute dialysis. Medical therapy should be tried first – to control blood pressure. If she is anuric and medical therapy fails to improve symptoms then dialysis may then be indicated. Absolute indications for acute dialysis include hyperkalaemia (≥6.5 mmol/l) or rapidly rising potassium that is resistant to medical treatment and pulmonary oedema with inadequate response to diuretics particularly in an oligo-anuric patient.

70793

Rate this question: ⓪☆☆☆☆☆

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Difficult

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	95
Responses Total:	95
Responses - % Correct:	0%

Back to Filters

Question 196 of 203

A 46-year-old man was admitted with 24 hours of severe pleuritic, retrosternal chest pain. He described a preceding 6-week history of malaise, fever, myalgia and 1 stone of weight loss. Over the last few days he had also developed a rash and had noticed some loss of sensation in his feet and ankles which was causing some difficulty with walking. On examination his temperature was 39°C, blood pressure was 190/108 mmHg, pulse 110 bpm regular. There was a friction rub audible at the left sternal edge. His chest was clear and his abdomen was soft and non-tender. He had paraesthesia in a stocking distribution bilaterally and purpuric skin lesions.

Investigation results are below:

Hb	8.6 g/dl
WCC	18 × 10 ⁹ /l
PLT	450 × 10 ⁹ /l
Mean corpuscular volume (MCV)	96 fl
Erythrocyte sedimentation rate (ESR)	102 mm/h
Na ⁺	132 mmol/l
K ⁺	5.8 mmol/l
Urea	26.0 mmol/l
Creatinine	860 μmol/l
Bilirubin	16 μmol/l
Aspartate transaminase (AST)	30 U/l
Alkaline phosphatase	80 U/l
Gamma-glutamyl transferase (GGT)	48 U/l
Urinalysis	Protein ++ blood +++
Electrocardiogram (ECG)	Saddle-shaped ST segments across all leads
Chest radiograph	Normal
Serum pANCA and cANCA (antineutrophil cytoplasmic antibody)	Negative

What is the most likely diagnosis?

- A

Cryoglobulinaemia
- B

Goodpasture’s syndrome
- C

Microscopic polyangitis
- D

Polyarteritis nodosa (PAN)
- E

Wegener’s granulomatosis

70794

Submit

Previous Question

Skip Question

Calculator

✔

Normal Values

✔

Back to Filters

Question 196 of 203

A 46-year-old man was admitted with 24 hours of severe pleuritic, retrosternal chest pain. He described a preceding 6-week history of malaise, fever, myalgia and 1 stone of weight loss. Over the last few days he had also developed a rash and had noticed some loss of sensation in his feet and ankles which was causing some difficulty with walking. On examination his temperature was 39°C, blood pressure was 190/108 mmHg, pulse 110 bpm regular. There was a friction rub audible at the left sternal edge. His chest was clear and his abdomen was soft and non-tender. He had paraesthesia in a stocking distribution bilaterally and purpuric skin lesions.

Investigation results are below:

Hb	8.6 g/dl
WCC	18 × 10 ⁹ /l
PLT	450 × 10 ⁹ /l
Mean corpuscular volume (MCV)	96 fl
Erythrocyte sedimentation rate (ESR)	102 mm/h
Na ⁺	132 mmol/l
K ⁺	5.8 mmol/l
Urea	26.0 mmol/l
Creatinine	860 μmol/l
Bilirubin	16 μmol/l
Aspartate transaminase (AST)	30 U/l
Alkaline phosphatase	80 U/l
Gamma-glutamyl transferase (GGT)	48 U/l
Urinalysis	Protein ++ blood +++
Electrocardiogram (ECG)	Saddle-shaped ST segments across all leads
Chest radiograph	Normal
Serum pANCA and cANCA (antineutrophil cytoplasmic antibody)	Negative

What is the most likely diagnosis?

- A

Cryoglobulinaemia
- B

Goodpasture’s syndrome
- C

Microscopic polyangitis
- D

Polyarteritis nodosa (PAN)
- E

Wegener’s granulomatosis

Explanation

- D

Polyarteritis nodosa (PAN)

PAN is a necrotising vasculitis which causes aneurysms of medium-sized arteries with subsequent thrombosis and infarction. It occurs most commonly in middle-aged men. It is a multisystem disorder, but the most common causes of death are from renal (hypertension, renal failure) and cardiac (coronary arteritis, myocardial infarction, pericarditis) involvement. Other features include mononeuritis multiplex, sensorimotor neuropathy, abdominal pain from infarcted viscera and skin lesions (purpura, livedo reticularis). Diagnosis is made by a combination of clinical features plus renal or mesenteric angiography which will demonstrate microaneurysms. ANCA is very rarely positive in classic PAN. Treatment options include with high-dose corticosteroids, azathioprine and cyclophosphamide.

- A

Cryoglobulinaemia

Cryoglobulinaemia is associated with a purpuric rash on the legs (leucocytoclastic vasculitis on biopsy); it may be associated with infections such as hepatitis C. Symptoms include fatigue, weakness, weight loss, arthralgia, myalgia and mononeuritis multiplex. Other symptoms include hepatosplenomegaly and Raynaud’s phenomenon. Renal manifestations include blood and protein in the urine. The proteinuria is frequently in the nephrotic range. There would be hypertension, renal impairment that is progressive or even present as an acute renal failure.

- B

Goodpasture’s syndrome

Goodpasture’s syndrome, also known as anti-GBM disease, is associated with pulmonary haemorrhage. There is pathogenic IgG anti-GBM antibodies against the α3 chain of the type IV collagen in the basement membrane of the glomerulus and alveolus. It may present gradually over months or years, or abruptly over days with fulminant pulmonary haemorrhage and renal failure.

- C

Microscopic polyangitis

Microscopic polyangitis is less likely given that ANCA is negative, because it is a small-vessel vasculitis characterised by ANCA. Symptoms include weight loss, fever, malaise and flitting polyarthralgia. There is no ear, nose and throat involvement. P-ANCA (MPO) is positive with no granulomata in the kidneys. There is a low likelihood of relapse.

- E

Wegener’s granulomatosis

Wegener’s is less likely given that ANCA is negative because, similar to microscopic polyangitis, it is a small-vessel vasculitis characterised by ANCA. However, in contrast to microscopic polyangitis, there are frequently ear, nose and throat symptoms. C-ANCA (PR3) is positive with granulomata in the kidneys. There is a high likelihood of relapse.

70794

Rate this question: ⚙️ ⭐️ ⭐️ ⭐️ ⭐️ ⭐️

Next Question

Previous Question

Tag Question

Feedback

End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	96
Responses Total:	96
Responses - % Correct:	0%

Back to Filters

Question 197 of 203

A 58-year-old man presents with haematuria and weight loss. He has a past medical history of hypertension. On physical examination his blood pressure is 160/70 mmHg, chest and abdominal examination was unremarkable. His renal function is normal.

His computed tomography (CT) scan is below:



CT abdomen confirms the presence of a renal carcinoma. CT thorax and additional PET scanning does not suggest other metastases apart from the one shown.

What is the most appropriate management strategy?

- | | |
|---|--|
| A | Combined nephrectomy and resection of brain lesion |
| B | Immunotherapy only |
| C | Nephrectomy and cranial irradiation |
| D | Nephrectomy and small molecule kinase inhibitor |
| E | Palliative chemotherapy only |

70795


Submit

Previous QuestionSkip Question

Calculator✔

Normal Values✔

Question 197 of 203



A	Combined nephrectomy and resection of brain lesion
B	Immunotherapy only
C	Nephrectomy and cranial irradiation
D	Nephrectomy and small molecule kinase inhibitor
E	Palliative chemotherapy only

D	Nephrectomy and small molecule kinase inhibitor
---	---

Costa et al. 1997.

A	Combined nephrectomy and resection of brain lesion
---	--

Immunotherapy with interleukin-2 or interferon- α can produce a transient increase in the CD8⁺ frequency but not in the CD4⁺ frequency.

C Nephrectomy and cranial irradiation

Radiotherapy and chemotherapy have no proven benefit in renal cell carcinoma. Renal cell carcinoma is resistant in most cases with

E	Palliative chemotherapy only
---	------------------------------

(1) Costa Rican development

Rate this question:

[Previous Question](#)
[Flag Question](#)
[Feedback](#)
[End Session](#)

Responses Correct:	0
Responses Incorrect:	97
Responses Total:	97
Responses - % Correct:	0%

Question 198 of 203

Investigation results are below:

Hb	14 g/dl
WCC	$6.0 \times 10^9/l$
PLT	$200 \times 10^9/l$
Mean corpuscular volume (MCV)	96 fl
Na ⁺	142 mmol/l
K ⁺	2.9 mmol/l
Urea	5.2 mmol/l
Creatinine	120 μ mol/l
Bicarbonate	16 mmol/l
Chloride	120 mmol/l
Phosphate	0.7 mmol/l (0.8-1.4)
Urate	0.32 mmol/l

pH	7.210
$p_a(\text{O}_2)$	12.6 kPa
$p_a(\text{CO}_2)$	4.8 kPa
Urinalysis	pH 6.3, blood ++
Abdominal radiograph	Speckled calcification in the left and right upper quadrants

11

- | | |
|---|--|
| A | Fanconi syndrome |
| B | Medullary sponge kidney |
| C | Type 1 (distal) renal tubular acidosis |
| D | Type 2 (proximal) renal tubular acidosis |
| E | Type 4 renal tubular acidosis |

Submit

Skip Question

Back to Filters

Question 198 of 203

A 23-year-old man was admitted to hospital with severe left loin pain and vomiting. Prior to this he described a 5-week history of worsening generalised weakness and anorexia. He was previously well and on no regular medication. On examination he was in obvious discomfort and hyperventilating. There was tenderness and guarding in the left renal angle, but no other significant findings.

Investigation results are below:

Hb	14 g/dl
WCC	6.0 × 10 ⁹ /l
PLT	200 × 10 ⁹ /l
Mean corpuscular volume (MCV)	96 fl
Na ⁺	142 mmol/l
K ⁺	2.9 mmol/l
Urea	5.2 mmol/l
Creatinine	120 μmol/l
Bicarbonate	16 mmol/l
Chloride	120 mmol/l
Phosphate	0.7 mmol/l (0.8-1.4)
Urate	0.32 mmol/l

Arterial blood gases (on air):

pH	7.210
<i>P_a</i> (O ₂)	12.6 kPa
<i>P_a</i> (CO ₂)	4.8 kPa
Urinalysis	pH 6.3, blood ++
Abdominal radiograph	Speckled calcification in the left and right upper quadrants

What is the most likely diagnosis?

- A

Fanconi syndrome
- B

Medullary sponge kidney
- C

Type 1 (distal) renal tubular acidosis
- D

Type 2 (proximal) renal tubular acidosis
- E

Type 4 renal tubular acidosis

Explanation

- C

Type 1 (distal) renal tubular acidosis

The kidneys are unable to create an acid urine due to failure of hydrogen ion excretion in the distal tubule. Urine pH will be greater than 5.5, often in the presence of a severe systemic acidosis. There is almost invariably hypokalaemia, which can cause severe weakness. Hypercalciuria and hyperphosphaturia occur due to the release of calcium phosphate from bone in order to buffer excess hydrogen ions during acidosis, and the direct effects of acidosis on tubular reabsorption of these ions. Other clinical features include bone pain due to osteomalacia, renal stones, constipation, fatigue and anorexia. Nephrocalcinosis may be seen. Renal function is usually preserved. In children it commonly presents with polyuria, thirst and failure to thrive. Biochemically, there is a hyperchloraemic metabolic acidosis with a normal anion gap. Causes of type 1 renal tubular acidosis include autoimmune diseases (eg systemic lupus erythematosus), drugs (eg amphotecerin B), or it may be inherited or idiopathic. Treatment is with bicarbonate replacement (1–3 mmol/kg/day); however hypokalaemia must be corrected before the acidosis in order to prevent a further fall in the potassium and the risk of cardiac arrest. Nephrocalcinosis is not a feature of proximal renal tubular acidosis.

- A

Fanconi syndrome

Fanconi syndrome is a generalised descriptive term for proximal tubular dysfunction. There is a failure of the proximal tubules to reabsorb many filtered substances and classically there is phosphate wasting. There is bone pain and osteomalacia. The causes are as for proximal renal tubular acidosis (see below). Investigations would show a metabolic acidosis, reduced serum phosphate, reduced serum urate, glycosuria, proteinuria and increased urinary phosphate and citrate.

- B

Medullary sponge kidney

Medullary sponge kidney is a benign common condition characterised by diffuse medullary cystic formation. Impaired calcium handling leads to formation of calcium-containing stones. There are no other biochemical abnormalities such as acidosis seen in this case.

- D

Type 2 (proximal) renal tubular acidosis

There is impaired retention of bicarbonate in the proximal tubule leading to bicarbonate wasting and systemic acidosis. There is hyperchloraemic metabolic acidosis. Common causes include myeloma, amyloidosis, cystinosis, Wilson’s disease and heavy metal toxicity.

- E

Type 4 renal tubular acidosis

This is known as hyporeninaemic hypoaldosteronism. Aldosterone promotes urinary potassium loss so its absence leads to hyperkalaemia. Investigations would show hyperkalaemia, low bicarbonate (rarely less than 16 mmol/l), normal anion gap with increased chloride, negative urinary anion gap, reduced urinary citrate. Urine pH is variable but often less than 5.3.

70796

Rate this question: ⚙️☆☆☆☆☆

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Difficult

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	98
Responses Total:	98
Responses - % Correct:	0%

Back to Filters

Question 199 of 203

A 38-year-old Asian woman presents with several weeks of increased urinary frequency and dysuria. She had already been treated with two courses of antibiotics from her general practitioner, which had not helped. She had been feeling unusually lethargic recently and had lost a small amount of weight, but had no other specific symptoms. On examination she was thin with a low grade fever of 37.6°C. Her abdomen was soft with mild suprapubic tenderness but no palpable masses. The rest of the clinical examination was normal.

Investigation results are below:

Hb	10.2 g/dl
WCC	10.8 × 10 ⁹ /l
Platelets	390 × 10 ⁹ /l
Mean corpuscular volume (MCV)	82 fl
Erythrocyte sedimentation rate (ESR)	30mm/h
Na ⁺	138 mmol/l
Ca ²⁺	2.52 mmol/l
K ⁺	3.6 mmol/l
Urea	5.2 mmol/l
Creatinine	98 μmol/l
Urinalysis	Blood + protein +

Urine microscopy:

WCC	22/mm ³
Red blood count	3/mm ³
Few hyaline casts	
No organisms seen	
Urine culture	No growth after 5 days

What is the next most appropriate investigation?

- A

Micturating cystourethrography
- B

Renal biopsy
- C

Plain abdominal radiograph
- D

Urine electrophoresis
- E

Urine for mycobacterial culture

70797

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 199 of 203

A 38-year-old Asian woman presents with several weeks of increased urinary frequency and dysuria. She had already been treated with two courses of antibiotics from her general practitioner, which had not helped. She had been feeling unusually lethargic recently and had lost a small amount of weight, but had no other specific symptoms. On examination she was thin with a low grade fever of 37.6°C. Her abdomen was soft with mild suprapubic tenderness but no palpable masses. The rest of the clinical examination was normal.

Investigation results are below:

Hb	10.2 g/dl
WCC	10.8 × 10 ⁹ /l
Platelets	390 × 10 ⁹ /l
Mean corpuscular volume (MCV)	82 fl
Erythrocyte sedimentation rate (ESR)	30mm/h
Na ⁺	138 mmol/l
Ca ²⁺	2.52 mmol/l
K ⁺	3.6 mmol/l
Urea	5.2 mmol/l
Creatinine	98 µmol/l
Urinalysis	Blood + protein +

Urine microscopy:

WCC	22/mm ³
Red blood count	3/mm ³
Few hyaline casts	
No organisms seen	
Urine culture	No growth after 5 days

What is the next most appropriate investigation?

- A

Micturating cystourethrography
- B

Renal biopsy
- C

Plain abdominal radiograph
- D

Urine electrophoresis
- E

Urine for mycobacterial culture

Explanation



- E

Urine for mycobacterial culture

This woman has a sterile pyuria which could be due to a partially treated bacterial urinary tract infection; however genitourinary tuberculosis has to be excluded, especially in the context of malaise and weight loss and the patient's ethnic origin. Genitourinary TB develops in approximately 5% of cases of pulmonary TB and is usually due to haematogenous spread to the renal cortex during the primary phase of infection. The cortical lesion may then ulcerate into the pelvis, ultimately involving the bladder, seminal vesicles and prostate. It tends to present between 20 and 40 years of age. Other clinical features include haematuria, urethral strictures, cold abscesses and chronic epididymo-orchitis. Renal failure may occur due to extensive destruction of the kidneys or by obstruction secondary to fibrosis. The diagnosis can be made with intravenous urography, ultrasound or computed tomography (CT) scan, in combination with several early morning urine samples for mycobacterial culture, in order to demonstrate active infection. The treatment is the same as for pulmonary TB.

- A

Micturating cystourethrography

This investigation visualises the urinary tract from bladder to urethra as the patient urinates. It is used for diagnosis of vesicoureteral reflux (VUR) where there is reflux of urine from the bladder up the ureters back into the kidneys. Radiologists may grade the degree/staging of VUR from the micturating cystourethrography. This investigation would be normal in renal TB.

- B

Renal biopsy

Although renal TB may reveal interstitial nephritis and caseating granulomata in the renal biopsy, there are easier and non-invasive methods of diagnosing this from urine cultures and investigations described above. Renal biopsy may have risks of bleeding, haematuria and, in severe cases if bleeding cannot be stopped, the patient may need kidney removal. Therefore a renal biopsy should not be carried out unless absolutely indicated.

- C

Plain abdominal radiograph

A plain abdominal radiography may show calcified lymph nodes from tuberculosis; however, it would not provide a diagnosis. Calcified lymph nodes may be idiopathic and be due to causes other than TB.

- D

Urine electrophoresis

The urine electrophoresis would be normal in TB. Electrophoresis would be helpful in diagnosis of myeloma where there may be light chain excretion and Bence Jones proteins.

70797

Rate this question:

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	99
Responses Total:	99
Responses - % Correct:	0%

Back to Filters

Question 200 of 203

A 65-year-old man presents with a 6-week history of anorexia, malaise and breathlessness which he relates to rapidly worsening asthma. He takes NSAIDs for osteoarthritis of his knees, paroxetine and also inhalers for mild asthma. He was on lithium carbonate 2–5 years ago. Examination reveals mild peripheral oedema, bilateral pleural effusions and a skin rash. He is hypertensive at 210/100 mmHg.

Investigations:

Hb	7.0 g/dl
Eosinophils	2.2 × 10 ⁹ /l
Serum IgG	8.0 g/l (6.0–13.0 g/l)
Corrected calcium	1.9 mmol/l
Serum IgA	1.2 g/l (0.8–3.0 g/l)
Phosphate	2.5 mmol/l serum
IgM	1.5 g/l (0.4–2.5 g/l)
Urea	79.0 mmol/l
Serum IgE	200 kU/l (<120 kU/l)
Creatinine	800 mol/l
Potassium	7.1 mmol/l
Urinalysis	Blood ++, protein ++
Abdominal ultrasound scan (USS)	No obstruction, kidneys 11 and 12 cm

The likely diagnosis is:

- A

Amyloidosis
- B

Chronic analgesic nephropathy
- C

Churg–Strauss syndrome
- D

Lithium nephrotoxicity
- E

Membranous glomerulonephritis

70798

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 200 of 203

Eosinophils	2.2 × 10 ⁹ /l
Serum IgG	8.0 g/l (6.0–13.0 g/l)
Corrected calcium	1.9 mmol/l
Serum IgA	1.2 g/l (0.8–3.0 g/l)
Phosphate	2.5 mmol/l serum
IgM	1.5 g/l (0.4–2.5 g/l)
Urea	79.0 mmol/l
Serum IgE	200 kU/l (<120 kU/l)
Creatinine	800 μmol/l
Potassium	7.1 mmol/l
Urinalysis	Blood ++, protein ++
Abdominal ultrasound scan (USS)	No obstruction, kidneys 11 and 12 cm

The likely diagnosis is:

- | | |
|---|-------------------------------|
| A | Amyloidosis |
| B | Chronic analgesic nephropathy |
| C | Churg–Strauss syndrome |
| D | Lithium nephrotoxicity |
| E | Membranous glomerulonephritis |

Explanation

- | | |
|---|------------------------|
| C | Churg-Strauss syndrome |
|---|------------------------|

The key to the diagnosis here is the significantly raised eosinophil count and raised serum IgE both of which are characteristic of Churg-Strauss syndrome. Churg-Strauss syndrome is characterised by asthma and blood eosinophilia together with an eosinophilic vasculitis. The initial phase of the disorder is one of asthma and allergic rhinitis, often followed by peripheral blood eosinophilia with eosinophilic tissue disease. The vasculitic phase that follows is life threatening; however, it can often be treated effectively with immunosuppression. Renal disease consists of focal glomerulonephritis in the vasculitic phase with non-selective proteinuria, haematuria and red cell casts on microscopy.

- | | |
|---|-------------|
| A | Amyloidosis |
|---|-------------|

nephrotic picture. Presentation involves heavy proteinuria in the form of nephrotic syndrome and chronic renal failure. There is fatigue, weight loss, bruising and easy bleeding (including the gastrointestinal tract). AL amyloidosis may affect other organs such as heart to produce cardiomegaly and liver with

- B Chronic analgesic nephropathy

Chronic analgesic nephropathy is incorrect. Analgesic nephropathy will not present with blood and protein in the urine and there is no skin rash. There may be papillary necrosis with visible blood in the urine and abdominal pain or insidious renal impairment with no blood or protein in the urine.

- | | |
|---|------------------------|
| D | Lithium nephrotoxicity |
|---|------------------------|

associated with nephrogenic diabetes insipidus and may be associated with tubule-interstitial fibrosis and tubular dilatation. The presentation would be gradual worsening of renal function over a period of years. There is no blood in the urine or skin rash.

- | | |
|---|-------------------------------|
| E | Membranous glomerulonephritis |
|---|-------------------------------|

Membranous glomerulonephritis is incorrect. Membranous glomerulonephritis is the most common cause of nephrotic syndrome in the adult population in the developed world. There would not be any blood in the urine or any skin rash. Progressive renal impairment with proteinuria is a common presentation. The diagnosis is by renal biopsy which would show glomerular basement membrane (GBM) thickening and 'spikes' of GBM extending around subepithelial deposits often best seen on silver stains. There is loss of podocyte foot processes with a variable degree of acute tubular necrosis and tubule-interstitial fibrosis.

Rate this question:

[Next Question](#)

[Previous Question](#)

Tag Question

Feedback

End Session

Session Progress

Responses Correct:	0
Responses Incorrect:	100
Responses Total:	100
Responses - % Correct:	0%

Back to Filters

Question 201 of 203

A 78-year-old woman was admitted to the Accident & Emergency Department with lethargy and lower limb swelling. On examination the ST2 noticed that her left leg was markedly more swollen, consistent with a deep vein thrombosis (DVT). Her blood pressure was 140/60 mmHg, chest and abdominal examination was unremarkable.

Investigations show:

Hb	11.0 g/dl
Urea	10.0 mmol/l
Creatinine	90 mol/l
Sodium	135mmol/l
Potassium	4.5 mmol/l
Cholesterol	7.0 mmol/l
Albumin	25 g/l
Urinalysis	Blood none, protein +++
24-hour protein	>3 g
Renal biopsy	Evidence of membranous disease

What is the most direct link between her DVT and membranous nephropathy?

- A

Antithrombin deficiency
- B

Chronic inflammation
- C

Hypoalbuminaemia
- D

Hypercholesterolaemia
- E

Underlying malignancy

70799

Submit

Previous Question

Skip Question

Calculator

Normal Values

Back to Filters

Question 201 of 203

A 78-year-old woman was admitted to the Accident & Emergency Department with lethargy and lower limb swelling. On examination the ST2 noticed that her left leg was markedly more swollen, consistent with a deep vein thrombosis (DVT). Her blood pressure was 140/60 mmHg, chest and abdominal examination was unremarkable.

Investigations show:

Hb	11.0 g/dl
Urea	10.0 mmol/l
Creatinine	90 mol/l
Sodium	135mmol/l
Potassium	4.5 mmol/l
Cholesterol	7.0 mmol/l
Albumin	25 g/l
Urinalysis	Blood none, protein +++
24-hour protein	>3 g
Renal biopsy	Evidence of membranous disease

What is the most direct link between her DVT and membranous nephropathy?

- AAntithrombin deficiency
- BChronic inflammation
- CHypoalbuminaemia
- DHypercholesterolaemia
- EUnderlying malignancy

Explanation

- AAntithrombin deficiency

Nephrotic syndrome is not only associated with albumin loss, but also with loss of a number of other plasma proteins. Relative deficiency of antithrombin is known to occur in nephrotic syndrome due to loss in the urine and this is thought to be one risk factor in increased development of DVT in these patients. Antithrombin deficiency may also be inherited in autosomal dominant fashion and occur in association with trauma, major surgery or use of the oral contraceptive pill.

- BChronic inflammation

Although chronic inflammation may be associated with hypoalbuminaemia, this is not the mechanism for DVT. Studies have shown that chronic inflammation is a result of DVT rather than a cause of the DVT.

- CHypoalbuminaemia

Nephrotic syndrome is associated with loss of albumin however this is not the mechanism by which DVT is formed. It is loss of other factors such as antithrombin that causes the DVT. Low albumin may be found in other conditions such as liver failure or malnutrition.

- DHypercholesterolaemia

A high cholesterol is found in nephrotic syndrome because there is both an increased in synthesis and decreased clearance of lipoproteinemia. There is an increased level of total and low-density lipoprotein (LDL) cholesterol as the most characteristic abnormality. This does not relate to DVT in any way.

- EUnderlying malignancy

Malignancies are frequently associated with membranous nephropathy and having membranous nephropathy may pre-date malignancy by 10 years. Malignancies may also lead to hyperviscosity and immobility that could lead to an increased risk of DVT. In this case there is no indication that there may be malignancy - there is no history such as weight loss, loss of appetite or change in bowel habit. The haemoglobin is normal.

70799

Rate this question: 1 2 3 4 5

Next Question

Previous Question	Tag Question
Feedback	End Session

Difficulty: Easy

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	101
Responses Total:	101
Responses - % Correct:	0%

Back to Filters

Question 202 of 203

A 28-year-old woman presents to the clinic with microscopic haematuria and proteinuria. Other past history of note is that she has suffered progressive bilateral hearing loss which was first noticed during a school medical in her early teenage years. She has also noticed that her vision has deteriorated over the past few years and she has gone through three pairs of glasses in the past 5 years. On examination she is a slim looking woman with a BMI of 21 and blood pressure of 155/90 mmHg.

Investigations reveal:

K ⁺	5.4 mmol/l
Na ⁺	138 mmol/l
Creatinine	195 mol/l
Urea	7.1mmol/l
Renal ultrasound	Bilateral small kidneys

Which of the following diagnoses fits best with this clinical picture?

- A

Alport’s syndrome
- B

Hypertensive renal disease
- C

IgA nephropathy
- D

Reflux nephropathy
- E

Wegener’s granulomatosis

70800

Submit

Previous Question

Skip Question

Calculator

Normal Values

Question 202 of 203

--	--

ture

-

- nting

2

- ake

ely.

- the

stick

- ct

a
in

- 44

disc

Rate this question:

10

Tag Question

End Session

1

0

102

0.70

Back to Filters

Question 203 of 203

A 23-year-old adopted Jewish man is referred to the Renal Clinic by his GP because of proteinuria. The GP also reports that he has suffered intermittent illnesses in the past, characterised by severe fevers which were thought to be due to recurrent viral illnesses. Other past history of note includes an appendicectomy, and arthritis affecting the hips and knees. On physical examination, blood pressure is 130/80 mmHg, chest and abdominal examination was unremarkable. ESR checked by the GP during one of the recent attacks of fever was raised at 65 mm/h.

Investigations on the day of clinic visit revealed:

K ⁺	5.4 mmol/l
Na ⁺	138 mmol/l
Creatinine	80 mol/l
Dipstick urine	Protein 3+
24-hour urine testing	>3 g protein

Which of the following diagnoses fits best with this clinical picture?

- A

AL amyloidosis
- B

Familial Mediterranean fever
- C

Minimal change disease
- D

Still’s disease
- E

Systemic lupus erythematosus (SLE)

70801

Submit

Previous Question

Calculator

Normal Values

Back to Filters

Question 203 of 203

A 23-year-old adopted Jewish man is referred to the Renal Clinic by his GP because of proteinuria. The GP also reports that he has suffered intermittent illnesses in the past, characterised by severe fevers which were thought to be due to recurrent viral illnesses. Other past history of note includes an appendicectomy, and arthritis affecting the hips and knees. On physical examination, blood pressure is 130/80 mmHg, chest and abdominal examination was unremarkable. ESR checked by the GP during one of the recent attacks of fever was raised at 65 mm/h.

Investigations on the day of clinic visit revealed:

K ⁺	5.4 mmol/l
Na ⁺	138 mmol/l
Creatinine	80 μmol/l
Dipstick urine	Protein 3+
24-hour urine testing	>3 g protein

Which of the following diagnoses fits best with this clinical picture?

- A

AL amyloidosis
- B

Familial Mediterranean fever
- C

Minimal change disease
- D

Still's disease
- E

Systemic lupus erythematosus (SLE)

Explanation

- B

Familial Mediterranean fever

The answer is familial Mediterranean fever (FMF). FMF is described in a number of families, particularly those from the region around the Mediterranean Sea of Jewish or Arab descent. The disease is characterised by intermittent paroxysms of fever, chronic arthritis, appendicectomy related to paroxysmal abdominal pain and nephrotic syndrome due to renal amyloidosis. Pleuritic involvement and pericarditis may also occur in around 25% of patients. Mortality is related to deteriorating renal function. Colchicine therapy has been shown in a number of studies to delay deterioration in renal function and is the mainstay of therapy. A large number of patients do still progress and require renal transplantation.

- A

AL amyloidosis

AL amyloid is due to mononclonal light chains usually lambda or light chain fragments produced by plasma cell dyscrasia. It is more common in women who are over 50 years old. Although nephrotic range proteinuria is a usual presentation, there may be other signs such as easy bleeding, cardiomegaly and hepatomegaly. The young age and high fevers are unusual in AL amyloidosis.

- C

Minimal change disease

Minimal change disease is the most common cause of nephrotic syndrome in the young, therefore the age and nephrotic range proteinuria would fit with this diagnosis. However, there are no associated symptoms of arthritis or recurrent fevers with minimal change disease.

- D

Still's disease

Still's disease is a rare systemic inflammatory disease that is characterised by a triad of persistent high spiking fevers, joint pains and a distinctive salmon-coloured raised rash. Nephrotic range proteinuria is not a feature. In this case, the lack of a rash would also be against the diagnosis of Still's disease.

- E

Systemic lupus erythematosus (SLE)

SLE may present with rash, fatigue and arthralgia. It is an autoimmune condition that mostly occurs in women of childbearing age. The condition is less common in people of European origin and more common in people of African, Caribbean or Asian origin. The lack of clinical symptoms of SLE and the patient being a Jewish man makes the diagnosis of SLE unlikely.

70801

Rate this question: ⚙️ ⭐ ⭐ ⭐ ⭐ ⭐

End Session

- Previous Question

Tag Question
- Feedback

Difficulty: Average

Peer Responses %

Session Progress

Responses Correct:	0
Responses Incorrect:	103
Responses Total:	103
Responses - % Correct:	0%